

E. H. RAINWATER, Tenneco Oil Co., Houston, Tex.
REGIONAL STRATIGRAPHY AND PETROLEUM POTENTIAL
OF GULF COAST LOWER CRETACEOUS

Lower Cretaceous sediments were deposited throughout the Gulf Coast during the Neocomian, Aptian, Albian, and early Cenomanian Epochs. Basal sandstones extend across the entire region, overlapping Upper Jurassic terrigenous strata. Following the initial stage of detrital rock deposition, a shallow epicontinental sea covered the western coastal plain and regions on the south and west. In this area, and also in the seaward parts of the eastern coastal plain, shallow-shelf carbonates were deposited contemporaneously with subsidence. In East Texas and adjacent areas, periods of peneplanation of the bordering land and deposition of carbonates alternated with regressive periods when the hinterland was uplifted and deposition of the land-derived material exceeded subsidence. The southern Appalachians rose at irregular rates throughout Early Cretaceous time, and furnished sediments (sand and clay) to the Mississippi embayment and the eastern Gulf Coast.

Lower Cretaceous rocks are the most widespread and have the greatest volume of any major Gulf Coast stratigraphic division. They are believed to underlie an area of about 340,000 sq mi and have a volume of more than 200,000 cu mi. The area presently productive of oil and gas, extending from Mexico to southwestern Alabama, has an area of 83,000 sq mi, and a volume of 60,000 cu mi. In this proved belt, 1½ billion bbl of oil and 10½ Tcf of gas have been produced from Lower Cretaceous sandstone and carbonate rock. Landward from the productive belt in Texas and Arkansas is a narrow belt that is considered to be non-prospective. This belt widens eastward across central Mississippi, southern Alabama, Georgia, and northern Florida, where continental "redbeds" with thicknesses to 4,000 ft are present. A prospective belt and a speculative belt are gulfward from the proved area.

Depositional conditions of the extensive and thick Lower Cretaceous sediments were favorable for the development and preservation of vast amounts of hydrocarbon source materials and for the formation of many reservoir rocks and stratigraphic-structural traps. An environmental analysis of each stratigraphic unit indicates a very large petroleum potential for this group of rocks. Many hydrocarbon accumulations will be found in the prospective belt, on land as well as under the continental shelf of Mississippi, Alabama, and Florida. The undiscovered accumulations are in deltaic sandstone, carbonate reefs, and shell mounds. In the speculative belt of coastal and offshore Louisiana and Texas, the objectives are limestone reefs which developed on the landward side of positive blocks. Many new fields will be found in the productive belt, and there will be lateral and deeper extensions of producing fields.

E. H. RAINWATER, Tenneco Oil Co., Houston, Tex.
STRATIGRAPHY AND PETROLEUM POTENTIAL OF PENINSULAR FLORIDA AND SOUTHERN GEORGIA

Cretaceous and Tertiary sediments, with a maximum thickness of about 8,000 ft in southwestern Georgia and 18,000 ft in the South Florida basin, are present under all of the area south of the Appalachian Piedmont belt of Precambrian (?) crystalline rocks. In addition, flat-lying early Paleozoic sandstone and shale underlie northern Peninsular Florida and southernmost Georgia; Triassic continental deposits, with diabase sills and dikes, are in grabens under parts of the Geor-

gia coastal plain and northern Florida; and Jurassic terrigenous clastics, carbonates, and evaporites probably are present on the western and southern flanks of the Peninsular arch.

Limestone and dolomite comprise most of the Cretaceous and Tertiary section. Anhydrite is abundant in the Lower Cretaceous and Paleocene deposits, and sandstone and shale are present throughout the section in Georgia near the northern edge of the coastal plain. Deposition was in shallow-marine environments on an extensive, slowly subsiding shelf. The southern Appalachians were of low relief and, after Early Cretaceous time, little sediment from this bordering land was transported to the shoreline of the shallow sea. There were, however, several minor marine transgressions and regressions in southern Georgia during the Late Cretaceous and Tertiary, because of changes in the rate of uplift of the bordering land and downwarp of the coastal plain and adjacent marine areas. The regressive deposits, with sands, were deposited slowly, and they were reworked during the succeeding advances of the sea.

Many wells have been drilled in nearly all parts of Peninsular Florida and southern Georgia in search for petroleum. Therefore, data are available for deciphering the depositional and tectonic history of the area and assessing its petroleum potential. Only 4 small oil fields, all in southern Florida, have been discovered.

Lower Cretaceous carbonates and quartzose sandstones of the continental shelf off western Peninsular Florida, and of the northern flank of the South Florida basin have the greatest petroleum potential. Upper Jurassic carbonates and quartzose sandstones, which are probably present in the South Florida basin and the inner part of the Gulf continental shelf, may have many accumulations, and Paleocene carbonates of northwestern Peninsular Florida and adjacent continental shelf are potentially productive.

GRAY S. ROBINSON, Chevron Oil Co., New Orleans, La.

CHANGE OF BATHYMETRIC DISTRIBUTION OF GENUS *Cyclammina*

Disparities between ancient and modern occurrences of *Cyclammina* spp., both in abundance and in the nature of the associated fauna, indicate that the bathymetric distribution of the genus *Cyclammina* in the Gulf of Mexico has changed with time. An examination of Gulf Coast fossil assemblages reveals that this distribution became more restricted during the late Pliocene, normal occurrences in the younger sediments being limited to upper bathyal and deeper zones. In the older section, the distribution extended into the neritic environment. This change was accompanied by a marked decrease in the relative and absolute abundance of *Cyclammina* spp. in the younger sediments.

ANTHONY D. SHINN, Pennzoil United, Houston, Tex.

POSSIBLE FUTURE PETROLEUM PROVINCES OF GULF COAST—UPPER MIOCENE-PLIOCENE

During the Miocene and continuing through the Pliocene, great volumes of sediment were delivered rapidly to an eastward-shifting depocenter in southernmost Louisiana. Sediments were introduced to the basin by rivers and were distributed farther by marine currents gulfward and laterally across persistently broad shelf areas; the amount and grain size of the