petitive sequences of thin superposed depositional systems. Deposition of fluvial and deltaic facies was rapid; deposition of destructional and transgressive facies represents most of the time consumed by deposition. Depositional systems shifted westward through time as the eastern flank of the basin was filled.

BUSCH, DANIEL A., Tulsa, Okla.

GENETIC UNITS AND DELTA PROSPECTING

Deltas generally are formed at river mouths during stillstands of sea level under conditions of either cyclic transgression or regression. Consequently, they are rarely isolated phenomena, but are present in multiples in a predictable fashion. Reservoir facies consist of both continuous and discontinuous, bifurcating channel sandstones that thicken downward at the expense of the underlying prodelta clays.

All the lithologic components of a deltaic complex are related to each other and are collectively referred to as one type of "genetic increment of strata" (GIS). The GIS is a sequence of strata in which each lithologic component is genetically related to all the others. It is defined at the top by a time-lithologic marker bed (such as a thin limestone or bentonite), and at the base by either a time-lithologic marker bed, an unconformity, or a facies change from marine to nonmarine beds. It generally consists of the sum total of all marginal marine sediments deposited during one stillstand stage of a shoreline, or it may be a wedge of sediments deposited during a series of cyclic subsidences or emergences. An isopach map of a GIS clearly shows the bifurcating trends of the individual distributaries and the shape of the delta, regardless of the variable lithology of the channel fills.

A genetic sequence of strata (GSS) consists of two or more GIS's and, when isopached, clearly defines the shelf, hinge line, and less stable part of a depositional basin. An isopach map of the McAlester Formation of the Arkoma basin is a good example of a GSS. The oil-productive Booch Sandstone is a good example of a deltaic complex occurring within a GIS of this GSS. The upper Tonkawa, Endicott, and Red Fork Sandstones of the Anadarko basin are deltaic accumulations within different GIS's.

A hypothetical model serves to establish the criteria for (1) recognizing successive stillstand positions of a shoreline, (2) predicting paleo-drainage courses, (3) predicting positions of a series of deltaic reservoirs, (4) locating isolated channel sandstone reservoirs, and (5) tracing related beach sandstone reservoirs.

COLLINS, STEPHEN E., Independent, Dallas, Tex.
EVOLUTION OF SMACKOVER PLAY, TEXAS TO FLORIDA

The Smackover limestone is the primary exploration target in the updip part of the entire Gulf Coast area, extending from East Texas to Florida. Accordingly, the Smackover provides most of the new reserves being found in this area and most major companies and many independents have virtually stopped Cretaceous exploration to concentrate on the Jurassic reservoirs (Smackover and Cotton Valley).

Smackover exploration started on a major scale during the 1940s in southern Arkansas where relatively shallow (above 9,000 ft) oil was found. Production was predominantly from oolitic limestones on relatively simple anticlinal structures that were located by reflection seismograph. Nearly 20 years later, in the 1960s,

major Smackover exploration began in East Texas. There, a different set of problems confronted explorationists—more radical facies changes, predominantly dolomite reservoirs, gas condensate-sulfur rather than oil pays, abundant inert gases, complex salt intrusions, greater depths (over 12,000 ft), and erratic porosity developments.

Nearly 5 years later, in the late 1960s, Smackover exploration expanded to Mississippi, Alabama, and most recently, to Florida. There, producing rates are higher and multipays of both oil and gas are common in the Cotton Valley and Norphlet sandstones, in addition to the Smackover. Broad limits of the favorable producing facies in the Smackover are fairly well recognized in Mississippi and presently are being extended eastward into Alabama and Florida where recent discoveries by Humble have started "hot" lease plays.

Although local structure and stratigraphic variations are extremely complex, the regional picture of Smackover deposition is viewed as a simple "wedge" of predominantly carbonate rocks rimming the Gulf Coast, and underlain by salt along the downdip area and an old Paleozoic landmass on the north in the updip area. Most Smackover fields have been found in areas of post-Smackover (Buckner anhydrite) "thins," seismic "salt swells," and very low relief Lower Cretaceous structures. Most of the recently discovered fields cover less than 1,200 acres; however, pay sections of more than 100 ft are not uncommon, yielding large reserves over small geographic areas. Consequently, several tests commonly are required to pinpoint the accumulation.

FISHER, W. L., Bur. Economic Geology, Univ. Texas, Austin, Tex.

TERTIARY DEPOSITIONAL SYSTEMS, GULF OF MEXICO BASIN

Thick, offlapping, terrigenous clastic wedges make up the principal fill of the Gulf basin. Proximal parts of these wedges consist of paralic deposits formed either as large-scale, high-constructive delta systems (with related strike systems) or as a series of smaller, high-destructive delta systems. Distal parts accumulated as continental slope deposits associated with salt diapir fields at the terminus of prograded paralic systems.

High-constructive delta systems (e.g., lower Wilcox, Yegua, and Jackson) are comparable in scale and facies to Holocene Mississippi deltas. They were supplied by rivers with large-volume sediment discharge; fluvial facies are concentrated locally along the basin margin. These deltas consist dominantly of fluvial and fluvially influenced deposits, with extensive coal-bearing delta-plain facies, thick progradational delta-front sandstone facies, and very thick, organic-rich, prodelta mud facies. Progradational sandstone facies show either lobate or elongate patterns in plan. Delta systems of this type supported extensive strike-fed systems comparable to strandplain and barrier bar systems of the Holocene northwestern Gulf Coast.

High-destructive delta systems (e.g., upper Wilcox and Frio) are analogous to the Rhone and other Holocene deltas with significant marine modification (chiefly wave action) of fluvially introduced sediments. These deltas were supplied by numerous, relatively small rivers with moderately high sand load; updip fluvial facies persist along the entire basin margin. High-destructive deltas are composed of a series of sandstone bodies with thickness axes roughly parallel with

regional strike. Each of these deltas consists of local progradational sandstone facies (channel and channel-mouth bars) flanked marginally by extensive sandstone units reworked from channel-mouth bars. Associated prodelta mud facies are moderately thick to thin. High-destructive deltas supported local rather than areally extensive strike-fed systems.

Principal oil and gas reservoirs in high-constructive deltas occur in the progradational delta front sandstones with trends controlled by geometry and distribution of these lobate or elongate sandstone bodies. Vertical stacking of sandstone bodies is common, resulting in multipay fields. Trends within these delta systems are discontinuous along strike, as facies between main prograded lobes consist mostly of muds and tight sands. Attendant growth faulting, salt doming, and mud intrusion cause structural traps. In related barrier bar and strandplain systems trends are regionally persistent with stratigraphic traps. Oil and gas trends in high-destructive deltas are defined by local cuspatetrending coastal barrier sands and downdip progradational channel-mouth bars; principal traps are stratigraphic.

Continental slope systems, making up the distal parts of Gulf basin terrigenous wedges, have been penetrated only in younger units of the basin or in very deep wells. Systems are comparable in scale, composition, and structural association to modern continental slope deposits of the northwestern Gulf. Thick and rapidly deposited delta systems of the Gulf basin mobilized underlying deep-seated salt. Principal flowage was toward the distal front of the prograding systems resulting in distinct diapir fields coextensive with continental slope systems; minor flowage was toward thinner interdeltaic areas. Salt mobilization was a significant control in termination of facies fabric and growth faulting. Younger offlapping units inherited and perpetuated the tectonic grain established by underlying systems.

GALLOWAY, WILLIAM E., Continental Oil Co., Ponca City, Okla.

FACIES OF CISCO-EQUIVALENT SLOPE DEPOSITION SYSTEMS AND THEIR ROLE IN CONSTRUCTION OF EASTERN SHELF

Uppermost Pennsylvanian rocks of the Eastern shelf, a paleogeographic feature of the eastern flank of the Midland basin, consist of 3 depositional systems. (1) The Cisco fluvial-deltaic system extends across the updip part of the shelf and consists of fluvial to deltaic facies flanked by interdeltaic embayment facies. (2) Downdip, along the margin of the shelf, the Sylvester shelf-edge bank system separates relatively flat-lying shelf units from slope deposits. (3) The Sweetwater slope system is composed of numerous broad, coalescing to restricted wedges of sediment 800 to 1,200 ft thick. The terrigenous wedges are bounded by carbonate aprons that extend upslope into the shelf-edge banks. Sandstone distribution within the slope wedges forms fan-shaped maxima that are elongate perpendicular to the shelf edges and grades from relatively narrow, restricted belts upslope to irregular, broad patches downslope. Principal facies include shelf-margin sandstone, which caps the slope wedges, slope-trough sandstone, which extends through the body of the slope wedge, and distal-slope sandstone. Slope-trough facies include thinly interbedded sandstone and mudstone and medium to thick-bedded, graded sandstone units. Sedimentary structures and textures indicate sporadic

deposition, probably by turbidity currents and associated traction carpets, and by subsequent reworking.

The slope system was fed from distal parts of the largest delta distributaries that prograded to the shelf edge, or where tidal or storm currents swept sediment through local breaches in the bank system. Sediment moved downslope by gravity transport mechanisms. The Eastern shelf was constructed by both simultaneous upbuilding (fluvial-deltaic and shelf-edge bank deposition) and outbuilding (slope deposition). Sites of deposition shifted widely across the shelf and slope in response to subtle variations in subsidence and consequent abandonment of active delta lobes.

Petroleum production is limited to lower slope-trough, distal-slope, and shelf-margin sandstones. The small number of fields in the slope system may be the result of (1) low permeability, characteristic of slope sandstones, (2) updip migration of petroleum through slope sandstone facies into facies of other systems, or (3) insufficient drilling.

HALBOUTY, MICHEL T., Consulting geologist and petroleum engineer, Houston, Tex.

ECONOMICS: WITHOUT WHICH—WHAT?

No organization or company with a product to sell can remain in the business of selling that product if the raw materials or parts which make up the product become unavailable. Nor can the company grow if the supply of inventory diminishes in the face of increasing sales ability and demand.

Basic to all such businesses, therefore, is the economics concerned with the availability of inventory, or raw materials. Without raw material for product, there is no business, and without business there can be no need for the economics of running a going concern.

The oil industry, like others, cannot continue to do business long without an adequate and reliable supply of inventory, namely large domestic reserves. Just as great copper reserves in foreign countries are being nationalized by those nations, it is possible that the huge foreign reserves on which the petroleum industry depends for much of its raw crude today may not be available to us someday.

It would be good economics for the domestic petroleum industry to insure its future by giving more consideration to finding domestic reserves. This would ensure a reliable future source of raw materials and would protect this nation from dependence on unreliable crude holdings in foreign countries.

Attainment of raw materials for the future will necessitate the use of economic wisdom in the exploratory search for reserves. For example, (1) exploration money should be used in favorable trends where profitable fields are likely to be found; (2) exploratory tools should be used effectively and efficiently and should supplement, rather than replace, fundamental geology and bold, creative, optimistic geologic thinking; (3) resources should not be placed in jeopardy because of undue risk (expensive drilling programs require partners in many cases); (4) manpower should be used to best advantage, and cooperation between the various disciplines should be stressed; and (5) management should encourage new ideas of exploration, especially those dealing with the search for paleogeomorphic and stratigraphic traps which probably contain the large, undiscovered reserves of the future.

These economic practices, combined with greatly increased domestic exploration and drilling, will satisfy