

beds and salt, representing lower to upper sabkha progradational couplets, reflect dominance of sabkha sedimentation during upper San Andres deposition. The overall genetic aspect of the stratigraphy is a general southerly facies shift through time.

Porosity is best developed in subtidal dolomite facies. Facies mapping delineates areas of potential porosity preservation and is important to explorationists predicting updip San Andres porosity trends and pinch-outs.

ROBERTS, W. H., III, Gulf Research and Development Co., Houston, Tex.

Design and Function of Oil and Gas Traps

Oil and gas are found in traps. If we can understand what is going on in traps, we should be able to look back along the migration trail with special insight as to what has happened. That insight could even extend all the way back to the "source."

Traps are the most logical places for hydrocarbon mixtures to be put together as distinct oil and gas fluids. It follows that traps are not just passive receivers or containers of hydrocarbon mixtures put together elsewhere. Effective oil and gas traps of different well-known styles have a very important feature in common: structurally and stratigraphically, they are designed to discharge waters from depth. Thus they function as active focal mechanisms to gather and process feedstock waters carrying hydrocarbons and other organic materials. It is a forced-draft system. The concept adds an exciting new dimension to the anticlinal theory. It honors all factual observations around oil and gas deposits.

Very simply, the most important function of a trap is to leak water while retaining hydrocarbons. The water can leak because the enclosing membranes and cover are water-soaked, like a wick. The hydrocarbons and other organic materials are separated from the waters as they pass through the trap. The separation is caused by abrupt changes in pressure, temperature, and possibly salinity—those changes being related to the basic change in direction of feedstock (water) movement from lateral to upward. Coalescence of hydrocarbons makes bubbles or globules which cannot move so easily as water. The ultimate composition of a trapped hydrocarbon mixture depends on the respective residence times of the various components of that mixture which in turn depend on (1) what the water carries to the trap, (2) what the trap retains, and (3) the pore-volume exchange rate.

SANNES, TORSTEIN, Saga Petroleum U.S., Inc., Houston, Tex., and E. D. MINIHAN, Marshall R. Young Oil Co., Fort Worth, Tex.

Waveland Field, Unique Structural and Stratigraphic Trap

Waveland field, located in Hancock County, Mississippi, and currently being developed, was discovered in 1965 by Humble Oil and Refining Co. Gas Unit 1, Sec. 22, T8S, R15W. One additional field well was drilled, a northwest diagonal in Sec. 16, T8S, R15W. After these

two field wells were completed and put on production, there was no additional development until 1975, when lease blocks were assembled by Phillips Petroleum Co., Saga Petroleum, Marshall R. Young Oil Co., and others.

There is no massive deposition of Ferry Lake Anhydrite in the area. Because of the lack of massive anhydrite, actual definition of the formations of the Trinity Group (Lower Cretaceous) is difficult to impossible. However, it is interpreted that the primary reservoir of Waveland, a porous limestone, is the Mooringsport Formation.

The depositional environment of the Mooringsport limestone reservoir is extremely complex because of the influence of the large regional carbonate banks on the south. For convenience, the Mooringsport porosity zones are lettered A through G. To date, the A and B zones are the primary contributors to production. The B zone may be subdivided within the field proper.

The reservoir rock is best described as a two-porosity system—matrix porosity (range 6 to 12%) and vugular porosity (range 7 to 16%) directly related to the mineralogy, lithology, and diagenetic history. Matrix permeability is generally low, not exceeding 2 md and usually less than 1 md. Fractures are essential for good productivity and intense fracturing is present in all the higher crestal positions.

The Waveland field comprises the crestal 19,000 productive acres (7,600 ha.) of a north-to-south-elongated nose having no apparent structural closure, fault closure, or north dip at the Mooringsport limestone level. Structural elevation at the Mooringsport level is a major factor in determining hydrocarbon saturation, but no direct relation has been found among pore-throat size, rock type, porosity, structural elevation, and hydrocarbon saturation. The Waveland field is a complex combined structural and stratigraphic trap.

WEBSTER, ROBERT E., Core Laboratories, Inc., Dallas, Tex.

Evidence for Wrench Faulting, Southern Val Verde Basin, Southwest Texas

A structural analysis in the southwest Edwards Plateau area included four 15-minute quadrangles in Edwards, Kinney, and Val Verde Counties, Texas. Seven rock units of the Cretaceous Comanche Series and two units of the Cretaceous Gulf Series crop out. Subsurface Paleozoic foreland facies rocks of the southern Val Verde basin are present in the northern study area; the Devils River uplift, partly blanketed by allochthonous upper Paleozoic metamorphic rocks, underlies the southern part. A pronounced east-west-trending fault zone which cuts through the center of the study area is at least 55 mi (88 km) long and 1 to 2 mi (1.6 to 3.2 km) wide. This trend, named the Carta Valley fault zone, is characterized by a complex series of en echelon, north-east-oriented faults which are cross-cut by a lesser number of northwest-trending faults. These faults bound a series of grabens, downthrown by 200 to 300 ft (60 to 91 m), which are as large as 3 sq mi (7.7 sq km). Nine gently dipping, northwest-plunging, en echelon anticlines parallel the south side of the fault zone. The con-

figuration of faults and folds strongly suggests that the surface fault zone formed by left-lateral wrench displacement in the basement rocks. Subsurface data show that the fault zone overlies a major late Paleozoic high-angle fault, downthrown to the north by over 10,000 ft (3,050 m), which marks the Devils River uplift-Val Verde basin boundary. The surface fault zone may thus be a later reflection of more pronounced wrench faulting along this basement fault during the Ouachita orogeny.

**1980 SYMPOSIUM ON GEOLOGY OF ROCKY MOUNTAIN COAL
APRIL 28-29, 1980
GOLDEN, COLORADO**

A 2-day symposium on the Geology of Rocky Mountain Coal will be held at Green Professional Center, Colorado School of Mines, Golden, Colorado, on April 28 and 29, 1980. The symposium is sponsored by the Energy Minerals Division of AAPG, Colorado Geological Survey, U.S. Geological Survey, Colorado School of Mines, and Colorado School of Mines Research Institute.

This symposium will be followed by a field trip by bus to one of the important Rocky Mountain coal-producing regions (the trip is anticipated to be of approximately 2-days' duration).

Details of the symposium and field trip will be published within a few months in the AAPG *Bulletin*. A registration flyer will be mailed early in 1980.

For additional information, contact: D. KEITH MURRAY, Co-General Chairman, Colorado School of Mines Research Institute, P.O. Box 112, Golden, Colorado 80401. Telephone: (303)279-2581, Ext. 256.

Call for Papers

Authors are invited to submit abstracts of papers dealing with all aspects of coal geology in the Rocky Mountain, Northern Great Plains, Interior, and Pacific Coast Provinces. Topics of interest include stratigraphy and paleoenvironmental modeling of coal and associated strata, geophysical exploration techniques, coal geochemistry, petrology and palynology, determination of the methane content of coal beds, methods of mine planning, and broad evaluations of the potential of individual coal basins or regions. Papers dealing with coal production and utilization are encouraged if they relate such aspects to coal geology.

As a general policy, presentations will be limited to 30 minutes. However, longer presentation time will be allowed for some papers. Please identify abstracts of longer papers at time of submittal.

The deadline for receipt of abstracts, not to exceed 250 words in length, is January 1, 1980. Abstracts should be sent to: THOMAS A. RYER, Program Chairman, U.S. Geological Survey, Box 25046, MS 972, Federal Center, Denver, Colorado 80225.

RECENT CONTRIBUTIONS TO AAPG FOUNDATION

General Fund

John P. Albanese

Thomas W. Angerman
M. E. Arnold, Jr.
Ellwood T. Barrett, II
Charles A. Barton, Sr.
James W. Baxter
B. W. Beebe, in memory of Thomas C. Hiestand
M. W. Bennett
Allan P. Bennison
Allan P. Bennison
Francis X. Bland
A. Edward Blanton
Richard W. Boebel
Douglas L. Bostwick
Feodor Bourgeois
Michael B. Brady
Joseph L. Brown
William F. Brown
R. S. Buchanan
Albert W. Bunsen
Earl P. Burke, Jr.
Leon G. Byerley, Jr.
Fred J. Campen, Jr.
William H. Cardwell
James W. Caylor
George H. Chase
Michael B. Clark
Clarence F. Conrad
John C. Craddock
Ronald Cumella
Wesley Earl Davis
Wesley Edward Davis
Frederik E. Dekker
Leon S. Ditzell, Jr.
Herbert D. Duey
William L. Effinger
Robert T. Elmore, Jr.
Marion S. Escallon
Leroy Esterak
James R. Fasbender
Kenneth J. Fellowes
John D. Ferguson
James L. Fisher
Jack Q. Frizzell
H. J. Funkhouser
James A. Gibbs
John Gilga, Jr.
Thomas Gilmartin
M. W. Goering
Serge Gonzales
Hugh Gerald Graham
Peter S. Gray
Charles Frederic Green
Mark A. Guinan
W. Dow Hamm, in memory of Van D. Robinson
W. Dow Hamm
Margaret H. Hawin
Quey C. Hebrew
Hollis D. Hedberg
Don K. Henderson
G. Warfield Hobbs, IV
W. H. Holman
William F. Howell
William E. Hupp