

continental crust. Throughout this program, particular thought will be given to involvement of a larger scientific community in potential research activities.

KIRKLAND, DOUGLAS W., Mobil Research and Development Corp., and ROBERT EVANS, Mobil Oil Corp., Dallas, TX

#### Source-Rock Potential of Evaporitic Environment

Examination of modern saline lakes, solar ponds, and lagoons shows that the evaporitic environment can be very productive of organic matter. Few species survive in the brines, but those that do survive commonly exist in abundance. In a model evaporitic embayment, the flow of surface currents is persistently toward regions of highest salinity, so that there is a continual supply and concentration of those nutrients being brought into the saline environment. Prolific growth of phytoplankton may occur, analogous to phytoplankton blooms in areas of upwelling in modern oceans. Only carbonates precipitate in the "mesosaline" part (4 to 12% salinity) of the marine evaporitic environment and no great dilution of organic matter by clastic or biogenic sediments occurs. Because of stratification of the brine and the chemically reducing conditions associated with brines, much of the organic matter produced is preserved. The result may be a potentially rich carbonate source rock, frequently unrecognized in the geologic column. In the Middle East, mesosaline conditions have occurred many times from Triassic to Cretaceous and may be responsible for the vast reserves of petroleum in the area. Evaporitic conditions may also have played a part in the petroleum productivity of many other basins, including the Michigan, Paradox, Williston, and Devonian of western Canada.

KITELEY, L. W., and M. E. FIELD, U.S. Geol. Survey, Menlo Park, CA

#### Patterns of Shallow-Marine Deposition, Upper Cretaceous of Northern Colorado

Economically important sandstone bodies encased in marine shales have been described from the Western Interior region of Utah, Colorado, Wyoming, and New Mexico. Our study of a part of an Upper Cretaceous shelf region in northern Colorado shows that the occurrence and distribution of shelf sands were dependent upon a large dynamic sediment system associated with both major and minor transgressive-regressive phases.

Discrete sandstone members of the Mancos and Pierre Shales represent beach, distributary mouth bar, shoreface, and mid-shelf bar deposits. Texture, sorting, bioturbation, thickness, and bed forms of these units are variable, reflecting the variation in rate of deposition and wave energy. Evidence of transgressive reworking is present locally on the caps of some units. Although some sandstone members (e.g., the Hygiene) have been previously correlated over seaward distances exceeding 50 mi (80 km), our detailed examination of cross-bedding patterns and composition indicates deposition in distinctly different shallow-marine environments. Bed forms (medium-scale tabular cross-beds) and lithology of easternmost exposures of the Hygiene are similar to

those in modern sand bars on the United States Atlantic, Bering Sea, and North Sea continental shelves, suggesting deposition at mid-shelf depths. Sources of sands were to the west; dominant direction of transport, however, was to the south.

Key factors in the Upper Cretaceous were the relation between nearshore depositional environments and mechanisms of sediment transport on the shelf. Nearshore sedimentation rates apparently were high but episodic; shelf areas were broad and shallow resulting in discontinuous seaward progradation of delta-front sheet sands; and minor, local transgressive phases, as well as major phases, had impact on the geometry and stratigraphic associations of discrete marine sand bodies during the Late Cretaceous.

KNEBEL, HARLEY J., U.S. Geol. Survey, Woods Hole, MA

#### Processes Controlling Characteristics of Surficial Sand Sheet, U.S. Atlantic Outer Continental Shelf

A surficial sand sheet covers almost the entire Atlantic outer continental shelf of the United States. Until recently, the processes controlling the characteristics of this sheet were inferred mainly from the texture and composition of bottom grab samples and from the bathymetry. Studies of these aspects outlined the general nature and age of major processes, but they were necessarily limited in scope.

With the advent of leasing of the outer shelf for petroleum exploration, many process-oriented studies were initiated by the U.S. Geological Survey. These studies included measurements of the velocity of bottom currents, the frequency of bottom-sediment movement, the kinds and amounts of suspended sediments in near-bottom waters, and the acoustic and sedimentary characteristics of the shallow subbottom strata. These new measurements, when used in conjunction with previous data, show that attributes of the surficial sand sheet such as the thickness, volume, composition, texture, and internal structure have been controlled by a variety of ancient and modern processes. Ancient processes include those associated with glaciers, ancestral rivers, nearshore-marine environments, and subaerial solution and erosion. Modern processes include tidal-, wind-, and wave-driven currents, internal waves, movement of water masses, regional circulation patterns, sediment bioturbation, latitudinal changes in biogenic components, and bottom fishing. A knowledge of the various factors effecting the sand sheet is fundamental to (1) an understanding of its general geologic history; (2) the paleoenvironmental interpretation of ancient sand strata; (3) a determination of the distribution and fate of anthropogenic sediments and pollutants; and (4) an evaluation of potential sand resources and geologic hazards.

KNOX, LARRY W., Tennessee Technological Univ., Cookeville, TN, and MOLLY F. MILLER, Vanderbilt Univ., Nashville, TN

#### Environmental Control of Trace Fossil Morphology Sandstones, shales, and coal of the Fentress Forma-