

panel were observed in data obtained from instruments installed for subsidence monitoring purposes on the surface, in the overburden, and underground. Instrumentation included strain measurement at the surface and both absolute and relative stress measurement at mine level. Data from tape extensometer readings, overcoring, and borehole pressure cells support a rotated fault-block hypothesis. In the mine area, frequent nearly vertical faults are present and can be located on satellite imagery. Two major faults are present and can be located on satellite imagery. Two major faults striking obliquely across the panel appear to have isolated a block of rock that rotated toward the gob at the north end of the panel as mining progressed, creating adverse loading conditions at mine level. The combination of satellite imagery, ground mapping during mining, and selected instrumentation may hold some potential for improved mine development in areas experiencing similar ground control problems due to the interception of geologic discontinuities.

PARKS, JAMES M., and ROBERT A. GALLAGHER, Lehigh Univ., Bethlehem, PA, and EDWARD COTTER, Bucknell Univ., Lewisburg, PA

Tuscarora Sandstone (Silurian), Central Pennsylvania: Preliminary Quantitative Grain Shape Analyses of Cotter's (1982) Facies—Fluvial, Estuarine, Beach, and Marine(?) Shelf

Cotter (1982) demonstrated a logical sequence of depositional facies in the Tuscarora Sandstone, both vertically and laterally, from braided river fluvial deposits to the southeast, progressing through local "estuarine" and regional beach facies, to marine(?) shelf sand waves to the northwest. Several examples of each facies were sampled and subjected to quantitative grain-shape analysis. Although the outcropping Tuscarora has suffered intense diagenesis (pressure solution and quartz overgrowth), these effects are less obvious in thin sections cut parallel to bedding, where the gross two-dimensional grain shapes are relatively unmodified. Preliminary results comparing the Fourier Analysis (FA) methods of Ehrlich and colleagues (1970 to 1982) and Boon (1982) with the Rotated Radials-Factor Score (RR-FS) method of Parks (1981) are presented.

In the RR-FS method, the two-dimensional silhouettes of several hundred grains from each sample are electronically digitized and the raw data (100+ points per grain) are stored on floppy disks. From this point on, the analysis is performed under operator control by a series of FORTRAN programs implemented on a microcomputer. For each grain, the digitized outline is rotated about a calculated "center of gravity" to a least squares best fit with an empirical "reference shape"; and 36 equal-angle spaced radials are calculated. The data set for each sample (400+ grains) is reduced by R-mode Factor Analysis. Computed factor scores are used to classify the gross grain shapes into a limited number of distinguishable categories. Associations of shape types are compared to specific facies.

PATCHEN, DOUGLAS G., MARY C. BEHLING, MICHAEL ED. HOHN, and KATHARINE LEE AVARY, West Virginia Geol. and Econ. Survey, Morgantown, WV

Computer-Generation of a Devonian Shale Production and Potential Atlas for West Virginia

In the final phase of an Eastern Gas Shales Project (EGSP) contract with the U.S. Department of Energy, the West Virginia Geological and Economic Survey compiled an atlas of Devonian shale production and potential. By using both the survey's existing oil and gas data base and a more detailed, computerized data file created during the project, 22 maps were computer-generated for areas of western and southern West Virginia. These multicolored

maps show all wells with known gas production from Devonian shales, all wells drilled to the shales that were dry holes, and all other unsuccessful shale wells that produce either from shallower Mississippian or Pennsylvanian units, or from deeper units below the shales (e.g., the Middle and Lower Devonian Huntersville and Oriskany Formations or the Upper Silurian Newburg sand).

In addition, a gray screen pattern on the maps indicates wells from which shows of gas were reported from the shales, and isopotential lines contour initial open flows from the shale gas wells. Thus, the atlas can be used to locate further shale wells in areas with known high productivity, as well as areas with potential for dual completions in one or more zones in addition to the shale. This atlas is one of many products that can be generated from high quality, detailed computer files.

PRECHT, WILLIAM F., Phillips Petroleum Co., Englewood, CO

Paleoecology and Structure of a Late Silurian-Early Devonian(?) Patch Reef, Northwestern New Jersey

Quantitative and qualitative field examination of an Upper Silurian-Lower Devonian(?) "patch reef" bioherm in northwestern New Jersey resulted in the identification of at least five distinct lithologic and biotic facies: (1) a basinal lithofacies of medium to thick-bedded pelagic limestones containing few discernable fossils; (2) a peloidal grainstone facies, which is dominated by pelmatozoan fragments and peloidal carbonate grains; (3) a reef-talus facies characterized by a high biotic diversity; (4) a coral-boundstone facies dominated by massive tabulate corals; and (5) a stromatoporoid reef facies composed of large juxtaposed hemispherical stromatoporoids (65%), tabulate corals (30%), and rugose corals (5%).

Facies evidence demonstrates that the stromatoporoids and corals organically cemented this buildup into a rigid, wave-resistant structure that controlled its surrounding environment. According to Heckel's 1974 classification of reefs, this type of boundstone lithofacies would indicate an "organic framework reef."

The patch reef grew in a moderate-to-high energy regime and was developed on a carbonate platform which was situated on the margin of an epeiric sea. Reef growth was progradational and upward in response to a relative rise in sea level. Investigation of the patch reef facies revealed compound growth and accretion zones, in addition to demonstrating distributional patterns similar to those of modern patch reef analogs.

REVETTA, FRANK A., Potsdam State Univ. College, Potsdam, NY, DENNIS HODGE, State Univ. New York, Buffalo, NY, and NORBERT O'HARA, Indianantic, FL

Geologic Interpretation of Gravity and Magnetic Anomalies in Western New York and Lake Ontario

The Bouguer gravity anomaly map of western New York State and Lake Ontario indicates a series of gravity anomalies extending north-northeast across New York State and Lake Ontario. These gravity anomalies are due to the extension of the Precambrian rocks of the Bancroft-Madoc area of southern Ontario, Canada. Mafic plutons intruded into a mafic-metavolcanic sequence produce gravity positives, while granitic batholiths cause negative gravity anomalies. The intermediate gravity values are due to a marble-rich metasedimentary group. In most places the gravity positives are located between a magnetic high and low, suggesting the center of the magnetized source is fairly identical with the center of mass surplus.

The Clarendon-Linden fault is the most prominent structural feature in the Paleozoic rocks of western New York. Some control of this Paleozoic structure by the Precambrian basement is suggested

by the location of the structure on the western flank of a series of gravity and magnetic positives. High resolution seismic reflection profiles, gravity, and magnetic-intensity data indicate the structure continues under Lake Ontario into Canada. The geophysical anomalies also suggest the fault has a northwest branch near Attica, NY, which may account for the moderate seismicity between Attica and Buffalo. The mafic intrusions associated with the fault probably indicate fracture zones in the Precambrian basement. The potential of these fracture zones for generating earthquakes in the area is of major concern in nuclear power plant site location.

ROBINSON, JOSEPH E., Syracuse Univ., Syracuse, NY

#### Computer Enhancement of Old Well Logs Suggests New Plays in New York State

There has been oil and gas exploration in New York State since the 1860s, and geophysical well logs have been recorded and deposited in state well information repositories since the 1930s. These logs contain a wealth of information that has largely been overlooked in the recently intensified search for new petroleum and natural gas reserves. The majority of old wells were drilled by small companies using numerous logging contractors. There were a wide variety of logging tools and an astonishing number of both vertical and horizontal scales employed. Often log interpretation was minimal. Regional exploration has been very difficult because of this general lack of well-to-well uniformity, as displayed by the various logs.

The Department of Geology at Syracuse University has compiled a data base of digitized computer processable well logs for a large part of central New York State. These logs have been standardized to a uniform lithology response and uniform scales for optimum correlation, porosity computation, and lithology evaluation. Evaluation of these corrected logs suggests that reservoirs which were either overlooked or considered uneconomic when the wells were drilled could, under today's economic conditions, provide extensive new and potentially profitable gas reserves.

SANTINI, RONALD, J., Amerada Hess Corp., Tulsa, OK, and ALAN H. COOGAN, Kent State Univ., Kent, OH

#### Newburg Porous Carbonate Zone of Lockport Group (Middle Silurian), Northeastern Ohio

The Ohio driller's "Newburg sand" or Newburg zone lies in a stratigraphic position approximately equivalent to part of the Guelph Formation of the Lockport Group (i.e., at the top of the Lockport and just below the Salina A Unit of the Cayuga). The Newburg is characterized by a higher overall porosity than the rest of the Lockport. The porous dolomite extends downward as much as 80 to 100 ft (25 to 30 m) below the top of the Lockport. More than one interval may be porous. Stratigraphically, the base of the Newburg zone parallels the top of the Lockport regardless of whether the Lockport is thick or thin. The thickening of the Lockport occurs locally, especially where there has been a thickening of the lower unit of that group, the Gasport Formation. The thick Newburg zone in the upper Lockport tends to overlie thick Gasport and may reflect reef or reefoid buildup in the Guelph Formation. Mapping of the Lockport surface shows that local structure is caused by the thickening. The small local structures can be detected several hundred feet up-section because of the draping effect of overlying beds on the more rigid Lockport dolomites. Structural mapping of the top of the Lockport or of the top of the Devonian "Big Lime" (Columbus Limestone) may reveal areas of potentially productive thickening of the Newburg zone.

The Newburg usually contains salt water. Accumulation of commercial gas is controlled both by local structural features and by

changes in the relative percent porosity in the Newburg and consequent change in the total porosity feet. Examination of production records from four Newburg gas pools in Summit County, Ohio, shows that the largest and longest yields come from wells located high on structure.

SEYLER, BEVERLY, Illinois State Geol. Survey, Champaign, IL

#### Depositional Environments of Silurian Whirlpool Sandstone (Medina) in Western New York

The Whirlpool Sandstone (Medina Group), quartzose, white, clean, well-rounded, and well-sorted, is the basal Silurian unit in western New York. Its source was the Queenston delta some distance to the southeast. Sedimentary structures indicate that deposition occurred in a beach-shore face environment. Fragmented phosphatic marine fossils are found throughout the sandstone.

Erosion of Upper Ordovician sediments in western New York was minimal and involved only the uppermost green (reduced) portion of the Queenston Shale, which supplied lags of flat, green shale pebbles to the Whirlpool Sandstone. Soft sediment deformation and burrowing in the Queenston substrate (the transitional contact between the red Queenston Shale and red Grimsby Sandstone [Silurian] in Rochester, New York, immediately to the east of the area of Whirlpool deposition) suggest a short period of nondeposition or possibly mild erosion following the deposition of the Queenston Shale. Thus, the Ordovician-Silurian boundary in western New York is essentially conformable and signified only by an abrupt influx of clean quartz sand.

Petrographic analysis shows that the Whirlpool is a second or multicycle sandstone composed primarily of quartz grains originally derived from igneous and metamorphic terranes with only a minor sedimentary component. Because the sandstone is cemented by quartz overgrowths, usually in an advanced stage of development, primary porosity is reduced and the overall porosity of the Whirlpool Sandstone is low (less than 5%). Higher porosities are attributed to the development of secondary moldic porosity caused by dissolution of detrital quartz grains and their original calcite cement. These zones of higher porosity are the most prolific natural gas producing areas in the Whirlpool Sandstone when hydraulic fracturing is used.

SMITH, CARL J., and HOBART M. KING, West Virginia Geol. and Econ. Survey, Morgantown, WV

#### Computer Mapping of Quality Data for Coals in West Virginia—An Aid in Matching a Specific Coal Grade to a Specific End Use

In 1907, the first studies of the state's coal reserves were initiated as part of the West Virginia Geological and Economic Survey's statewide geological mapping program. Since then, extensive work has been conducted to map and characterize the state's 62 minable seams. This effort has shown that the coals exhibit a wide diversity of quality, and this diversity provides the coal user a choice of grades to meet specifications for varied applications. Approximately 6,000 coal samples have been analyzed and a computer data base of coal-quality information is now maintained and continues to grow. An extensive coal-quality mapping project makes this information convenient to use.

The objective of coal-quality mapping is to produce a series of contour maps showing the variations in coal quality for the most important seams. Parameters being mapped include sulfur, ash, Btu, fuel ratio, Hardgrove grindability, volatile matter, fixed carbon, and kilocalories per kilogram. This type of information is extremely valuable for someone interested in buying, selling, evaluating, or developing West Virginia coal.

The maps are computer-generated at a scale of 1:500,000 and