organic matter, whereas for blanket peats, 14% of the section has more than 70% organic matter.

Mineralogic data collected on low-temperature ashes indicate that major minerals in brackish peat consist of kaolinite, quartz, and water-soluble salts, whereas minerals from freshwater samples are primarily kaolinite and quartz. X-ray diffraction and electron microscopy show the presence of smectite, illite, abundant siliceous spicules, framboidal pyrite, biotite, pyroxene, and rutile in the insoluble fraction.

Suitability of these deposits as coal precursors has been discussed by several workers. Superficial examination of data generated so far suggests that only thin coal seams with high ash would be produced. However, the presence of decomposing siliceous sponge spicules in the ash and the possibility of ash being leached by fresh ground water, suggest that much ash may be lost during early diagenesis. Hence, this environment may have more potential as a modern coal-forming model than it first appeared.

LEVIN, DAVID M., Levin Consulting, San Antonio, TX

Deep Wilcox Structure and Stratigraphy in Fandango Field Area, Zapata County, Texas

The Fandango field in Zapata County, Texas, is a new deep Wilcox trend extension. The deep Wilcox sands are commonly found at depths of 15,000 to 20,000 ft (4,500 to 6,100 m). Enough well log and seismic control now exists to make an accurate integrated interpretation of regional deep Wilcox structure and stratigraphy.

Deep Wilcox structure and stratigraphy are controlled by regionally extensive shale anticlines. These shale uplifts control deep Wilcox sand distribution, create large anticlines, and cause regional growth faults which commonly influence local structure. Each regional uplift presents a new exploration frontier holding the promise of vast reserves in the deep Wilcox.

The history of Frio-Vicksburg exploration is an analogy to the deep Wilcox trend today. It took 40 years to expand Frio exploration from shallow stratigraphic traps down into the enormous reserves in the Gulf of Mexico, because each new fault-block extension was considered to mark the downip limit of Frio production. This was, of course, not true and is not true in the deep Wilcox today. The deep Wilcox trend remains virtually unexplored and it is my belief that continued work will prove the existence of much more deep Wilcox potential than is currently thought to exist.

LEVIN, DOUGLAS, Louisiana State Univ. and Louisiana Geol. Survey, Baton Rouge, LA, DAG NUMMEDAL, Louisiana State University, Baton Rouge, LA, and SHEA PENLAND, Louisiana Geol. Survey, Baton Rouge, LA

Tidal Inlet Variability in Mississippi River Delta Plain

Stratigraphic sequences of deltaic and shallow marine origin commonly contain sand bodies transgressively overlying lower delta-plain and delta-front deposits. Although generally ascribed to barriers formed during the destructive phase of the delta cycle, most of this sand is probably of tidal-inlet origin because of the high preservation potential for sediment deposited below the base of the retreating shoreface in deep migratory tidal channels and their associated tidal deltas. To facilitate the identification of such units, this paper reviews the temporal evolution of the inlet sand bodies found along the rapidly transgressive shoreline of the abandoned Holocene Mississippi River deltas. This study also reveals that tide dominance or wave dominance of a coastline is not simply a function of tide range and wave height; it depends largely on the tidal prism, an inlet parameter which, in Louisiana, changes rapidly over time.

Three distinct stages can be identified in the evolutionary sequence for Louisiana tidal inlets: (1) wave-dominated inlets with flood-tidal deltas, (2) tide-dominated inlets with large ebb deltas, and (3) wide, "transitional" inlets with sand bodies confined to the throat section.

Stage 1.—Tidal inlets ranging in age from 50 to a few hundred years are associated with flanking barrier systems attached to erosional deltaic headlands. The barriers enclose restricted interdistributary bays. Small inlets occur at the entrance to abandoned distributary channels within the headland section proper. The tidal prism being exchanged through either of these inlet types is small; the morphology of the inlets and adjacent

coastline is wave-dominated, and most of the inlet sand is associated with a flood-tidal delta. The inlets are generally shallow.

Stage 2.—The Holocene Mississippi River deltas are subject to rapid subsidence and consequent local sea level rise. One gage at Grand Isle indicates a sea level rise of 30 cm (12 in.) over the past 20 years; however, the long-term average is somewhat less. Subsidence leads to an expansion of back-barrier open water environments, an increase in tidal prism, and an evolution of the inlet into a tide-dominated morphology with a deep main channel and large ebb-tidal delta. The recent evolution of Pass Abel and Quatre Bayou Pass represents the transition from wave dominance to tide dominance. Sand bodies developed in stage 2 inlets have the greatest preservation potential because they generally lie below the base of the retreating shoreface.

Stage 3.—Further subsidence generally leads to development of an open sound, permitting efficient tidal exchange with the gulf along the sound margin (Chandeleur Sound). As a consequence, the inlets play only a minor role in the tidal exchange pattern. At this stage, the inlet sand bodies evolve along two distinctly different paths, apparently controlled by sediment supply. Barriers with adequate coarse sediment produce many small well-defined inlets with large flood-tidal deltas (washover fans) and only transient, post-storm ebb deltas. The island shore is distinctly wave dominated. Along coastal segments where coarse sediment is scarce, one finds rapid island deterioration, shoaling of the inlet channel, and reworking of the ebb-tidal deltas into a "transitional" configuration with the sand tied up in throat section shoals.

As the inlets migrate during the transgression, they will leave behind on the continental shelf, tidal sand bodies with a landward succession of facies changing from those characteristic of wave dominance, into tide dominance, and back again to "transitional" or wave-dominated inlets.

LIEBER, ROBERT B., Tesoro Petroleum, Houston, TX, and MARSHALL C. CAROTHERS, El-Can Exploration, Metairie, LA

Harmony Field, Clarke County, Mississippi: A True Stratigraphic Trap

Harmony field in Clarke County, Mississippi, has produced approximately 2 million bbl of oil since its discovery in 1968. Production has been from oolitic grainstones in the upper Jurassic Smackover Formation.

The trapping mechanism at Harmony field is a complex stratigraphic trap. Porous oolitic grainstones pinch out updip into tight carbonates and anhydrite. Structure contour maps on top of the Smackover Formation indicate a low relief structural nose associated with the field. Additional structure maps contoured at the top of the Haynesville anhydrite, approximately 500 ft (150 m) above the top of the Smackover, reveal only regional southwesterly dip. An isopach map of the interval between the two structural markers shows a thinning of Haynesville section coincident with the field area. Evidence suggests, therefore, that the porous Smackover in Harmony field was deposited with depositional relief above the surrounding sediments. This relief had been completely masked by the time the Haynesville anhydrite was deposited.

Stratigraphic and structural cross sections using the Haynesville anhydrite as datum indicate the Smackover in Harmony field consists of not one, but multiple, thin oolitic zones which are productive in various portions of the field. These zones grade laterally as well as updip into nonporous anhydritic carbonates.

The Smackover Formation is often considered to be a chronolithologic unit. In the Harmony field area it is a lithostratigraphic unit, i.e., a unit defined not by time but by a particular rock type, in this case a porous limestone.

During the Late Jurassic, the Harmony field area was near the updip limits of the Mississippi salt basin. Porous oolitic grainstones are interpreted as washovers or storm deposits into lagoons containing tight carbonate muds. Such a model can provide excellent exploration targets in similar areas within the Mississippi salt basin. As Harmony field demonstrates, we are looking for features with little structural expression and great variations in rock type—true stratigraphic traps.

LOCK, BRIAN E., Univ. Southwestern Louisiana, Lafayette, LA, BRUCE K. DARLING, Pennzoil Co., Midland, TX, and ILENE D. REX, Univ. Southwestern Louisiana, Lafayette, LA

Marginal Marine Evaporites, Lower Cretaceous of Arkansas

The mixed evaporite/carbonate/terrigenous clastic sediments of the DeQueen formation, in southwestern Arkansas, were deposited at the