

thrust the West Texas crustal block against the Delaware block, with local denudation of the uplifted edge and eastward-directed backthrusting into the Midland basin. Later in the Permian, the area was the center of a subcontinental bowl of subsidence—the Permian basin proper. The disturbed belt formed a pedestal for the carbonate accumulations which created the Central Basin platform.

The major pre-Permian reservoirs of the Permian basin lie in large structural and unconformity-bounded traps on uplift ridges and domes. Further work on the regional structural style may help to predict fracture trends, to assess the timing of oil migration, and to evaluate intrareservoir variations in the overlying Permian giant oil fields.

FALKENHEIN, F. U. H., R. M. BARROS, I. G. DA COSTA, and C. CAINELLI, *Petróleo Brasileiro S/A, Rio De Janeiro, Brazil*

Potiguar Basin: Geologic Model and Habitat of Oil of a Brazilian Equatorial Basin

The Potiguar basin integrates the eastern part of the Brazilian equatorial Atlantic-type margin. The rifting stage of this basin occurred during the Neocomian and Aptian. The drifting stage and sea-floor spreading began in the Late Albian. The rifting stage clearly was intracratonic during the Neocomian and is recognized as a mosaic of half-grabens trending mostly northeast-southwest and filled with syntectonic lacustrine siliciclastics. The half-graben pattern exhibits rotation of beds into the major fault zone, and the preserved uplifted margins display either paleostructures or paleogeomorphic features with hydrocarbons. A regional pre-Aptian unconformity preceded the Aptian proto-oceanic rifting stage which was characterized by syntectonic fluvio-deltaic sediments. The Aptian tectonics were represented by reactivation of former lineaments superimposed by predominant east-west normal faulting. Structural highs during this stage are so far the most prolific oil accumulations. The most important source beds and reservoir rocks are both Neocomian and Aptian sediments. Geochemistry and hydrodynamics have shown that hydrocarbon migration was driven through fracture or fault zones in both Aptian or Albian plays. Lithofacies maps support this interpretation because pools occur whenever adjacent downthrown blocks present a high shale content.

FANG, J. H., *Univ. Alabama, University, AL*, and LEVY ZEVIN, *Univ. Ben Gurion, Beer Sheva, Israel*

New Method in X-Ray Quantitative Mineralogical Determination of Sedimentary Rocks

A new method in quantitative x-ray diffractometry has been developed which uses no "standard" minerals for calibration. Thus, this new method is diametrically different from the classical internal method of Klug and Alexander, which requires the similarity (in chemical composition, crystallinity, and degree of preferred orientation) of minerals used as "the standards" and in the unknowns. The method is based on the interrelationship between the weight percentages of constituent minerals in a rock and relative intensities of x-ray peaks arising from these minerals.

If the concentration in percent of mineral m in sample s is denoted as X_{ms} , the mass absorption coefficient as μ_s , the intensity of a diffraction peak as I_{ms} , and the instrumental (including structural and compositional characteristics of the constituent minerals) constant as K_m , then the weight percentages of minerals in a rock can be obtained easily through a least-squares analysis, $X_{ms} = K_s / I'_{ms}$, where $I' = I_{ms} / \mu_s$. The μ_s can be measured readily from the Compton scattering intensity with an x-ray spectrometer.

Examples from a synthetic mixture, carbonate rock, and mudrock illustrate this new approach.

FARRELL, KATHLEEN, M., and DAG NUMMEDAL, *Louisiana State Univ., Baton Rouge, LA*, and SUDHIR B. BASUMALLICK, *India School of Mines, Dhanbad, India*

Sedimentation in a Transgressed Interdistributary Bay on Mississippi-Lafourche Delta Lobe

Caminada Bay is a shallow south Louisiana bay, located between the distributary levees and beach-ridge plain of the late Lafourche Mississippi

delta (900-500 y.B.P.) on the west and the Bayou Des Families levees (3,000-2,000 y.B.P.) on the east. Small distributaries traversed the region of the present-day bay, probably during both early and late Lafourche time. The Lafourche delta actively prograded until about 500 y.B.P. when it was abandoned by the Mississippi River; the area of the present bay then entered its transgressive phase. Grand Isle, which separates Caminada Bay from the Gulf of Mexico, was subsequently built by sand derived from local distributary-mouth bars and erosion of the Lafourche beach-ridge plain to the west. As a consequence of the cutoff in fluvial sediment supply and continued subsidence, the lower delta plain of the Lafourche lobe is currently being transformed into the rapidly expanding Caminada Bay.

Visual description, x-ray radiography, and resin peels of 23 vibracores (4 to 8 m, 13 to 26 ft, in depth) reveal the following vertical succession of sedimentary units beneath the bay: (1) basal paleobay deposits, (2) an intermediate unit representing Lafourche delta lobe progradation, and (3) upper transgressive bay deposits.

The deepest unit penetrated (paleobay deposits) consists of mottled or bioturbated very fine sand and clay and shell beds. The intermediate unit (delta lobe) consists of intricately interlaminated very fine to fine sand, silt, and clay. It features lenticular, wavy, and flaser bedding. The upper transgressive unit consists of banded organic-rich clay with in-situ and detrital peat. Expansion of the bay by wave erosion of the many marsh islands removes most of the organic-rich upper unit. A thin unit of bioturbated muds is currently being emplaced on the bay floor.

FEELEY, MARY H., *Texas A&M Univ., College Station, TX*, RICHARD T. BUFFLER, *Univ. Texas at Austin, Austin, TX*, and WILLIAM R. BRYANT, *Texas A&M Univ., College Station, TX*

Seismic Stratigraphic Interpretation of Mississippi Fan, Gulf of Mexico

Examination of extensive multichannel and single-channel seismic data across the Mississippi Fan, Gulf of Mexico, reveals that at least 7 seismic sequences comprise the upper Pliocene-Pleistocene section of this giant fan. These sequences are divided into 2 groups based on continuity and amplitude of reflectors. The lower 3 sequences are generally characterized by high-amplitude, parallel to subparallel, continuous reflectors overlain in places by a hummocky clinoform reflection configuration. The reflection patterns suggest distal turbidites deposited in a relatively low-energy environment. In contrast, the upper 4 sequences are generally thicker and characterized by regionally extensive chaotic units interbedded with thin, high-amplitude, parallel to subparallel reflection packages. The chaotic zones grade laterally into more continuous, parallel to gently diverging reflection patterns, probably a lower energy, more distal turbidite facies. Isopach and structure maps of each sequence indicate a seaward and eastward migration in the Pliocene-Pleistocene depocenter during fan development.

Channel, levee, slump, turbidite, and hemipelagic deposits are interpreted within each sequence. Channel/levee deposits are extensive, showing great variability in their morphology and distribution. On the upper fan, the channels are large with well-developed levee sequences. On the middle and lower fan, the channel sequences are smaller, confined mainly to the apex of lobes, and show little evidence of migration and abandonment during lobe development. Slumping off the slope apparently contributed a significant percentage of the material deposited on the upper fan. In addition, the truncation of prominent reflectors and the mounded, chaotic and diffracted patterns on the middle and lower fan suggest that slumping was continually active during fan development.

FELDMAN, SANDRA C., *Univ. Nevada, Reno, NV*

Appraisal of Coal Resources from Uranium Drill-Hole Logs, Southern San Juan Basin, New Mexico

Geophysical logs from uranium drill holes in the Grants region are a valuable source of information on coal resources. Coal occurs in the southern San Juan basin of New Mexico in the Upper Cretaceous Gallup Sandstone, Crevasse Canyon Formation, and Menefee Formation. Uranium has been mined from the Upper Jurassic Morrison Formation that underlies the coal-bearing Cretaceous formations and is separated from them by approximately 1,000 ft (300 m) of section.

Permission was obtained from Santa Fe Mining, Inc., Pathfinder