

episodic concept it follows that the best way to determine the distribution of sedimentary rocks within a basin is to understand facies relationships and the tectonic setting of the basin. This concept makes good sense and obviously applies to the vast majority of sediments. The weakness of the concept is its inability to explain the "rare event." For these rare events, the episodic-oriented geologist commonly calls on the 1,000 yr storm, the 500-ft (150-m) waves from meteorites, the blanket of dust that extinguishes life.

Our experience, based on seismic stratigraphic studies tied to well and outcrop sections, indicates that yes, sediments are deposited episodically, but they are packaged in genetically related depositional packages or sequences that are shifted back and forth in a predictable global cyclic pattern. We believe this global cyclic pattern is caused by rapidly fluctuating eustatic changes of sea level superimposed on more slowly changing tectonics. Each sequence is composed of all the rocks deposited during a complete cycle of sea level starting with the fall and progressing through the succeeding low, rise, and high before the next fall. We believe orderly cyclic sedimentation caused by eustatic sea level changes is a better explanation for many of the rare events. Deep-marine massive sand fans and debris flows commonly ascribed to 1,000 yr storms or 500-ft (150-m) tidal waves may be explained better by rapidly falling sea level or sea level lows. Rapid rises of sea level and their associated condensed stratigraphic sections offer an alternative explanation for the massive faunal extinction and rare deposits associated with the Cretaceous-Tertiary or Eocene-Oligocene boundaries.

WANLESS, HAROLD R., Univ. Miami, RSMAS, Virginia Key, FL and JEFFREY J. DRAVIS, Exxon Production Research Co., Houston, TX

#### Comparison of Two Holocene Tidal Flats—Andros Island, Bahamas, and Caicos, British West Indies

Minor climatic and physiographic differences have caused dramatic differences in sedimentation between the tidal flats of Andros, northwestern Bahamas, and Caicos in the southeastern Bahama chain.

Both tidal flats are leeward of large islands and adjacent to broad, shallow platforms that provide carbonate mud to the flats. Each flat forms a sediment wedge 4 m (13 ft) thick (Andros flats are 200 × 30 km, 125 × 19 mi; Caicos, 60 × 10 km, 37 × 6 mi), and each contains an outer channeled flat and an inner algal marsh. Both flats have cementation and protodolomite forming on channel-margin levees and the inner algal marsh.

Differences in rainfall, wind regime and orientation promote important sedimentological differences. Andros receives nearly twice as much rainfall as Caicos; gypsum forms in cemented crusts in Caicos, but not on Andros. Andros receives brief strong pulses of northwest wind following passage of 40 to 60 winter cold fronts per year. The northwest exposure of the margin of the Andros flat thus is flooded several times each year by sediment-laden waters. This flooding focuses on the shore and channel margins, building broad, strongly laminated levees. Caicos flats, in contrast, face south, and neither winter cold fronts nor summer trade winds blow onshore. Levees are poorly developed. Caicos is dominated by brisk easterly trade winds, causing persistent turbidity in nearshore waters. This is dispersed through tidal channels, largely filling intertributary ponds to a level at which organic-rich intertidal algal mats can flourish. Lower energy conditions on Andros provide insufficient sediment to fill ponds.

WARESBACK, DAMON B., JOHN G. MCPHERSON, and STEPHEN SELF, Univ. Texas, Arlington, TX

#### Volcanogenic Alluvial Fan Sedimentation, Puye Formation, New Mexico

The Pliocene Puye Formation of north-central New Mexico represents a rift-filling volcanoclastic alluvial fan sequence developed on the east flank of the Jemez Mountain volcanic pile. This coarse-grained sequence can be separated into two compositionally distinct members that are genetically unrelated. The basal Puye (here referred to as the Totavi Member) consists of a 25-m (82-ft) thick, clast-supported conglomerate with interbedded sandstone lenses. The Totavi is a coarse-grained braided stream deposit composed of reworked Precambrian terrane of the Sangre de Cristo Mountains and mafic lavas of the Taos basin. The sequence displays cross-bedded sandstones and conglomerates, and well-developed clast imbrication and long-axis orientation, which indicate a paleoflow

direction to the southwest. The Totavi member is an axial-stream gravel deposit of the Rio Grande rift.

The Totavi member is conformably overlain by a 100+ m (330+ ft) thick volcanogenic alluvial fan deposit here referred to as the San Ildefonso Member. This member represents the eroded and reworked detritus of the growing silicic volcanic complex of the Jemez Mountains. The San Ildefonso Member consists of four major interstratified lithofacies associations. These include: (1) clast-supported, massive and horizontally stratified conglomerates, with interbedded sandstones; (2) matrix-supported conglomerates; (3) laminated claystones and mudstones; and (4) primary pyroclastic deposits of both silicic and mafic composition. The clast-supported conglomerates and interbedded sandstones are typical coarse-grained, braided-stream deposits representing longitudinal bar and minor transverse bar accumulations. Internal stratification and clast imbrication points to an easterly flow direction.

The matrix-supported units display a complete spectrum of types from clast-rich (debris flows) to clast-poor (mudflow) varieties. These deposits are sediment gravity flows developed in response to steep slopes, abundant unconsolidated volcanic detritus, and excessive loading by water uptake. A number of debris flows and mudflows are underlain by primary volcanic airfall material and were undoubtedly initiated by volcanism.

The laminated claystones and mudstones are lacustrine deposits that resulted from temporary damming of the alluvial fan and ancestral Rio Grande drainage systems. The San Ildefonso Member contains at least 7 primary silicic air fall ash beds and 3 basaltic ash beds. These not only provide a time framework for sedimentation but can be used to delineate lateral facies changes.

WARREN, JOHN, Univ. Texas, Austin, TX, DAVID LOCK, CRA Exploration, Pty. Ltd., Research Corp., Canberra, Australia, and PATTI GRANGER, Univ. Texas, Austin, TX

#### Coorong Dolomites (South Australia) and Models of Ancient Coorong-Style Dolomites

The Coorong Dolomites and associated magnesian and calcian carbonate muds form sporadic deposits over a 90 × 200 km (56 × 125 mi) area of coastal dune strand plain in southeastern South Australia. They are deposited wherever the Pleistocene dune surface intersects the water table. A reconstruction based on the hydrological setting of the Coorong is a facies mosaic model. It includes the ideas that: (a) dolomite and magnesian carbonate deposition was widespread and associated with resurging continental groundwater; and (b) although widespread, the actual volume of "primary" dolomite is insignificant compared to the total volume of lacustrine and other carbonate sediment.

The lacustrine sediment in the interdunal depressions builds up to the highest level attained by the water surface, equivalent to the outcropping water table. The upper portion of these carbonate muds often contains one, sometimes two, indurated crusts. Such crusts contain subtle, small-scale structures called extrusion tepees. These tepees form as layers of mud are injected from below into megapolygonal cracks in the crust. Successive layers of mud are cemented to the sides of the crack as the lake desiccates each summer. Ongoing episodes of injection and cementation expand the crust volume until it overthrusts into large-scale tepee structures.

An ancient "Coorong-type" dolomite is a shallowing upward mudstone sequence which shows increasing evidence of subaerial exposure in the upper levels of the cycle. The lower portion of the cycle is a relatively thick, organic rich, laminated, calcitic mudstone (2–3 m, 7–10 ft, thick). This passes up into a lighter, highly bioturbated and pelleted dolomudstone (1–2 m, 3–7 ft, thick). The upper zone is characterized by crusts containing extrusion tepees, intraclast breccias, and siliceous cements. The succession is capped by a well-developed palaeosol with portions of the uppermost zones sometimes showing evidence of dedolomitization.

One should be wary of the idea that ancient Coorong-type dolomites are never associated with evaporites. In the modern Coorong lakes, gypsum and halite grow as an ephemeral phase in the near surface and surface muds during late summer. The same crystals are dissolved out by winter rains, and their molds destroyed by bioturbation and the thixotropic nature of fresh dolomite mud. In Precambrian analogs, the lack of bioturbation could lead to the retention of some evaporite pseudomorphs. Also the modern hydrology of the Coorong is highly compartmentalized. A laminated gypsarenite sequence at least 2 m (7 ft) thick is

forming in a coastal salina (Halite Lake) less than 2 km (1 mi) from the area where the thickest Coorong dolomites are forming. Obviously evaporite filled ponds can occur in an ancient Coorong-type facies mosaic, and the distribution of evaporites versus dolomites could be a good palaeoslope indicator.

WASHINGTON, PAUL A., Univ. Connecticut, Storrs, CT

#### Variability of Internal Geometry of Duplexes

Duplexes are imbrications of the footwall of a thrust fault (or series of connected thrust faults) in which the minor faults connect to or intersect with the master thrust surface at both their leading and trailing edges. Within this closely constrained definition, there is considerable geometric variation. The current classification scheme recognizes the variation in postdeformational arrangement of horses in two dimensions, but ignores variations in the size and shape of the horses and in the displacement on the imbricate fault surfaces. Compilation of the geometric arrangement of horses in known duplexes shows that such variations are common and result in several geometric variations that do not fit into the current classification. Additional geometric variations can be found in 3-dimensional framework.

These geometric variations can be correlated with mechanical properties of the footwall material in which the duplex formed. Since the deformational style within the horses is also controlled by the same mechanical factors, the internal structure of the horses can be correlated with the gross geometry of the duplex. Recognition of the range of duplex geometries and the correlation with deformational patterns should aid exploration in complex thrust belts.

WATKINS, DAVID K., Florida State Univ., Tallahassee, FL

#### Calcareous Nannofossil Paleobiogeography of the Cretaceous Greenhorn Sea

Two distinctive, laterally traceable bentonite beds were used to construct two isochronous time slices through the marine sediments of the Upper Cretaceous Greenhorn cyclothem of the United States Western Interior basin. Calcareous nannofossil assemblages from these time slices were examined from more than 40 outcrop localities. Nannofossil presence and assemblage diversity and composition were statistically analyzed to examine the paleoceanographic conditions within the basin.

The lower time slice (X bentonite) is at the stratigraphic horizon which approximately corresponds to the time at which free communication between the basin and open oceanic systems first occurred. The most striking trend in the nannofossil distribution is delineated by the presence or absence of nannofossils. The presence of common to abundant nannofossils in the center of the basin (i.e., near the hingeline of the basin) and the absence of nannofossils from the eastern and western basin margins indicate that open marine conditions conducive to significant standing crops of calcareous phytoplankton occurred only in a narrow, centrally located zone. Conditions at the basin margins were probably unsuitable for large populations due to environmental instability. Evidence indicates that *Watznaueria barnasae* was the most ecologically tolerant form.

The upper time slice (HL-3 bentonite) samples the nannofossil distributions at a point at or near the maximum transgression of the Greenhorn Sea. Similar, basinward increases in diversity and abundance are evident but more subtly expressed. Paleolatitudinal differences in diversity and the relative abundances of some taxa indicate contributions from both tropical and boreal oceanic water masses.

WATSON, MICHAEL P., BP Petroleum Development Ltd., Dyce, Aberdeen, Scotland

#### Submarine Fans in a Developing Extensional Regime—Their Significance in the North Sea Hydrocarbon Province

The North Sea is a major hydrocarbon province with estimated proven recoverable reserves of approximately 23 billion bbl of oil and 50 tcf of gas. A significant proportion (22%) of these reserves occurs in reservoirs interpreted as submarine fan deposits. These include both oil and gas fields that are among the largest discovered in the province.

The North Sea basin provides a classic example of long-term development of an extensional basin. Major tectono-sedimentary sequences can be broadly matched to pre-, syn- and post-rift phases of the Viking and Central grabens. Development of submarine fan systems within the basin can in turn be related to this framework.

Submarine fans developed in the Viking graben within the syn-rift tectonic setting during the late Jurassic. The major controlling factor at this time is considered to be internal tectonics of the graben with subordinate influence from sea level changes. Small, coarse, scarp-fed fans formed a sediment apron along the fault-controlled graben margins. Finer-grained, rather better developed fan systems were also formed during this period, reflecting the relative importance of different controlling influences.

The post-rift cycle in the North Sea basin is characterized by Late Cretaceous-Tertiary thermal subsidence of the graben as well as the former rift margins. The major influences on fan development in this phase were external tectonics in the source area, and, of equal importance, sea level changes which controlled sediment input to deeper parts of the basin. Late Paleocene and early Eocene fans developed in association with major prograding delta systems.

The variability in space and time of submarine fan types in the North Sea highlights the importance of identifying the most appropriate fan model. This is critical to reservoir distribution in field development, as well as in future exploration where predictive models will be required to identify time periods and areas most likely for the development of submarine fans.

WEDOW, HELMUTH, GENE E. TOLBERT, and HOWARD A. PRICE, JR., MAPSS Corp., Oak Ridge, TN

#### Computer-Assisted Attribute-Coincidence Mapping (ACM) in the Search for Massive-Sulfide Exploration Targets in Appalachian Devonian Rocks

Attribute-coincidence mapping (ACM) is a powerful tool commonly used by geologists when they visually compare map data from diverse sources by use of a series of transparent overlays. The same geologists using computer-assisted ACM (particularly the MAPSS=MAGIC software system) can consider, evaluate, and replot or store for later retrieval many more kinds of data with less time and effort than with manual methods. Because the MAPSS=MAGIC system organizes and manages information by orthogonal patterns, standard CRT terminals or high-speed line printers enable users to view complex grid-cell patterns of derivative maps almost instantaneously, and thus facilitate rapid decision making.

ACM with the MAPSS=MAGIC system can be used at any scale to compute, analyze, and plot derivative patterns. For regional analysis the standard U. S. Geol. Survey 7 1/2-minute quadrangle is used as the information cell for data manipulation. A demonstration of the system shows map patterns of some of the various attributes of sediment-associated massive-sulfide ore deposits that have been combined through ACM to identify potential exploration targets in Appalachian Devonian rocks. The massive-sulfide model was developed by the West German Federal Geol. Survey and recently modified by Wedow for computer use by the U. S. Geol. Survey. This ACM study of the Appalachian Devonian has revealed several significant exploration targets.

WEICHMAN, BEN, Beneco, Inc., Houston, TX

#### A View on Oil Shale Development

An examination of energy statistics shows clearly that the United States is on the edge of an impending energy crisis. This appears preposterous in the face of a world oil glut and probable world oil price decreases over the next year or so. However, oil supply, oil demand and United States economy statistics support the scenario of a rapidly occurring energy crisis in the United States should oil imports be shut off even partially. This is not an unlikely happening under present world tensions.

Oil and gas exploration and production statistics support the view that domestic oil reserves will continue to decline even if concentrated oil exploration activity were maintained.

The shortest avenue for United States energy independence is through the development of synthetic fuels. However, a detailed study shows that, under present technology, synthetic fuels cannot be produced economi-