

The informal term "Langtry formation" is applied to these strata, which flank shallower water platform facies to the east, north, and west (Austin Chalk and San Vicente Member of the Boquillas Formation), and pass into basinal limestones of the San Felipe Formation in the La Mula basin to the south.

LOWRIE, ALLEN, RFD Rte. 1, Box 164, Salem Community-E.Z. Goss Road, Picayune, MS

Sediment Distribution About Salt Domes and Ridges on Louisiana Slope

Salt ridges and domes underlie much of the present Louisiana slope. The bathymetric expression of underlying salt could be either a mound or a flattening of the normal rate of descent down the slope. The "mounded" salt features form barriers to the gravity-driven sediments from the shelf break. Much industrial research has been done in the search for reservoir sands about such an obstruction. Clues to depositional patterns about salt features may be obtained from studies in the deep ocean about seamounts located in pathways of ocean-bottom-following currents. Parallel-bedded sediments form foredrifts on the upcurrent side of a seamount. These foredrift sediments were deposited where the prevailing ocean bottom currents were locally decelerated by the obstructing seamount. In waters overlying the obstruction, a Taylor column of dead water or a slow cyclonic eddy provides tranquil oceanographic conditions, permitting a greater fallout of sediments. Moats are found on the sides of the obstruction and are the result of erosion or non-deposition owing to acceleration of deflected waters. Leedrifts are found on the downcurrent side of the obstruction. Current gyres result from deceleration of accelerated currents along the obstruction's flanks, and a complex sedimentation pattern results. Flow over the obstruction's top is determined by size and shape of the obstruction relative to size and velocity of the bottom-following current. A turbulent wave will be set up which may have sufficient amplitude to influence sedimentation on the downcurrent side. There are appreciable differences between the sedimentation patterns about a dome and a ridge.

If ocean bottom currents equal gravity-driven terrigenous sediment movement and seamounts equal salt domes and ridges, then the result of deep ocean surveys are directly applicable to sedimentation on slopes with underlying salt basement. The salt-related sedimentation pattern of the present slope should be applicable to similar paleoenvironments.

MANCINI, ERNEST A., ROBERT M. MINK, and BENNETT L. BEARDEN, Alabama State Oil and Gas Board, Tuscaloosa, AL

Paleoenvironments and Hydrocarbon Potential of Upper Jurassic Norphlet Formation of Southwestern Alabama and Adjacent Coastal Water Area

Upper Jurassic Norphlet sediments in southwestern Alabama and the adjacent coastal water area accumulated under arid climatic conditions. The Appalachian Mountains of the eastern United States extended into southwestern Alabama, providing a barrier for air and water circulation during Norphlet deposition. These mountains not only contributed to the arid climate but also affected sedimentation. Norphlet paleogeography was dominated by a broad desert plain rimmed to the north and east by the Appalachians and to the south by a developing shallow sea. The desert plain extended westward into eastern and central Mississippi.

Initiation of Norphlet sedimentation was a result of erosion of the southern Appalachians. Norphlet conglomerates were deposited in coalescing alluvial fans in proximity to an Appalachian source. The conglomeratic sandstones grade downdip into red-bed lithofacies that accumulated in distal portions of alluvial fan and wadi systems. Quartzose sandstones (Denkman Member) were deposited as dune and interdune sediments on a broad desert plain. The source of the sand was the updip and adjacent alluvial fan, plain, and wadi deposits. Wadi and playa lake sediments probably also accumulated in the interdune areas. A marine transgression was initiated late in Denkman deposition, resulting in the reworking of previously deposited Norphlet sediments.

Norphlet hydrocarbon potential in southwestern and offshore Alabama is excellent with four oil and gas fields already established. Petroleum traps discovered to date are primarily structural traps involving salt anticlines, faulted salt anticlines, and extensional fault traps associated with salt movement. Reservoir rocks consist of quartzose sandstones,

which are principally eolian in origin. Porosity types include both intergranular and secondary dissolution. Smackover algal carbonate mudstones were probably the source for the Norphlet hydrocarbons.

MAZZULLO, JIM, and JEFFREY CRISP, Texas A&M Univ., College Station, TX

Sources and Distribution of Silt, South Texas Shelf

Fourier grain shape and mineralogic analyses were conducted on the coarse silt fraction of the surficial sediments on the south Texas continental shelf to determine the sources and distribution of the silt. The distribution patterns were evaluated in light of the late Pleistocene paleogeography and modern hydrodynamic conditions prevailing on the shelf to determine whether the coarse silt fraction was relict, palimpsest, or modern in origin.

Two coarse silt-grain shape types are present in varying proportions in the samples from the south Texas shelf. One is associated with quartzose coarse silt and is considered to represent multicyclic coarse silt-size detritus derived from the Texas coastal plain and high plains. The second type is associated with more feldspathic coarse silt and is considered to represent first-cycle, coarse, silt-size detritus derived from igneous and metamorphic rocks of Texas, New Mexico, and Mexico.

Each of the major river systems that supplied sediment to the south Texas shelf during the late Pleistocene and Holocene is characterized by distinctive proportions of first- and multicyclic coarse silt. The Brazos-Colorado and Rio Grande systems, which drain igneous, sedimentary, and metamorphic rock terranes, deposited sediment enriched in first-cycle coarse silt on the shelf during the late Pleistocene low sea level stand. At the same time, the Guadalupe and the Copano-Nueces-Baffin coastal plain river systems, which drain sedimentary rock terranes only, deposited sediment enriched in multicyclic coarse silt on the shelf.

In the southern part of the other south Texas shelf, the distribution patterns of first-cycle and multicyclic coarse silt define the late Pleistocene alluvial plains of the Rio Grande and the south Texas coastal plain rivers. The locations of the alluvial valleys, as defined by grain shape analysis, coincide precisely with the locations of these valleys as defined by shallow seismic studies; therefore, the coarse silts in this area are considered to be relict in origin, unmodified by the weak (0-10 cm/sec) semipermanent bottom currents that prevail in the southern part of the outer south Texas shelf. The preservation of paleogeographic features in the surficial sediments of this area of the shelf indicates that no modern coarse silt, which might bury this relict surface, has been deposited here during modern times.

In the northern part of the south Texas outer shelf, first-cycle coarse silt originally deposited on the ancestral Brazos-Colorado delta is found not only in the delta area, but also overlying most of the alluvial valley of the Guadalupe system lying west of the delta. Therefore, the coarse silt on the northern outer shelf in the vicinity of the Brazos-Colorado delta and the ancestral Guadalupe valley is palimpsest in origin and apparently is being reworked by the strong semipermanent bottom currents that prevail in this region.

In the southern part of the south Texas inner shelf, the patterns of distribution of first-cycle and multicyclic coarse silt indicate that the inner shelf coarse silt is also relict in origin. The apparent absence of modern coarse silt on this part of the inner shelf is thought to reflect the paucity of this size fraction in the major sediment sources of the southern inner shelf—the Rio Grande and Padre Island. In the northern part of the south Texas inner shelf, evidence indicates that the inner shelf coarse silt is a mixture of palimpsest silt reworked from the late Pleistocene substrate and modern silt provided by the Colorado River.

MCBRIDE, EARLE F., Univ. Texas, Austin, TX

Rules of Sandstone Diagenesis Related to Reservoir Quality

The reservoir quality of sandstone is almost entirely controlled by diagenetic events. The chemical and physical processes responsible for diagenesis are complex and they influence sand during all stages of burial and, in some basins, during subsequent uplift. Petrographic studies by many workers in the past 10 years provide the basis for formulating rules of sandstone diagenesis that help in predicting reservoir quality in differ-