

mosaic of fault blocks that were differentially active through geologic time. The buildups are elongate northwest, and their distribution and overall shape appear to be controlled by northwest-trending paleostructures. Some larger buildups (i.e., ls may) show local northeast-trending thickens within an overall northwest-trending buildup.

Examination of Landsat imagery revealed an extensive network of northwest- and northeast-trending lineaments that parallel linear patterns apparent from aeromagnetic, gravity, and subsurface isopach data. Additionally, outcrops along selected lineaments contain fractures that parallel these lineaments, suggesting that the lineaments are related to fundamental (i.e., basement) fracture zones along which algal buildups may have developed. Comparison of the fracture network to the distribution of algal thickening reveals these buildups occur predominantly along northwest-trending lineaments. Local disruptions within and apparent terminations of the buildups correspond to cross-cutting northeast-trending lineaments. This relationship provides guidance to locating prospective algal buildups. Integration of these data with detailed subsurface mapping can refine some leads into prospects. Several of these features have been successfully drilled.

MERIN, I. S., Earth Satellite Corp., Chevy Chase, MD, and W. R. MOORE, Petroleum Geologist, Denver, CO

#### Application of Landsat Imagery to Hydrocarbon Exploration in Niobrara Formation, Denver Basin

The Niobrara Formation produces commercial quantities of oil from fractures in several places in the Denver basin. The Niobrara in this basin is an oil-prone, mature source rock having as much as 3.4% TOC, and has been in the generating window since early Eocene. This implies that hydrocarbon generation from the Niobrara is partly contemporaneous with the Laramide orogeny. The Laramide was a multiple-phase orogenic event that began with compression directed to the east-northeast during the Late Cretaceous to Paleocene and ended with compression directed to the northeast during the Eocene. We believe the Eocene phase activated northeast-trending extension fractures that may have acted as loci for storage and migration of hydrocarbons, locally generated in the Niobrara. The auto-fracturing pressures related to hydrocarbon generation in the Niobrara theoretically would preferentially open and fill this northeast-trending fracture system.

Examination of Landsat imagery shows that zones of northeast-trending lineaments are present throughout the basin. Numerous northeast-trending faults are present in the basin, and many overlie older zones that were reactivated during the Laramide. This suggests that these lineaments are previously unrecognized fracture zones. We have defined an exploration fairway within the basin based on subsurface isopach and resistivity mapping. We believe that mapping of northeast-trending fractures can help identify leads (within this fairway) prospective for Niobrara production. Support of this concept is the location of several apparently productive Niobrara wells along a zone of northeast-trending lineaments.

MESCHER, PAUL K., and JAMES C. POL, ARCO Exploration, Denver, CO

#### Sedimentation and Tectonic Implications of Cambrian-Ordovician Clastics, Renville County, North Dakota

Cambrian-Ordovician clastics of the Deadwood Formation were studied in detail from Newporte field in Renville County, North Dakota. This small Cambrian-Ordovician oil pool was extensively cored, often to the Precambrian basement, allowing close examination of clastic deposition influenced by local basement tectonics.

In Renville County, the basal unit consists of a well-rounded, fine to medium-grained glauconitic quartz sandstone. Paleohighs appear to have had a pronounced effect on Deadwood sedimentation. Sands, from quiet water settings, show poor to moderate sorting, are commonly finely laminated, and/or show traces of minor small-scale cross-bedding. In places, bedding planes are highly disrupted, suggesting intervals of intense bioturbation (Skolithos). Sands associated with paleohighs are clean, well sorted, and commonly friable. Their association with basement structure is suggestive of beach-barrier-bar sequences related to irregularly upthrown basement blocks. In one example, this clean basal

sand is associated with an upthrown basement block and is sharply truncated by the pre-Winnipeg (early Ordovician) unconformity.

The first unit above the basal sandstone in structurally lower wells is an anomalous conglomerate unit. Large angular basement clasts up to cobble size were viewed in core. This unit grades upward into a fine sand sequence and distally grades into a marine sand. It terminates abruptly in upthrown wells and indicates rapid fault movement and offset during middle Deadwood deposition, with development of localized fanglomerate sequences associated with fault scarps.

Immediately capping this sequence is a dark-gray marine shale that thins depositionally toward paleohighs.

MILLIKEN, K. L., P. B. GOLD, and L. S. LAND, Univ. Texas at Austin, Austin, TX

#### Feldspar Diagenesis in Neogene Sediments, Northern Gulf of Mexico

Alteration of feldspars in the youngest of the Gulf Coast Cenozoic sands and sandstones is dominated by dissolution and albitization. Volumetrically significant amounts of alteration are only observed below burial depths of about 4.5 km in sands of Pliocene and Miocene age. Only trivial amounts of plagioclase dissolution were observed in Pleistocene units. In general, plagioclase exhibits minor amounts of secondary dissolution at all depths, and greater amounts in the deepest samples. Potassium feldspar is subject to very little dissolution to depths of about 3.5 km; by 4.5 km K-feldspar removal is virtually complete. Albitization affects only plagioclase and appears to be operative, in these sediments, over temperatures of 110°C-140°C.

Compared to older Cenozoic units elsewhere around the Gulf of Mexico feldspar dissolution and albitization in Neogene sands have advanced to a lesser degree, at least in the sense that they affect a smaller proportion of the total section. Comparison of pre-alteration feldspar composition, temperatures of alteration, and geothermal gradients for Gulf Coast sandstones of different ages suggests that the main controls on feldspar alteration are temperature, pre-alteration plagioclase composition, and possibly the amount of fluid flow. Time per se seems to be a factor of negligible importance, at least over time spans greater than 10<sup>6</sup> yr. Thus, the lesser volume of Neogene sand affected by feldspar dissolution and albitization can be attributed primarily to the lower geothermal gradients of the northern Gulf.

MITCHELL, STEPHEN M., Mobil Producing Texas & New Mexico, Houston, TX

#### Reservoir Description Applied to Iatan East Howard Field, Mitchell County, Texas

A reservoir description has been completed to help improve recovery over Mobil Producing Texas and New Mexico's leases in the Iatan East Howard field in Mitchell County, Texas. The following 5 phases of study have been used: core description, log analysis, stratigraphy, mapping, and follow-up.

Two 850-ft cores were used to study depositional environment and diagenetic history. Clear Fork (Permian Leonardian) pay is characterized by thin, discontinuous porosity zones cut by vertical fractures. Porosity development is controlled by facies zonation from a tidal flat similar to that on Andros Island today. Porosity developed in carbonate-sand accumulations along tidal channels that meander and bifurcate across the flat. Thickness and continuity increase northwestward toward the basin. Six major zones of transgressive channel belt sediments separated by regressive supratidal marsh sediments were identified in the upper Clear Fork.

Digitized logs were analyzed and log analysis was accepted where porosity matched whole-core porosity measurements. Histograms of log response were used to normalize uncored wells to cored wells for equivalent results.

Stratigraphy was studied by cross sections of each zone. Facies-biased contouring was used to map reservoir parameters. Pay thickness was mapped using 2 porosity cutoffs with the higher identifying the best reservoir. Maps were digitized and used to calculate the original oil in place. The fracture orientation was recognized as N60°E from injection-water breakthrough, water sampling, and water-oil ratio mapping. Injectors will be placed along this trend perpendicular to porosity trends. Follow-up with reservoir engineering will continue with comparisons of mapped

volume versus production histories and will influence pattern alignment and well spacing.

MITCHELL, STEVEN W., California State College, Bakersfield, CA

#### Quaternary Eustatic Sedimentary Accretion of Southern Bahamas Archipelago

Surficial geologic mapping indicates that the southern half of the Bahamas Archipelago is forming by the accretion of discrete depositional sequences resulting from successive eustatic sea level changes: (1) multiple beach and dune ridges, (2) estuarine, (3) lacustrine, (4) shallow subtidal, (5) reef and reef rubble, and (6) megadune complexes. The lithologies are accreted along unconformable erosional-solutional contacts—marine terraces and subaerial caliche crusts. During periods of significant transgression, sequences 1-5 are accreted. Sediments are predominantly skeletal and peloid. During periods of significant regression, megadune complexes are accreted. Ooids are the dominant sediment.

Erosional-solutional features reflect areas of subaerial exposure and/or coastline erosion. Terraces at 10, 20, and 40 ft elevations are preserved along arid eastern Great Inagua Island. The calichification of Bahamian Quaternary carbonates has concentrated insoluble residues (quartz, feldspar, heavy minerals, crandallite, micrometeorites). Insoluble residue analysis provides a basis for the correlation of accreted eustatic sedimentary sequences.

MOLNIA, BRUCE F., U.S. Geol. Survey, Sioux Falls, SD, and WILLIAM C. SCHWAB, U.S. Geol. Survey, Woods Hole, MA

#### Potential Geologic Hazards of North Aleutian Shelf, Bristol Bay, Alaska

Federal OSC lease sale 92, North Aleutian shelf, Alaska, is scheduled for April 1985. The area, located in the southeastern Bering Sea, has 3 basins with sedimentary thicknesses in excess of 4 km.

Six geologic conditions that could cause problems during petroleum development are: (1) seismicity, (2) recent faulting, (3) gas-charged sediment, (4) bed forms and active sediment transport, (5) scours, and (6) volcanism.

Since 1953, the region has a history of at least 10 shallow earthquakes, including a 1971 back-arc event with a Richter magnitude of 5.2. The largest event impacting the entire region, a Richter magnitude 8.7 earthquake, occurred in 1938. Normal faults are located along the southern edge of the St. George basin, and on the northeastern edge of the Amak basin. Many exhibit increased offset with depth, surficial sags, and small surficial cracks. Surprising was the absence of any evidence of sea-floor sediment instability. "Sonar bright spots," and possible, near-surface gas-charged sediment occur west of Amak Island and north of Unimak Island. An area of megaripples and dunes covers more than 1,500 km<sup>2</sup>. Bed forms have spacings of 20-50 m and heights of 1-3 m. Observations suggest that coarse sand may be actively transported. Thousands of scours, many linear and parallel, some greater than 800 m long, 250 m wide, and incised up to 5 m, were identified. Pavlof, an Alaskan Peninsula active volcano, located 45 km northeast of Cold Bay, has a continuous history of steam release and occasional eruption. Lahars, nuée ardentes, and other volcanogenic activities are possible, but their probabilities are unknown. None of the geologic conditions identified precludes petroleum development or production. The potential impact of these factors must, however, be included in planning for future petroleum activities.

MONTZ, MELISSA J., and DAG NUMMEDAL, Louisiana State Univ., Baton Rouge, LA, and DON SWIFT, ARCO Exploration and Production Research, Dallas, TX

#### Shelf Storm-Deposited Sandstones, Upper Mancos Shale, San Juan Basin, New Mexico

Hummocky cross-stratified and amalgamated sandstone beds in the upper Mancos Shale of the San Juan basin record sedimentation on a storm-dominated shelf. Interbedded silty shale, siltstone, and sandstone form coarsening-upward cycles in Mancos shelf deposits. Abundance of cycles suggests storms were frequent, and preservation potential of shelf storm deposits is high.

Mancos storm beds typically have an erosional base with tool marks, and planar laminations overlain by hummocky cross-stratification

(HCS). HCS comprises 20-90% of the primary sedimentary structures in these beds. Wave-ripple lamination commonly overlies HCS in ≤2-m thick beds high in the section. Less commonly, wave ripples overlie planar lamination in some thinner beds (5-15 cm) low in the section. In comparison, planar lamination commonly overlies HCS in storm sequences. Paleoflow trends are northeast to southwest, normal to presumed regional paleoshoreline.

Distal to proximal changes in the beds are seen in vertical section. Increases in sand to shale ratio, amalgamated beds, abundance of sole marks, average grain size (from 0.05 to 0.125 mm), and wavelength of HCS are seen upsection. Bioturbation of beds decreases upsection. These trends reflect progressive shoaling of the Mancos shelf.

Upward-coarsening interbedded silty shale, siltstone, and sandstone intervals comprise the bulk of upper Mancos shelf deposits. A hummocky storm bed 30-100 cm thick caps each cycle, and contact with the overlying interval is sharp. Contact with the overlying Point Lookout shoreface sandstone is abrupt. The cycles indicate that episodic storm sedimentation characterized the Santonian-Campanian shelf in the study area.

MOORE, CLYDE H., Louisiana State Univ., Baton Rouge, LA, KATHY MCGILLIS, Shell Oil Co., Houston, TX, SARA STEWART, Pennzoil, Houston, TX, SCOTT WILKINSON, Superior Oil, Lafayette, LA, and GILLIAN HARWOOD, Univ. Newcastle upon Tyne, Newcastle upon Tyne, United Kingdom

#### Upper Jurassic of East Texas, a Stratigraphic Sedimentologic Reevaluation

The Smackover-Haynesville of east Texas has long been modeled as a simple progradational carbonate-evaporite ramp. Recent data indicate that the conventional ramp model for this sequence should be abandoned in favor of an evolving rimmed shelf to platform model, forming in response to changes in rate of relative sea level rise during the Late Jurassic. Evidence for Smackover-Haynesville shelves include: (1) thick high-energy carbonates along the basin margin in the Smackover and throughout the Haynesville, (2) low-energy pellet-dominated lagoonal carbonates, evaporites, and evaporitic siliciclastics occurring landward of, and interfingering with, the Smackover and Haynesville basin-margin carbonate barriers, (3) deeper water, open-marine low-energy limestones with black shales seaward of the basin-margin barriers (Smackover-Gilmer undifferentiated), and (4) the Gilmer shale forms a siliciclastic wedge seaward of the Haynesville basin margin and its zero isopach defines the Kimmeridgian shelf margin. The Smackover and Haynesville seem to represent 2 distinct sedimentologic cycles, with each cycle reflecting an initial relative sea level rise during which a rimmed shelf and lagoon are developed, and a terminal sea level standstill during which the shelf evolved into a high-energy platform. Although these sedimentologic patterns seem compatible with accepted Jurassic sea level curves, they may also reflect differential basin-margin subsidence combined with variable carbonate production rates. Finally, the shelf-platform model more clearly defines future exploration strategies for Smackover-Haynesville targets in east Texas and perhaps across the Gulf of Mexico, if eustatic sea level changes were the dominant causative factor for shelf development in the Late Jurassic.

MOORE, GEORGE T., and R. W. JONES, Chevron Oil Field Research, La Habra, CA

#### Organic Facies and Petroleum Potential of Eastern North American Margin

Sufficient data now exist to permit reconstruction of the organic facies for Late Jurassic and middle Cretaceous times when the potential for source rock deposition was greatest on the eastern margin of North America. Three distinct and one mixed organic facies have been mapped for these 2 geologic intervals: facies B (oil-prone) amorphous organic material plus alginite and exinite deposited under marine anoxic conditions, facies C (gas-prone) terrestrial plant debris deposited in mildly oxic environments, facies B-C (waxy oils and gas condensate) mixed facies, and facies D (nonsource) degraded and/or recycled organic material.

Distribution of the Late Jurassic organic facies was greatly influenced by the relatively arid subtropical climate and the shelf-edge carbonate