

ISPHORDING, WAYNE C., and JOHN A. STRINGFELLOW, Univ. South Alabama, Mobile, AL, and GEORGE C. FLOWERS, Tulane Univ., New Orleans, LA

#### Sedimentary and Geochemical Systems in Transitional Marine Sediments in Northern Gulf of Mexico

The coastal zone of the northern Gulf of Mexico is marked by a series of bays and estuaries that serve as the principal depositional basins for rivers draining an area of greater than 160,000 km<sup>2</sup>. These rivers annually contribute a sediment load to the basins in excess of 12 million tons. Because each river drains a watershed of different lithologic character and each river is further characterized by a different flow regime and hydraulic properties, the sediments deposited in the marginal basins have unique characteristics.

Extensive municipal and industrial dumping of effluent over the years has also acted to imprint geochemical differences on each of the depositional basins. Depending on the degree of industrialization within the watershed, the bays and estuaries may be described as heavily impacted (Mobile Bay), moderately impacted (Apalachicola Bay, Mississippi Sound), or slightly impacted (Pensacola Bay). A strong correlation was observed between the degree of heavy metal contamination and the textural and organic content of the sediments. Analyses further indicated that most metals were partitioned in the bottom sediments in forms that would permit their subsequent release back into the water column or would allow transference of the metal to fauna by ingestion.

JACOBS, J., Sohio Petroleum Co., San Francisco, CA

#### Fluvial Responses to Hydrologic Changes on Red River in Northeast Texas

Changes in meander patterns of the Red River in northeast Texas from 1860 to 1980 were identified from various maps, aerial photographs, and fieldwork by a 430-km stretch from the Denison Dam on Lake Texoma downstream to Texarkana. Denison Dam closure in 1943 resulted in increased daily base flow and substantially reduced frequency and intensity of flood peaks. The Red River responded to the closing of the dam by increasing width, depth, meander wavelength, amplitude, radius of curvature, and channel length.

Changing only discharge and sediment load downstream from the dam reveals a geologically instantaneous fluvial response to the dam closure. Qualitative prediction of these recent changes on the Red River is generally confirmed by empirical studies in the literature.

Three ancient meander patterns preserved on the Holocene Red River flood plain record a different hydrologic regimen in which well-sorted, clay-rich sediment was transported in a paleoriver having low wavelength, amplitude, width, and depth with high sinuosity. Bankfull discharge is estimated to have been quite low. Archeological remains suggest these features formed 5,000 ± 1,000 years ago.

In perspective, the isolated hydrologic changes that occurred as a result of the closure of the Denison Dam are minute by comparison to the climate-related changes the Red River has undergone over the last 5,000 years.

JANKS, J. S., Robertson Research U.S., Inc., Houston, TX (present address: Texaco USA, Houston, TX), and T. SANNESS and B. A. RASMUSSEN, SAGA Petroleum A/S, Hovik, Norway

#### Diagenesis of Cotton Valley Sandstones, Catahoula Creek Field, Southern Mississippi

Three conventional cores from the deep Cotton Valley clastics in south Mississippi were analyzed by thin-section petrography, scanning electron microscopy, x-ray diffraction, and cathodoluminescence. The  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  values were determined on carbonate cements in sandstones. The sandstones are predominantly subarkose, arkose, and quartzarenites, with minor amounts of quartz and feldspathic wackes. K-feldspar, muscovite, and plagioclase are relatively fresh. Authigenic clay in sandstones is almost exclusively chlorite that occurs as a pore-lining and pore-filling phase, most commonly associated with porous zones. Feldspar diagenesis includes overgrowths of both potassic feldspar and albite. Typically, fresh detrital K-feldspar grains have overgrowths of K-feldspar that are

partly dissolved, whereas plagioclase grains (with partly dissolved interiors) have unaltered albite overgrowths. Quartz overgrowth development is pervasive where not hindered by chlorite. Carbonates include intergranular calcite, dolomite, and iron-rich dolomite. The earliest phase is a pore-lining dolomite cement, followed by a later ferroan calcite. Ferroan dolomite (or ankerite) is associated with chlorite in relatively porous intervals. The  $\delta^{13}\text{C}$  values of mixed carbonates in sandstones range from -0.8 to -4.4 ‰ (PDB), which is typical of normal marine carbonates. The  $\delta^{18}\text{O}$  values range from -5.0 to -12.3 ‰ (PDB). Formation temperatures of approximately 115°-160°C are suggested by the oxygen isotope data from almost pure late-stage ferroan dolomite.

Petrographic evidence suggests that coarser grained sands were cemented early in the burial history by dolomite and calcite. These cements were later leached by formation waters, possibly related to the generation of organic acids and/or carbon dioxide released during kerogen maturation. Resultant secondary porosity was preserved by formation of bladed chlorite. The source of the iron, manganese, and magnesium necessary for chlorite formation was presumably the release of ions from smectite to illite conversion in shales, because ferromagnesian rock fragments in the sandstones are rare. The feldspar volume and stability and the  $\delta^{13}\text{C}$  values of carbonate cements suggest the pore water was probably initially seawater and later brine that migrated upward from the Louann Salt. Present brines contain 7,300 ppm of potassium and 51,113 ppm of sodium. Meteoric diagenesis of these rocks was likely minimal. These data suggest that original depositional porosities and permeabilities and carbonate cementation are significant controls in secondary porosity formation in the Cotton Valley sandstones.

JOHNSON, D. JAY, Conoco Inc., Oklahoma City, OK, JERRY ALLINGER, Conoco Inc., Larkspur, CO, and ROBERT K. AAKER, Conoco Inc., Littleton, CO

#### Evaluation of Radioelement Geochemistry for Detection of Petroleum Reservoirs

We tested the hypothesis that radioelements are either enriched or depleted over petroleum accumulations. Total count surveys are subject to signal variations due to changes in soil composition and to variations in soil moisture. Methods have been determined to lower the variations due to soil composition and to eliminate variations due to soil moisture. We conclude that, if hydrocarbon-related anomalies are present, they are overwhelmed by changes in soil composition.

Changes in soil composition can cause: (1) a fourfold change in total count and eU, (2) a twofold change in K, and (3) a thirteenfold change in eTh. Modified eU/K, eTh/K, and eU/eTh ratios reduce the variations caused by soil composition to ±50% of normal conditions. This permits detection of anomalies as low as 1.0 ppm eU, 2.0 ppm eTh, or 0.7% K. Statistical detection limit for the surveys is ±0.24 ppm eU at 2.0 ppm eU concentration.

The detected anomalies are of equal amplitude and frequency both on and off the fields. This, plus the lack of anomaly repeatability from line to line, precludes a hydrocarbon source. The majority of all anomalies are attributed to soil or lithologic sources. The others are caused by uranium mineralization.

KINGSTON, JIM, Baylor Univ., Waco, TX

#### Long-Term Effects of In-Situ Leach-Mining Restoration in Oakville Aquifer System near George, West Texas, with Implications for Shallow Waterflooding

The Miocene Oakville Formation is a major Gulf Coast freshwater aquifer and uranium host. The impact of long-term pumping during the restoration phase of leach mining depends on the efficiency of water-quality improvement methods, characteristics of the aquifer, and meteorological fluctuations affecting recharge. Newly acquired data from this industry have improved understanding of fluid migration in a fluvial depositional system.

Development of the in-situ leach-mining industry has added vast amounts of subsurface hydrogeological data obtained from operation of an extensive network of wells. Both surface and subsurface data were combined in this investigation, which included analysis and interpretation of outcrop samples, uranium drill logs, cross sections loaned by com-

panies, pumping test data, water-well data furnished by the Texas Department of Water Resources, and geological and soils maps. This information was incorporated into the U.S. Geological Survey three-dimensional finite-difference computer model to predict ground-water behavior in a heterogeneous and anisotropic aquifer. Results indicate leakage through confining beds and communication across growth faults, suggesting that the Oakville is a complex aquifer system rather than a simple confined aquifer.

Understanding the development of intricate flow paths and controlled movement of fluids through designed and controlled systems may improve methods of shallow waterflooding of petroleum reservoirs as well as management of water resources.

KONTROVITZ, MERVIN, Northeast Louisiana Univ., Monroe, LA

#### Ocular Sinuses in Genera of Ostracod Family Trachyleberididae

The ocular sinuses of seven genera of ostracods (Trachyleberididae) were studied through the use of internal molds and thin and polished sections. The three-dimensional aspects of the ocular sinus in the genera *Actinocythereis*, *Henryhowella*, *Malzella*, *Orionina*, *Puriana*, and *Radi-mella* have not been studied before. *Echinocythereis*, previously studied, was included for comparison.

Morphology of the sinuses varies from the low moundlike form in some specimens of *Henryhowella*, to the long structure in *Actinocythereis*. With the exception of *Henryhowella*, all other genera have ocular sinuses with many features in common. These properties include a stalk-like appearance marked by a distinct constriction someplace along the length of the stalk. Most of the sinus is circular in cross section except the distal portion, which is expanded and irregular in shape, presumably to accommodate the portion of the eye with the lens cell, retinal cells, and rhabdomes. The surface of the distal portion is dominated by a concavity that is the complement of the convexity of the adjoining eyespot. Adjacent to the convexity is an anterior lobe or rim and usually a higher posterior rim. Channels with diameters of 2 to 6  $\mu$ m probably are passageways for axons. Ontogenetic development includes an early, low, cone-shaped sinus with a central concavity. Later juveniles in most taxa have a stalk, as do adults.

Despite overall similarities of the ocular sinuses, detailed examination indicates that genera and species can each be distinguished using this feature. Variations in size, shape, proportions, and secondary topography are diagnostic; thus, another character is available for taxonomic purposes.

KOSTERS, ELISABETH C., Louisiana Geol. Survey, Baton Rouge, LA

#### Deeply Buried Tertiary Sand Bodies in Northern Gulf of Mexico: Examples from Lower Hackberry (Oligocene) and Houma Embayments (Miocene)

Ninety-three electric logs were used to analyze deep-seated sand bodies in two fields in the Hackberry embayment of southeastern Texas and the Houma embayment of southern Louisiana. The geometry and occurrence of these sand bodies can be related to the regional structural and sea level history of the study areas. The Hackberry embayment originated by shelf-edge retreat after a major drop in sea level. Large amounts of sediment were then supplied by the Frio delta systems that built out to the shelf margin. The Houma embayment originated during a slight increase in the rate of sea level rise, while the Miocene deltaic systems built out over the continental shelf, triggering mass movement on the continental slope.

Both areas are characterized by large-scale synsedimentary faulting and salt tectonics. Large arcuate growth faults form the updip boundary of each embayment. Displacement across faults is from 900 to 2,000 ft (300 to 650 m). In both embayments, the sand bodies are overlain by thick shale masses that contain similar "flysch-type" faunas. The sands have a blocky appearance on the SP curves of electric logs, are slightly more than 100 ft (35 m) thick, and can be correlated for only a few miles. The structures show typical rollover, or reverse drag, into the main growth faults. In the Hackberry embayment, it can be shown that some intervals on the expanded downthrown side of the fault contain additional section, whereas marker beds below and above continue across the fault. These expansions developed during a time interval represented by 400 ft (130 m)

of section and are interpreted as erosional and depositional events across the growth fault.

Comparison with present-day large-scale rotational slumps on the continental slope off the Mississippi delta shows that the overall characteristics of the Tertiary sands are similar to such modern slump masses; therefore, these sands are interpreted as having originated in a similar fashion. Thus, it appears that these analogous sand bodies accumulated as a result of comparable processes, even though the associated changes in sea level in the two areas differed considerably.

LEHMAN, THOMAS M., Univ. Texas at Austin, Austin, TX

#### Transgressive-Regressive Cycles and Environments of Coal Deposition in Upper Cretaceous Strata of Trans-Pecos Texas

Paralic deposits of Late Cretaceous age are represented in Trans-Pecos Texas by the Aguja Formation in the Big Bend region and the correlative San Carlos Formation in the Sierra Vieja region. Although both units are broadly progradational, each records a single widespread transgressive pulse. This transgressive event began in the early Campanian and reached a peak during the middle Campanian. Regressive (progradational) deposits consist of prodeltaic shale and delta-front siltstone and claystone, overlain by coalesced lenticular fluvial-dominated deltaic sand bodies. Transgressive deposits consist of local storm-dominated shoal sandstone, thin laterally extensive oyster-rich sublittoral sand sheets, and shelf mud. Transgression occurred relatively rapidly and was expressed differently depending on the distance traversed landward of the previous shoreline. Coal and lignite deposition occurred in restricted interdistributary areas during progradation, and in more widespread areas during peak transgression when the strandline was neither advancing nor retreating. No evidence is found for barrier island development during transgressive events; hence, the coal and lignite accumulated along a muddy nonbarred coast in marshes directly adjacent to the sea. The coal-bearing strata consist of an alternating series of dark-gray root-mottled carbonaceous claystone with sideritic "ironstone" concretions, coal or lignite, and well-laminated light-gray pyritic shale with calcareous concretions bearing marine invertebrates. The cyclic alternation of these facies suggests the repeated inundation of coastal marshes by marine waters, followed by mud accumulation, subsequent regrowth of the marsh, and the deposition of lignite or coal.

LEVIE, DONN, JR., Highland Resources, Inc., Houston, TX

#### Interdomal Sediment Ponding: A New Lower Hackberry Play?

Lower Hackberry (mid-Frio) reservoirs are highly unpredictable in the subsurface due primarily to variations in sand thickness and sand distribution. Most exploration for lower Hackberry reservoirs has been limited to turbidite sand-filled channels cut between sea-floor paleotopographic highs, and lower Hackberry sands on the northern flanks of salt domes, where excellent stratigraphic traps exhibit north dip—in places with significant closure. Two other existing types of reservoirs have received little attention, both in the literature and in exploration. The suprafan lobe in the midfan position lying at the base of the slope is a prime exploration target, where massive lower Hackberry sands should be found. Several wells in northwestern Cameron Parish, Louisiana, penetrated thick lower Hackberry sands in a downdip position from the channel plays, suggesting that a mechanism for large-scale downslope movement was present during Hackberry deposition.

Another potential reservoir that deserves more exploration is the interdomal sediment "ponding" of lower Hackberry sands into topographic lows created by the scouring of the sub-lower Hackberry surface. These sands generally are concentrated in the center of the pond and may not be present in either the traditional lateral or updip pinch-out positions. Many of the ponds occur independently from ideal structural location, commonly with a pinch-out of the sands serving as the trap. Ideal prospect locations are where the sub-lower Hackberry unconformity and top of the lower Hackberry marker diverge, indicating the presence of a sand-filled scour feature. Southwestern Calcasieu Parish provides excellent examples of this ponding feature in the subsurface. Successful exploration efforts depend on careful attention to paleontological information,