

Westchase Hilton • 9999 Westheimer
Social 5:30 p.m., Dinner 6:30 p.m.

Cost: \$25 Preregistered members; \$30 non-members & walk-ups

The HGS prefers that you make your reservations on-line through the HGS website at www.hgs.org. If you have no Internet access, you can e-mail reservations@hgs.org, or call the office at 713-463-9476. (include your name, e-mail address, meeting you are attending, phone number and membership ID#).

by *Raymond P. Sorenson*
Anadarko Petroleum Corporation
Houston, Texas

A Dynamic Model for the Permian Panhandle and Hugoton Fields, Western Anadarko Basin

Panhandle-Hugoton, the largest North American gas field, has long been controversial because of extreme subnormal pressures, variable gas composition and tilted fluid contacts, commonly attributed to hydrodynamic flow despite the absence of an effective updip aquifer. These anomalies are explained in terms of a basin-scale petroleum system history, largely independent of the geographically underlying pre-Permian system.

Giant Amarillo uplift drape structures trapped hydrocarbons immediately following Permian evaporite deposition.

Panhandle-Hugoton pressures, upon discovery, were subnormal relative to drilling depth but normal relative to reservoir outcrop elevations in eastern Kansas, indicating that pressures are controlled by aquifer communication with the surface rather than burial depth. Variations in fluid contacts, pressure, and gas composition suggest that reservoir fluids are still moving, driven by decompression and the rapid volumetric expansion of a supergiant gas accumulation. ■

Hydrocarbons were already being generated in the deep Anadarko basin during the Early Permian, with efficient southward migration from all potential source rocks via bounding faults and Pennsylvanian-Permian alluvial fans. Giant Amarillo uplift drape structures trapped hydrocarbons immediately following Permian evaporite deposition. The pre-Laramide Panhandle field, at maximum burial depth and pressure, contained most of the oil and gas now found in mid-continent Permian reservoirs.

A more detailed version of this presentation can be found in the July 2005 *AAPG Bulletin*, v. 89, no. 7, p. 921-938.

Biographical Sketch

RAY SORENSON received his BS degree in geology from Michigan State University in 1972 and his MA degree in geology from the University of Texas at Austin in 1975. He worked for Texaco from 1974 to 1975, and has been employed by Anadarko, in Oklahoma City and Houston, since 1976. The majority of his career has focused on the geology of the United States mid-continent.



The Early Tertiary Laramide orogeny redistributed Panhandle field fluid columns, possibly spilling gas into the Hugoton embayment. Subsequent erosion of Permian reservoir facies in eastern Kansas allowed water discharge to outcrops at elevations below the regional hydraulic head. As regional pressure dropped in response, the Panhandle field gas cap expanded rapidly, forcing a Late Tertiary-Quaternary mass movement of gas northward to fill Hugoton and associated fields.

Sept 15 Dues Renewal Deadline

Members are reminded that the deadline for dues renewal is September 15. Members who have not renewed their dues (\$24.00 active/associate; \$12.00 emeritus/student) by that date will be dropped from the mailing list for the HGS *Bulletin* and be charged the non-member rate for HGS events. There will be a \$5.00 reinstatement fee for members renewing their membership after 9/15. Active status will be placed on hold until dues are received by the HGS office.