

chroism especially pronounced in the short-wave portion of the visible spectrum. Brown biotite-phlogopite from Old Chelsea, P.Q. is optically similar. The effect is caused by a broad transmission maximum for light vibrating parallel with the mica sheet and a narrow transmission maximum for light vibrating across the sheet. Green phlogopite from Old Chelsea shows less marked reverse pleochroism. With thick sheets transmission maximum Y is at 6200 Å and Z at 5800 Å. No absorption maxima were noted in these varieties. Chemical data suggest the reverse pleochroism may be associated with ferric iron in tetrahedral sites.

HOUSE, M. R., University Museum, Oxford, England
 SUCCESSION OF *TORNOCERAS* AND RELATED GENERA IN THE DEVONIAN OF NEW YORK AND ADJACENT STATES

A study of the species of *Tornoceras*, *Parodiceras*, and *Tornoceras* and *Epitornoceras* from the western U.S.A. provides an independent stratigraphical zonation for the New York State Devonian. It also provides a study of allomorphosis in *Tornoceras*. Comments are made on the protoconch and the significance of the metamorphosis marked by the nepionic constriction in *Tornoceras*.

Where possible in twenty successive faunas, ontogenetic details have been elucidated, and this has been possible from protoconch upward at eleven successive stratigraphic levels. These successional ontogenies shed light on the evolution of the stock. Faunas at each level may be defined morphologically, but few consistently maintained evolutionary trends have been observed. Shell form seems particularly subject to independent, and probably phenotypic, variation. Through the equivalents of the Middle Devonian to the Lower Frasnian, protoconch width appears to increase progressively. Similarly, the suture becomes more undulating, particularly with regard to the ventrad face of the lateral lobe. Later species show reversion to early characters in these respects.

The origin of *Tornoceras* from *Parodiceras* is argued, and it is considered that *Tornoceras* gave rise to all later members of the Tornoceratidae.

HOWE, HERBERT J., Louisiana State University, New Orleans, Louisiana

MONTOYA GROUP (ORDOVICIAN) CARBONATE PETROLOGY OF TEXAS AND NEW MEXICO

The Montoya Group (Middle and Upper Ordovician) of west Texas and New Mexico consists of several carbonate types including crinoidal calcarenite with a microcrystalline matrix, crinoidal calcarenite with sparry calcite cement, calcilitute with abundant chert, chert-free calcilitute, shell limestones, calcirudites, and completely or partly dolomitized equivalents of the former.

Dolomitization and silicification are volumetrically important. Recrystallization and (or) replacement obscure much of the original texture and fossils. Dolomitization begins with the formation of sporadic small crystals, which increase in number until a complete dolomitic mosaic results. Coarse fossil debris becomes progressively reduced in texture as the process advances to completion. Texture and degree of silicification are not related in many places. Fluctuating silica supply in the sea water is strongly indicated. Intervals of abundant chert are separated by less cherty strata.

Study of the unaltered rocks show considerable range in environment from high-energy, shallow-water to low-energy, deep-water conditions. Montoya sedimentation can be compared with correlative strata deposited dur-

ing the Middle and Upper Ordovician submergence of the North American continent. Montoya deposition in the Cincinnati is estimated to be 320 feet in 10 million years—very slow when compared with 4,500 feet in 10 million years for the Cayugan autogeosyncline of Michigan (Kay, 1951).

Present lithologic and stratigraphic data support rejection of a disconformity within the Montoya Group, although several minor erosional breaks occur.

ILLING, L. V., Cheam, Surrey, England, and WELLS, A. J., Koninklijke/Shell Exploratie en Productie Laboratorium, Rijswijk, The Netherlands

PENECONTEMPORARY DOLOMITE IN THE PERSIAN GULF

Carbonate sediments dominate the shallow waters along the arid southwest side of the Persian Gulf. In the more protected parts of the west coast of the Qatar Peninsula, the processes of near-shore sedimentation have created lagoons and embayments with high chlorinity (30–35 g/l) and reduced tidal range; they are separated from the normal Gulf waters (22–24 g/l) with their average 4-foot tides, by many miles of sea less than 2 fathoms deep.

The lowest parts of the lagoonal shores are fringed by salt flats—"sebkhas"—varying in width from a few tens of yards to several miles. The sebkhas pass seaward into the intertidal zone, commonly via an intermediate algal flat. This is just covered by normal high tides, but only with favourable winds can very occasional spring tides reach far onto the sebkha surface which is a few inches higher.

Sedimentation is gradually filling the lagoons by the seaward advance of the environmental belts, so that sebkha sediment overlies stromatolitic algal laminae, and these are underlain by intertidal mud-pellet sands resting on lagoon muds.

The chlorinity of the pore waters increases landward and upward in response to surface evaporation losses. It increases rapidly within the algal flat (50–130 g/l), where small selenite crystals form beneath the higher, landward parts. Together with the continuing precipitation of aragonite, this causes an increase in the Mg/Ca ratio of the pore waters from the normal marine value of 3 in the lagoon to more than 10 at the sebkha edge. Within the sebkha, the ratio falls gradually to below 5, while the chlorinity continues to rise slowly to more than 150 g/l. The water table is close to the sebkha surface, and, beneath the uppermost layer subject to large daily temperature changes, the wet sediment reaches well over 40°C. in summer time. Its pH is low (around 6.7) and decreases downward.

These warm magnesium-rich brines cause diagenetic changes in the aragonite sebkha sediment. Dolomite appears. It occurs as a stiff, sticky, tan or tan-gray mud composed of rhombs 1–5 microns in size. Associated with it are turbid flattened crystals of gypsum up to 5 inches across, enclosing, displacing and sometimes replacing aragonite sediment. Depositional textures tend to become obscured, but both macro- and microscopic evidence of relic structures and the changing chemistry of the pore waters make it clear that both the dolomite and the associated coarse platy gypsum are replacing aragonite. They increase in abundance away from the lagoon until they make up the bulk of the sebkha sediment. The dolomite normally appears a few inches beneath the surface, increases rapidly, and almost disappears again in a more irregular fashion within a depth of 2–4 feet.

Carbon-14 determination on two dolomite samples collected within 9–18 inches of a sebkha surface gave

ages of 2,670 and 3,310 years, confirming that the dolomitization is a penetemporary phenomenon related to the present sedimentary environment.

IMBRIE, JOHN, Columbia University, New York, N. Y.

SEDIMENTARY STRUCTURES IN MODERN CARBONATE SANDS OF THE BAHAMAS

Sedimentary structures of carbonate sand bottoms near Bimini, Sandy Cay, and the Berry Islands have been examined in depths ranging to 10 ft. below MLW. Box cores with a vertical face 18 in. high and 12 in. wide were impregnated in the field with epoxy resin to preserve and accentuate the structures.

Four distinct types of internal structures occur in contrasting depositional environments. *Type I*, characterized by 3-10 cm.-thick sets of cross-strata and either horizontal or curving truncation surfaces, is found in intertidal, rippled, oolite shoals. *Type II*, with 20-40 cm.-thick sets of tabular cross-strata and horizontal truncation surfaces, is found in intertidal-subtidal, unrippled, grapestone bars. *Type III*, with even, parallel laminae inclined at low angles to the horizontal, is found in the backshore and foreshore of beaches. *Type IV* is mottled, lacks stratification, and is found in unrippled level bottoms.

Bahamian field observations confirm pertinent hydrodynamic conclusions of J. R. L. Allen, H. W. Menard, and D. B. Simons. Type I is observed to form by small ripple trains induced at low flow intensities by tidal currents. Type II is judged to form by a large-scale ripple induced at high flow intensities by rare storms. According to M. M. Ball, deposits with similar structure were formed by Hurricane Donna in Florida Bay. Type III is observed to form from swash and backwash currents on beaches. Type IV reflects an environment in which organisms destroy stratification as fast as it is produced.

These four types should be considered as end-members representing hydrodynamically pure situations. Many actual sites represent alternating conditions (beach-shoal, beach-bar, etc.), as is clear from preserved structures.

IRVINE, T. N., Geological Survey of Canada, Ottawa, Ontario, Canada

SEDIMENTARY STRUCTURES IN LAYERED ULTRABASIC ROCKS*

Many igneous intrusions show layering formed by gravitational accumulation of crystals that is, both in variety and detail, remarkably similar to the bedding of sedimentary rocks. Such layering occurs in most compositional types of intrusions but especially in mafic and ultramafic bodies. The examples considered specifically are from the Duke Island ultramafic complex in the Alaskan panhandle; the rocks are composed of olivine and clinopyroxene but, where layered, look much like graded bedded turbidity current deposits. The intrusion is composite, and the layering, which has been tilted and folded, has two ages—thus, layered blocks and fragments of olivine pyroxenite are included in stratified peridotite. Over-all, stratification is developed intermittently through an original vertical thickness of 2 miles. Individual layers have been traced for 300 feet, and one continuously layered section is 1,500 feet thick and extends 1,000 feet.

* Published by permission of the director of the Geological Survey of Canada.

A typical layer is 2 inches to 2 feet thick and is graded from grain sizes of 2-10 mm. at its base to 0.1-1 mm. at its top; some mineralogical sorting results because the olivine crystals are generally finer and thus concentrated upward. The base of a layer is sharp; the top may terminate against the next stratum or may pass into an intervening zone of thin laminae. Some of the layers of younger age resemble graded beds of slide conglomerate, their lower part being largely olivine pyroxenite fragments.

Other layering features include: cross-layering, slump and deformation concurrent with accumulation, streamlining of layers over irregularities, load casts, general correlation between layer thickness and particle size, lateral grading, and "diagenetic" recrystallization.

The layering has undoubtedly formed because of magmatic currents during extremely unstable conditions. The currents were probably a convective type of overturn marked by the descent of dense, crystal-laden magma from the cooling roof and wells of the pluton, and it is likely that they were initiated and perpetuated by repeated slumping. The described phenomena illustrate that features of water-laid sediments can form in a vastly different environment in terms of the specific gravities of particles and transporting liquid, and the viscosity of the liquid.

JOHNSON, J. HARLAN, Colorado School of Mines, Golden, Colorado, and DANNER, W. R., University of British Columbia, Vancouver, British Columbia, Canada

PERMIAN CALCAREOUS ALGAE FROM NORTHWESTERN WASHINGTON AND SOUTHWESTERN BRITISH COLUMBIA

Limestone units of the Tethyan Permian eugeosynclinal sequences of northwestern Washington and southwestern British Columbia contain abundant specimens of calcareous algae. The limestone bodies are mostly lenticular in shape and range from a few inches to approximately 1,000 feet thick. They are interbedded with volcanic flows and breccias, tuffs, ribbon cherts, argillites, conglomerates, graywackes, and lithic sandstones which may reach a thickness of 10,000 feet or more.

Limestones of Wolfcampian age on Orcas Island, Washington, contain *Mizzia* in association with bryozoa and the fusulinids *Schwagerina* and *Pseudofusulinella*. Limestones of probable early Guadalupian age on the south side of Mount Pilchuck in Snohomish County, Washington, contain *Mizzia* and *Gyroporella* associated with the fusulinids *Cancellina* and *Schubertella*. Upper Guadalupian age limestones of San Juan Island and western Snohomish County, Washington, contain *Mizzia* and other green algae associated with fusulinids of the *Neoschwagerina-Verbeekina* zone. The youngest Permian limestone, containing the *Yabeina* fusulinid zone (Ochoan?), near Granite Falls, Washington, and near Ashcroft, British Columbia, contains *Mizzia* and several other dasyclads.

JOLLIFFE, A. W., Queen's University, Kingston, Ontario, Canada

STRATIGRAPHY OF THE STEEP ROCK GROUP, STEEP ROCK LAKE, ONTARIO, AND EVIDENCE FOR EVOLUTION OF THE PRECAMBRIAN ATMOSPHERE

Knowledge of the stratigraphy has provided the surest guide in iron ore exploration and development at Steep Rock Lake. Most and possibly all of the 33 million tons produced has come from a single stratigraphic member.