

on the shallow banks flanking the Gulf has produced relatively well-sorted sand, low in layer silicates and organic matter. Some well-sorted material has also been deposited on the flanks of Georges and Browns Banks and in some coastal areas.

Most winnowed fine detritus has accumulated on the flat floors of the deep basins. The sediment there is silty clay composed dominantly of mica, chlorite, and mixed layered mica-montmorillonite. Kaolinite occurs only as traces in a few samples. The minor amounts of sand-size material are composed predominantly of mica and of foraminiferal tests and other biogenic debris. Organic matter exceeds 4 per cent in some of the basins. The carbon/nitrogen ratio of the deposits tends to increase with distance from shore.

HAUN, JOHN D., Colorado School of Mines, Golden, Colorado

CRETACEOUS DELTAS IN THE DENVER BASIN AND RELATIONSHIP TO PETROLEUM

Early Cretaceous and early Late Cretaceous deltaic sediments of the Denver basin include the oil-producing "D" and "J" sandstones (upper Dakota Group). The very fine- to coarse-grained quartzose sandstones are in part cross-bedded and ripple marked, in part marine, and in part nonmarine; contain coal and carbonaceous or lignitic beds, mud pellets, siderite concretions, interbedded claystones; and in some areas grade laterally and vertically through siltstone into marine shale.

Within the "J" sandstone there is a well-defined delta extending northwestward into the Denver basin in Colorado. There are smaller deltas in western Nebraska, and, between these two areas, there is a narrow marine embayment. A similar deltaic pattern is developed in the "D" sandstone. Petroleum is concentrated in the parts of the deltas that have the greatest interfingering of marine and deltaic deposits.

HEEZEN, BRUCE C., Lamont Geological Observatory, Palisades, New York

ABYSSAL BASIN SEDIMENTATION

Clastic sands and silts transported and deposited by turbidity currents have created the vast abyssal plains of the ocean basins and have constructed the abyssal cones and natural levee systems of the continental rise. Clastic silts and lutites, largely transported by ocean currents, have created much of the continental rise and the outer ridges which parallel the continental margins. Biogenous sediment, resulting from near-surface productivity, more or less redistributed by currents, has created the rolling abyssal swales of productive mid-oceanic areas and has contributed to the continental rise and marginal trench sediments.

Although turbidity currents are responsible for the leveling of the abyssal plains, turbidites constitute less than one-third of the sediments beneath the plains. Horizontal size grading in turbidites away from source areas is detectable but small. Apparently a much more important control on size is imposed by the lower courses of the major rivers. Turbidity currents originate near the mouths of several major rivers at the rate of 50 per centum but in many other likely areas none have occurred for thousands of years.

HERRICK, STEPHEN M., and WAIT, ROBERT L., U. S. Geological Survey, Atlanta, Georgia

SUBSURFACE STRATIGRAPHY OF COASTAL GEORGIA AND SOUTH CAROLINA

Sediments ranging in age from Pleistocene to Early Cretaceous (?) overlie metamorphosed basement rocks in most of coastal Georgia, but in southeastern Charlton County there are Paleozoic strata between the Lower Cretaceous (?) and the basement complex. The thickness of sediments ranges from 2,674 feet in Screven County, in the north, to 4,700 feet in Glynn County at the coast; the section thins somewhat toward Florida.

The Quaternary unit includes the surficial clastics of Recent to Pleistocene age. The Tertiary units include the predominantly clastic section of Miocene age; the Oligocene sandstones and their limestone equivalent, the Ocala Limestone; the Lisbon and Tallahatta clastics and their Avon Park-Lake City Limestone equivalents, the Wilcox clastics and their Oldsmar Limestone equivalent, all of Eocene age; and the Clayton and Tamesi Formations and their Cedar Keys Limestone equivalent of Paleocene age. The Upper Cretaceous units include the Lawson Limestone and the equivalent clastics of Navarro age; beds of Taylor and Austin age, both in clastic and limestone facies; and the clastic Tuscaloosa Formation. The oldest unit consists of clastics of Early Cretaceous(?) age; these sediments overlie metamorphosed basement rocks in the north and Paleozoics in the south. Two sections, one coastal and the other inland, show lithologic and faunal relationships which reflect changing depositional environments. Regionally, facies change from clastics in the north to limestones in the south with corresponding changes in microfossil suites. Tentative correlations of the clastic sections with their limestone equivalents are based on lithologic and fossil evidence.

There are no local geologic structures, but the sedimentary rocks of the area show the collective effects of the Carolina arch, the Southeast Georgia embayment, and the Ocala uplift.

HOYT, JOHN H., Marine Institute, University of Georgia, Sapelo Island, Georgia, SMITH, DAVID D., and OOSTDAM, B. L., Ocean Science and Engineering, Inc., Bethesda, Maryland

SEDIMENT DISTRIBUTION ON THE INNER CONTINENTAL SHELF, WEST COAST OF SOUTHERN AFRICA

"Sparker" surveys show that the inner continental shelf along a 400-mile stretch of the west coast of South and South West Africa (Olifants River to Luderitz) consists of large areas of virtually sediment-free bedrock and two well-defined elongate bodies of unconsolidated sediment of probable Late Pleistocene-Holocene age.

The sediment distribution pattern is dominated by a strikingly continuous wedge-like body of silts and clays that averages 7 miles in width. This wedge has a maximum thickness of about 80 feet along its inshore edge which lies approximately 3 miles offshore in 200-300 feet of water. Location of the wedge is controlled by a marked steepening in slope of the bedrock surface which coincides with the contact between the Precambrian basement and a sedimentary section of unknown age which dips gently seaward. At the Orange River, the wedge merges with and is overwhelmed by the river's submerged delta.

A second sediment body of lesser extent is the 1 to 1½-mile-wide inshore lens which lies just seaward of the surf zone along a 90-mile stretch south of Luderitz. This inshore lens averages 20 feet in thickness and consists primarily of silty sand and shell material. Gravels, where present, are generally concentrated near the base of the section, and fill irregularities in the dissected bedrock surface. The inshore lens has accumulated shoreward of a series of low discontinuous ridges, and includes re-