

worked and possibly downwarped Pleistocene shoreline deposits exposed onshore farther south.

HSU, K. JINGHWA, Department of Geology, University of California, Riverside, California

FOUNDERING OF CONTINENTS AND ITS RELATION TO ISOSTASY, CRUSTAL THINNING, AND MANTLE DENSITY CHANGES

Current theories on geosynclinal subsidence and on the disappearance of ancient land masses postulate crustal thinning by subcrustal processes. This idea was a necessary implication of the two premises popular a decade ago, namely: (1) surface elevation is related to crustal thickness according to Airy's model of isostatic compensation; (2) either the crust as defined by the Mohorovicic discontinuity is comparable with the crust envisioned by Airy, who separated a solid crust from a fluid substratum, or there is no significant regional difference in mantle density, and there has been none during the past. Recent geophysical investigations have indicated that isostatic adjustment could result not only from changes in crustal thickness, but also from variation in upper mantle density. This new discovery permits the postulate that thinning of continental crust can be related to removal of surficial sialic material by supracrustal processes, such as erosion, gravity sliding, and overthrusting from elevated regions which owed their surface elevation to an abnormally low mantle density. Subsequent increase in mantle density because of variations in mantle temperature leads to isostatic subsidence of regions of thin crust. Such a combination of crustal thinning by supracrustal processes and isostatic subsidence related to mantle density changes could explain adequately (1) the disappearance of ancient land, (2) the formation of a new geosyncline at the site of an ancient land, and (3) the estimated chemical composition of the earth's crust. The efficacy of supracrustal processes to remove large quantities of sialic materials has been discussed. It is not necessary to postulate subcrustal processes which either must assume lateral transfer of vast quantities of crustal material or that the Mohorovicic discontinuity is a phase change boundary. An isostatic model relating surface elevation, crustal thickness, and mantle density variation during an orogenic cycle is presented.

JOHNSON, J. HARLAN, Department of Geology, Colorado School of Mines, Golden, Colorado

LOWER CRETACEOUS ALGAE FROM SOUTH TEXAS

Appreciable numbers of algae occur in the Lower Cretaceous limestones of reef and near reef facies. They include red algae belonging to the families Solenoporaceae, Permocalculus, and Corallinaceae, and numerous green algae of the family Dasycladaceae. The flora is quite similar to the Lower Cretaceous flora of the Mediterranean region, with identical or closely related species. A number of the genera are recorded for the first time in the western hemisphere.

JONES, DANIEL J., Earth Science Curriculum Project, Boulder, Colorado

THE GILBERT-TYPE DELTA

The pioneer work on deltaic sedimentation was done by G. K. Gilbert, and published in 1885, and in 1890. The deltas which he so carefully described were built into Pleistocene Lake Bonneville by rivers and creeks entering the lake from the rugged Wasatch Mountains to the east.

The classic concept of the delta, as described by Gilbert, with distinctive topset, foreset, and bottomset beds, was incorporated into virtually all introductory textbooks in geology as a "typical delta". It has persisted until the present time, in spite of the publication of many papers on such complex deltas as the Mississippi and others quite different from Gilbert's classic, simple delta. The term "Gilbert-type Delta" was introduced by Bates in 1953, who characterized it as a product of homopycnal flow into lakes.

The present paper describes and illustrates some typical Lake Bonneville deltas as "Gilbert-types". The deposits are characterized by:

1. Essentially homopycnal flow by mountain streams of steep gradient into deep water immediately offshore.
2. Occasional turbidity flows (mud-rock flows); hyperpycnal.
3. Coarse gravels dominant in foreset beds.
4. Fine sands and silts in bottomset beds, prograded locally to receding lake levels.

Some of the deltas of Lake Bonneville are true counterparts of alluvial fans.

JORDAN, WILLIAM M., Jersey Production Research Co., Tulsa, Oklahoma

REGIONAL ENVIRONMENTAL STUDY OF THE NUGGET AND NAVAJO SANDSTONES

The late Triassic (?) and early Jurassic Nugget and Navajo Sandstones are compositionally and texturally extremely homogeneous, mature, feldspathic quartz arenites extending over vast areas in the western United States. They have long been considered as "classic" eolian sandstones, deposited within great interior, sub-tropical deserts.

Trough-shaped "festoon" cross-bedding predominates and most tabular-planar and irregular sets are modifications of this basic style. Large scale simple, non-erosional, wedge-shaped sets are subordinate and are restricted stratigraphically to the middle part of the Navajo Sandstone. Ripple-marked, wavy-bedded horizons, plane bedded to massive units, and interbedded shale seams associated with horizontal truncation planes are common, particularly in the Nugget Sandstone. Very thin dolomitic carbonate lenses are found in the Navajo. While the sparse fossil evidence is indicative of terrestrial, but not necessarily extremely arid, conditions, these latter characteristics, as well as widespread contortion of inclined laminae, are apparently the result of subaqueous processes.

Sediment volume, grain composition, and textural maturity indicate a predominantly sedimentary provenance. No areal compositional or textural trends are discernible but south- and southeasterly-directed paleocurrents suggest a source along the western margin of the Canadian Shield. Textural parameters derived from recent sediments fail to clearly identify Nugget and Navajo depositional environments.

The weight of all evidence indicates, however, that both formations consist of a complex of shallow-marine, littoral, and coastal dune deposits laid within, and in advance of, an east- and southward-transgressing sea. Although temporary regressive oscillations may have at times exposed large areas of marine sand to the action of the wind, the majority of Nugget and Navajo cross-bedding is aqueous in origin. Preserved eolian structures are remnants of transverse coastal dune belts rather than vast interior dune fields, and paleocurrent directions thus reflect both marine current and coastal on-

shore wind regimes rather than simply ancient trade-wind circulations.

KANES, W. H., WELLS, DANA, and DONALDSON, A. C., Humble Oil & Refining Company and West Virginia University, Morgantown, West Virginia

FACIES AND DEVELOPMENT OF THE COLORADO RIVER DELTA IN TEXAS

The Recent delta of the Colorado River of Texas is an exceptional model for analyzing sediment and faunal facies relationships in environments associated with deltas. The small size facilitates close spacing of cores, which is essential for detailed facies study. Moreover, air photographs document growth from a straight shoreline to the lobe-shaped deposit formed prior to artificial channeling of the river through a barrier island. A 6- to 8-foot platform of deltaic sediments now dividing Matagorda Bay was deposited in the 6-year period following removal of an upstream log jam. Continuous cores (37) and borings (10) define the deltaic facies complex and the underlying bay facies in the northeastern section of the delta, which is unaffected by man-made modifications. The sequence of facies and environments encountered in sections is: (1) clays and silts, 1 to 5 feet thick (delta plain), including silty clays and clayey silts containing numerous roots and root fibers (natural levee), silty clays with roots and "algal-encrusted" washout pans (marsh), and massive to burrowed clays (channel fill); (2) sands, 2 to 6 feet thick, characterized by small-scale cross bedding (delta front); (3) laminated silty clays, 1 to 5 feet thick, (prodelta); and (4) laminated to burrow-mottled silty clays and clayey silts, 10 to 14 feet thick (bay), unconformably overlying the Pleistocene. Color variations, together with X-ray radiographs recording laminations and burrows, are most useful in distinguishing between facies of bay and prodelta environments. The faunas characterizing the facies and environments are more variable than are the sediments; juvenile and small mollusks, less than $\frac{1}{4}$ inch, seem more definitive of environments here than do Foraminifera. Subsidence of 1 to 2 feet is estimated to have resulted from compaction since initial delta formation.

KELLER, GEORGE H., U. S. Navy Oceanographic Office, Washington, D. C.

SEDIMENTARY ENVIRONMENTS IN THE MALACCA STRAIT, MALAYSIA

A preliminary investigation of the waters and bottom sediments in the Malacca Strait and southern portion of the Andaman Sea, made in 1961 by the U. S. Naval Oceanographic Office, resulted in the collection of bottom sediments as well as salinity, temperature, and water transparency observations at 23 stations. Current measurements were made at selected sites for periods of 24 to 36 hours. The Malacca Strait is a narrow, shallow passage between the Malay Peninsula and Sumatra with oceanographic and bottom sediment characteristics closely related to the strong currents, debouching rivers, climatic variation, and the close proximity of bordering land masses. The strait assumed its present configuration as a result of the post-glacial rise of sea level which drowned the Sunda Shelf. The strait is located in a typical equatorial climate although the monsoonal effects are not as severe as in the more open neighboring areas. A northwest current flow, essentially tidal, prevails in the strait throughout the year, and is largely responsible for the hydrographic and oceanographic conditions in the area. Surface salinities

and temperatures are found generally to be lower than in the surrounding seas. A wedge of cold, high-salinity bottom water extends from the Andaman Sea into the Strait. Bottom sediments primarily consist of muddy sands, with large areas of mud occurring in the vicinity of debouching rivers and in the Andaman Sea Basin. Calcium carbonate, primarily in the form of mollusk shells and foraminiferal tests, and organic carbon are found only in minor amounts in the Strait. Higher concentrations of calcium carbonate generally were associated with the finer sediments of the Andaman Sea, while the higher concentrations of organic carbon were found in the vicinity of debouching rivers. The non-calcareous detrital fraction is dominated by quartz, with minor amounts of orthoclase and plagioclase feldspars. The heavy mineral suite is complex due to the varied geology of the bordering land areas. The heavy minerals present consist primarily of leucoxene, ilmenite, magnetite, biotite, and amphiboles.

KLEIN, GEORGE DeVRIES, Department of Geology, University of Pennsylvania, Philadelphia, Pa.

PALEOCURRENTS AND OCEANOGRAPHY

The rationale of paleocurrent studies suggests that directional current structures identify the provenance, transport direction, basin architecture, and paleoslope of sedimentary sequences. Numerous examples from fluvial deposits have reinforced this view.

In marginal marine environments, directional current structures are fashioned by a combination of tidal, wave, and wind-driven currents which flow randomly with respect to slope and sediment source. Directional structures formed by these currents are oriented perpendicular or parallel to both slope and sediment source. The resulting structures, therefore, reveal only the direction of transport of the last current to act upon the sediments prior to burial.

The direction of flow of bottom-scouring currents in continental shelves and deep ocean basins is determined by changes in density, temperature, and salinity, by wind action, and by the rotation of the earth. Such changes tend to drive currents in random directions in the ideal case, but because of the earth's rotation, most ocean currents flow parallel to topographic strike. Resulting directional current structures of shelf and deep marine sediments cannot define paleoslope or provenance, but they do define topographic strike.

Because the direction of flow of ocean currents depends on many variables, several combinations of current systems can be recognized, including converging systems, diverging systems, and stratified systems. Where two currents converge, one system will override the other system because of differences in density. Converging directional data have been reported from ancient turbidites, indicating that similar converging current systems existed in the past. These data also suggest that many "turbidite" directional criteria were formed by bottom scouring ocean currents.

KNOPOFF, L., Department of Physics and Institute of Geophysics, University of California, Los Angeles, California

ATTENUATION OF SEISMIC WAVES IN THE MANTLE

Field and laboratory measurements of attenuation in metals, non-metals, and rocks over a wide range of frequencies indicate that the specific attenuation factor, $1/Q$, is substantially independent of frequency in homogeneous material, whereas it varies as the first power of frequency in liquids. This suggests that the mechanism