

definition of carbonate sediments to include those containing more than 50 per cent carbonate minerals, many new parameters can be studied and new and useful information can be incorporated in studies of carbonate rocks.

It is obvious that, by enlarging the definition, consideration of insoluble residues and stratigraphic associations is possible. Further, the geographic range of carbonate sediments is enlarged, permitting study of a wider variety of environmental conditions. Finally, a greater variation in thickness of Recent carbonate sediments is found, a fact which should allow a better understanding of sedimentation rates.

CIFELLI, RICHARD, Smithsonian Institution, Washington, D.C.

AGE RELATIONS OF MID-ATLANTIC RIDGE SEDIMENTS

Age relations of Mid-Atlantic Ridge sediments were established by means of planktonic Foraminifera in several selected areas between 22° N. lat. and the equator. Sediment types containing planktonic Foraminifera include unconsolidated ooze, consolidated ooze ranging from loosely friable aggregates to hard limestone, breccia, and palagonite-rich rock.

In the 22° N. area, consolidated sediments ranging in age from late Miocene to probable middle Pliocene were dredged from the flank of the ridge. Associated with the consolidated sediments in the dredges were basalts. Where an age relation could be inferred, the basalt overlies the late Miocene. Studies of Sr/Rb isotopes in the basalts are still in progress at M.I.T. Results thus far, however, indicate that these isotopes do not permit an age assignment. On the crest of the ridge there is no evidence of the presence of sediments older than Quaternary. Materials examined from the crest of the ridge include indurated detrital tuff and palagonitic rock.

Examination of sediments collected from the 11° N. area, Romanche trench and St. Paul's Rocks, are still in progress. In the Romanche trench, late Tertiary planktonic Foraminifera mixed with Quaternary assemblages were recovered from foraminiferal ooze in a core. In the St. Paul's Rocks area, the matrix of a conglomerate containing pebbles of the St. Paul's Rocks type yielded a Pleistocene foraminiferal assemblage. In the same area a vesicular basalt was filled with limestone. The limestone in the vesicles contained a mixed assemblage of late Tertiary and Pleistocene.

It is difficult to arrive at definite conclusions on the basis of the relatively little material thus far examined. Yet, it is interesting to note that the oldest assemblages recognized are late Miocene. Though not conclusive, the present data agree with other evidence suggesting a relatively young tectonic and volcanic history for at least parts of the Mid-Atlantic Ridge.

COLBURN, IVAN P., Department of Geology, California State College at Los Angeles, Los Angeles, Calif.

GRAIN FABRICS IN TURBIDITE SANDSTONE BEDS AND THEIR RELATION TO SOLE-MARK TRENDS ON SAME BEDS

Grain orientations were determined quantitatively for 62 turbidite sandstone beds that exhibited sole marks (19 beds with flute casts and 43 beds with groove casts). Statistically significant alignment of elongate grains within these beds tend to parallel the trends of the sole marks on the base of the beds.

Grain fabrics of six beds were determined quantitatively for multiple levels (a minimum of four levels in each bed). The mean grain trends were observed to parallel approximately the trend of the sole marks on the base of these beds.

Deviations in the mean grain trend from the trend of the sole marks were noted for most of the beds studied, but the mean deviation for all beds was less than 10°. Moreover, there was no systematic sense of deviation such as that reported by Bouma (1962), Spotts (1964), and Spotts and Weser (1964). The deviations were grouped in normal fashion about the zero deviation point.

Dip direction of imbricate grains was determined at three levels in one of the sandstone beds studied. The imbricate grains in the middle and lower intervals dip in one direction whereas those in the upper interval dip in the opposite direction. However, statistically significant data were obtained only from the lowest interval in the bed. The current sense indicated by the dip direction of the imbricate grains in the lower and middle intervals is the same as that inferred from the flute casts on the base of the bed.

Grain fabric exhibited in the rock slices was determined according to the technique described by Spotts (1964). The statistical technique used to evaluate the grain trends was patterned after the method described by Curray (1956). The writer and his assistant, Monty Hampton, devised the statistical technique used to evaluate the dip direction of imbricate grains.

COLEMAN, JAMES M., and CLARA HO, Coastal Studies Institute, Louisiana State University, Baton Rouge, La.

EARLY DIAGENETIC CHANGES IN FRESH-WATER CLAY DEPOSITS

Cored borings of Recent fresh-water swamp deposits that accumulated in the Atchafalaya River basin, Louisiana, revealed the existence of numerous early diagenetic changes in clay deposits and the formation of various types of syngenetic and epigenetic inclusions. The deposits, approximately 100 feet thick, range in age from contemporary to slightly greater than 10,000 years and were deposited in four major environments of deposition: poorly drained (stagnant) swamp, well-drained swamp, lacustrine, and lacustrine delta fill. Many of these environmentally controlled facies are repeated several times in a single vertical sequence and offer the unusual possibility of studying diagenetic changes at different stages of development within a particular environment. The most common diagenetic change is the replacement of plant rootlets and other organic fragments by pyrite and calcium carbonate. Pyrite replacement is most common in the poorly drained swamp sediments, whereas carbonate replacement occurs most commonly in the well-drained swamp deposits. These changes took place rapidly, probably within a few years after deposition. Both pyrite and calcium carbonate tend first to be formed in the open spaces within the organic fragments, second to invade the pore spaces, and last to replace most of the original organic material. Vivianite ($\text{Fe}_3\text{P}_2\text{O}_8 \cdot 8\text{H}_2\text{O}$) forms fairly rapidly and also replaces organic material.

Nodules are abundant throughout the section, but are more common in lacustrine sediments. The size of the nodules differs, generally being smaller in the younger units, and increases in size in the older units. The shape changes, ranging from round, flattened, lenticular masses (lacustrine) to round and irregular-