

1. Know the goals—of your company or your client;
2. Strive for good ideas—they communicate better than poor ones;
3. Improve your communication mechanics—understand vocabulary, use meaningful illustrations, and organize any presentation emphasizing salient points;
4. Know the weak points and alternate solutions, and
5. Try to communicate.

The Toastmasters organization, though commonly considered to be only preparation for public speaking, actually trains a man to organize his thoughts and present them clearly with poise and enthusiasm in any communication situation.

The above approach should enable geologists to perforate or bend, if not shatter, communication barriers.

R. A. RUDKIN, Imperial Oil Ltd., Edmonton, Alberta

LOWER CRETACEOUS OF MONTANA, NORTH DAKOTA, AND CANADA

(No abstract submitted)

W. A. SEARS, JR., Independent Geologist, Casper, Wyoming, and JOHN J. SULLIVAN, Independent Geologist, Casper, Wyoming

TIERTARY FORT UNION FORMATION OF NORTHERN ROCKIES

The Tertiary Fort Union Formation of Paleocene age has yielded oil and gas fields of major significance in the northern Rocky Mountain area during the past few years. Notably, the LaBarge platform area on the west flank of the Green River basin in Wyoming is the "classic" example of productivity and an excellent subject area for stratigraphic and environmental study of the Paleocene. As of January 1, 1967, the fields on the platform had a cumulative production in excess of 43 million bbls of oil and 770 Bcf of gas, produced primarily from sandstones in the Fort Union Formation.

After the uplift and erosion of the Late Cretaceous Lance Formation in the platform area, Laramide orogenic movements began with gentle uplift and truncation which cut into the Cretaceous Hilliard Formation and established an unconformity surface which subsequently was folded in response to Laramide adjustments. Following these events, general Tertiary deposition began with the formation of coaly swamps. In the steeper synclinal parts of the Green River and other intermontane basins, large bodies of water developed and thick sequences of source-bed shales filled these troughs which approached marine conditions as inland seas. The significant Fort Union production that has been found to date in the northern Rockies is associated with a beach-type environment developed along the shorelines of these inland seas. Toward the end of the time of deposition of the Fort Union, these seas were filled with sediment and only the coaly sequence remained until additional orogenic movement folded and eroded these beds—in some places as deep as the older Fort Union-sea units. The Wasatch Formation of Eocene age was deposited on this unconformity surface.

The depositional and producing environment of the Fort Union in the LaBarge platform area is not

unique to the Green River basin of Wyoming. Similar conditions are evident in many other basins in the northern Rocky Mountains where little or no exploratory effort has been made to locate lucrative production similar to that found and continuing to be developed in the Green River basin.

DONALD C. SWANSON, Humble Oil Co., Oklahoma City, Oklahoma

PENNSYLVANIAN GEOLOGY OF WESTERN MID-CONTINENT

Pennsylvanian sediments deposited between the Sierra Grande—Frontrangia positive elements and the Central Kansas uplift—Nemaha ridge trend contain large amounts of hydrocarbons. In the Anadarko basin alone almost 2 billion bbl of oil, more than 20 trillion cu ft of gas, and up to 700 million bbl of gas liquids eventually will be recovered from Pennsylvanian reservoirs. Mokane—Laverne, Camrick, Hansford, Greenwood, Keyes, and Elk City are six major Pennsylvanian fields in the western Anadarko basin. They have up to 2 billion equivalent bbls of recoverable oil in traps ranging from anticlines to purely stratigraphic. The reservoirs range from carbonate banks to alluvial sandstone and are of Morrowan to Virgilian ages.

The Anadarko and Denver basins always have been considered to be separate geologic provinces. However, during Pennsylvanian time they were part of the same geological regimen and properly should be considered together. The Amarillo-Wichita Mountains, Sierra Grande uplift and Frontrangia comprise a tectonic rim with a common orogenic history during the Pennsylvanian. The distribution of facies and depositional environments from this rim toward the stable-shelf area is similar and closely related across a broad area covering eastern Colorado, western Kansas, western Oklahoma, and parts of Nebraska, Wyoming, and Texas.

Only the Anadarko basin part of this large area has been explored actively and most of eastern Colorado and western Kansas remains a relatively virgin area for Paleozoic exploration. Because the relations of facies and environments in the whole area are similar, because the basin structure and geologic history are closely related, and because the Anadarko basin has been the scene of prolific discoveries, it could follow that the Pennsylvanian potential of the relatively unexplored area of western Kansas and eastern Colorado is very good.

L. W. VIGRASS, Western Resources Consultants Ltd., Calgary, Alberta

GEOLOGY OF CANADIAN HEAVY OIL SANDS

Obvious barriers that have been or could be broken by western Canadian heavy oil are in the fields of exploitation, transportation, and marketing. A geological barrier to be broken is the problem of the origin of the oil. The heavy oils appear to be at or very near the site where a discrete oil phase first was formed. An understanding of the origin of these vast accumulations of heavy oil conceivably would supply important clues to the origin of light crude oil, condensate, and gas.

There are about 750 billion bbl of heavy oil in place in western Canada and most of this occurs in sandstones of the Lower Cretaceous Mannville Group.