

iting all the complexity of real processes and some of the control of the laboratory.

A variety of structures was observed along stream courses. With few exceptions, all of the structures resulted from infrequent floods and were developed during waning phases. In contrast, most of the bedding was deposited during maximum flood conditions. Erosional structures observed include various types of scours, rills, tool marks, microterraces, and mud balls. Structures produced by transportation and deposition were ripple marks (10 varieties), lineations (3 varieties), and imbrication. Postdepositional structures include desiccation cracks (6 varieties), surface markings, salt-crust features, algal mats, and sand volcanoes. Bedding types observed were horizontal parallel and discontinuous, small- and medium-scale trough cross-stratification, medium- and large-scale low angle wedge cross-stratification, inclined channel fill-bar complex, ripple stratification, high angle avalanche front cross-stratification, and convolute bedding. Within the channel, inclined bedding is dominant, but on bars, low angle cross-stratification is the most abundant type of bedding.

Several factors combined to control the occurrence of specific structures and bedding types. Stream velocity is the most important factor, but particle size, time, water depth, and local channel irregularities are also significant.

PRAY, LLOYD C., Dept. of Geol. and Geophys., Univ. Wisconsin, Madison, WI 53706

SUBMARINE SLOPE EROSION ALONG PERMIAN BANK MARGIN, WEST TEXAS

Recent sea floor investigations, such as those of the western Atlantic-Blake Plateau region by Heezen and Stetson and coworkers, indicate extensive erosion by deep ocean currents on flat surfaces and on steeper submarine slopes. Evidence of small-scale submarine erosion of lithified crusts or skeletal material is increasingly being recognized, but few examples of ancient unconformities are known that represent appreciable erosion in a deep marine environment. Forming the 2 major unconformities of the Leonardian and Guadalupian strata of the Guadalupe Mountains of Texas required extensive erosion; that this erosion and the creation of numerous small channels and diastems in associated strata occurred in a continually submerged, relatively deep, marine environment best fits the present geologic data.

The 2 unconformities are roughly parallel. They form the upper and lower boundaries of the dark basinal carbonates of the Cutoff Formation. The lower unconformity, disconformable shelfward, steepens basinward to 5–10° and truncates over 200 m of essentially flat-lying Victorio Peak bank margin carbonates before flattening basinward. This unconformity probably inherited the relief of a gentler bank margin depositional profile. Erosion steepened and caused bankward slope retreat for possibly hundreds of meters. The upper unconformity locally intersects the lower and is overlapped by about 300 m of the Brushy Canyon Formation, composed largely of deep water sandstones.

These unconformities and associated diastems are generally smooth, sharp surfaces, showing scant evidence of the environment of erosion, but broad gentle undulations, and spoon-shaped, possibly closed depressions, some with tens of meters of relief, appear characteristic of both unconformities. Several V- to U-shaped basin-trending channels are present locally.

The mid-Permian bank margin of the classic Guadalupe Mountain area appears to furnish an outcrop analog for some of the processes of both deposition and erosion along deep submarine slopes. The possibility that deep marine erosion created many unconformities of the geologic record should be considered seriously.

RACKLAY, RUFFIN I., Consulting Geologist, Casper, WY 82601

SEDIMENTARY ENVIRONMENTAL CONTROL OF SANDSTONE-TYPE URANIUM DEPOSITS

Uranium deposits of the Rocky Mountain area are present in fluvial sandstones that were derived from granitic highland source areas and deposited in adjacent intermontane basins. The sandstones are arkosic to arkose in composition and contain a small percentage of carbonized plant debris and pyrite, medium grained to conglomeratic in size, subrounded to angular in shape, and are poorly to moderately sorted. The sandstones intertongue and are interbedded with green mudstone.

The host sandstone sequence represents a change in tectonic and/or climatic conditions. A local or regional unconformity is present at the base of the sequence and the character of the rocks is commonly substantially different. The climate was temperate and humid, and the groundwater level was very near the surface. The oxidation-reduction potential was low, which caused the carbonization of woody debris and the formation of pyrite, a condition also favorable for the precipitation of uranium and the other commonly associated metals. The depositional pattern of the aggrading streams was influenced by both tectonic and topographic elements of the area being buried.

Following deposition and burial further tectonic events induced changes in the hydrodynamic system which caused an invasion of the reduced sediments by an oxidizing cell that obtained much of its energy from the oxidation of the pyrite. The oxidizing cell mobilized the uranium and other susceptible elements and swept them ahead. Where the cell encountered resistance from variations in thickness of the sandstone units its progress was impeded, therefore areas of pinchouts or intertonguing of sandstone and mudstone tend to be favorable areas for the localization of uranium mineralization.

RAINWATER, E. H., Tenneco Oil Co., Houston, TX 77001

LOWER CRETACEOUS ROCKS IN GULF COASTAL PLAIN OF UNITED STATES

Rocks of Early Cretaceous (Comanchean) age underlie most of the Gulf coastal plain. Their thickness varies from zero at the updip subsurface pinchout in the eastern region and less than 2,000 ft at the outcrop in central Texas to about 7,000 ft at the downdip limit of control.

Carbonates make up most of the section west of the East Texas embayment; alternating carbonates, terrigenous clastics, and some anhydrite compose the section in eastern Texas and adjacent parts of Louisiana and Arkansas; and sandstone and shale, with some beds of limestone and anhydrite, represent the Lower Cretaceous in the eastern region. The percentage of carbonates increases seaward in all segments of the Gulf Coast.

Deposition was in shallow open-sea to restricted marine, deltaic, coastal-interdeltaic, and coastal-plain