

QUARLES, M., JR., Petty Geophys. Engineering Co., San Antonio, Tex.

INTERPRETATION OF UNUSUAL GEOLOGIC FEATURES ON CONTINENTAL SLOPE OFF LOUISIANA

Many complex structural features exist on the continental slope off eastern Louisiana that appear to have no land equivalents. Highly contorted near-surface structures overlie complex patterns of deposition. Salt or shale masses apparently intrude into sediments in horizontal planes as well as vertically. The total assemblage may account for new types of oil fields that may be very prolific.

A technique of analysis called "comprehensive interpretation" is used to illustrate structural anomalies. It involves carrying the entire geologic section in color-coded display of the 10-20 most reliable reflections available.

QUARLES, M., JR., Petty Geophys. Engineering Co., San Antonio, Tex.

SEDIMENTARY STRUCTURE ACROSS BOTTOM OF GULF OF MEXICO

A continuous profile shot across the abyssal depths of the Gulf of Mexico shows sedimentary features that could constitute stratigraphic traps for hydrocarbons. Faulting is present but of minor importance. The edges of the abyssal plain are abrupt and difficult to explain. Continuous correlation of beds across the section demonstrates these structural features.

QUIGLEY, M. D., Exploration Surveys Inc., Dallas, Tex.

PROGRESS IN APPLYING POTENTIAL FIELDS TO SEDIMENTARY SECTION

In 1964 a major oil company research laboratory set up a long-range program to improve the usefulness of potential field surveys in reconnaissance exploration. Investigations of field techniques, instrumentation, and interpretation were parts of the long-range program. The results of the program include the adaptation of the helium magnetometer to an air-borne instrument, improved location and data density control, and innovations of interpretation methods.

Specific application of gravity and magnetics to particular exploration problems is used to show that positive definition of subsurface structure can be resolved when proper field techniques and interpretation procedures are used. Exploration Surveys Inc. has incorporated these developments and procedures into its surveys with surprising results. The high sensitivities and location controls now possible allow a direct comparison of gravity and magnetics surveys. The comparisons lead to the determinations of lithologies, depositional environments, and age relations, as well as the delineation of subsurface structure.

Given highly sensitive data and accurate location control, the interrelation of gravity and magnetic anomalies can be used by the geophysicist and the geologist to eliminate many of the ambiguities normally attached to the interpretation of potential fields.

RASCOE, B., JR., Phillips Petroleum Co., Bartlesville, Okla.

LATE MISSISSIPPIAN EPEIROGENY AND THE WICHITA OROGENY IN THE MID-CONTINENT

Widespread epeirogeny in Late Mississippian time uplifted elements of the Transcontinental arch and bordering regions. Late Mississippian deposition in the Mid-Continent was confined to the Anadarko, Ardmore, and Arkoma basins where maximum thicknesses of 5,000, 3,000, and 2,000 ft of sediments (respectively) accumulated. Erosional processes accompanied regional uplift and continued after the cessation of Late Mississippian deposition. Consequently, over the Mid-Continent an erosional surface was etched on a terrane of predominately carbonate Paleozoic sediments and igneous Precambrian rocks.

A maximum thickness of 4,000 ft of Morrowan (Early Pennsylvanian) sediments was deposited in the Anadarko basin. From there, northward to southwestern Nebraska, Morrowan beds onlap the Late Mississippian erosional surface and truncate rocks ranging in age from Late Mississippian to Precambrian. This depositional onlap accounts for the limits and much of the thinning of the Morrowan Series from the Anadarko basin toward the north and northeast. Morrowan rocks on the order of 3,000 ft and 5,000 ft in thickness were deposited in the Ardmore and Arkoma basins, respectively. In these basins Morrowan sediments lie on Late Mississippian rocks with minor unconformity.

The post-Morrowan, pre-Atokan Wichita orogeny brought the aligned Criner Hills-Amarillo-Wichita uplift-Cimarron arch into existence through strong vertical uplift accompanied by high-angle faulting with an estimated 10,000-15,000 ft of maximum vertical displacement. Concurrently a foredeep began to form in the Arkoma basin which would subside rapidly and receive phenomenal thicknesses of Atokan rocks—12,000 ft in the western Arkoma basin and 19,500 ft in the eastern part of the basin.

Meanwhile, an area on the western margin of this developing Atokan foredeep became mildly positive, possibly in response to the strong downwarping of the foredeep. From the Criner Hills this area trends northward across the Ardmore basin and the Arbuckle uplift, and turns northeastward toward the Ozark uplift, a distance of approximately 175 mi. This belt was about 125 mi wide and it encompassed the southeastern flank of the Anadarko basin and the northwestern flank of the Arkoma basin. Morrowan rocks were eroded from this positive area with the exception of those in the Ardmore basin, those on the southeastern flank of the Anadarko basin, and those on the northwestern margin of the Arkoma basin. These rocks exhibit an angular, unconformable relation with overlying Atokan or Desmoinesian beds. Elsewhere in the Arkoma basin, and in the Anadarko basin and on its shelf area, the contact of Morrowan and Atokan rocks is nonangular and probably disconformable.

In western Oklahoma, western Kansas, and southwestern Nebraska, the depositional limit of Morrowan sediments is overstepped eastward by Atokan beds, and the depositional limit of Atokan beds is in turn overstepped eastward by Desmoinesian rocks. The uninterrupted advance of these sediments eastward toward and onto the Nemaha ridge and the Cambridge arch-Central Kansas uplift is evidence that the age of these features is pre-Morrowan (Late Mississippian) rather than post-Morrowan, pre-Atokan (Wichita orogeny) as often claimed. In addition Atokan and Desmoinesian rocks of these areas do not exhibit lateral facies changes involving the influx of coarse clastic sediments