

and Sacate Formations. That this westerly shift continued is documented by Oligocene (Sespe) isopach trends and paleocurrent-slope indicators. In late Paleogene-early Neogene time, Vaqueros sands, unconformably overlying the nonmarine Sespe, were winnowed over time-persistent, east-west structural highs.

The culmination of the development of the western Transverse Ranges is marked by middle Miocene rifting, foundering, and volcanism. Deep-basin sedimentation essentially began with this event. The synclinal distribution of late Miocene turbidite sands is evidence for the continued growth of the early formed structural highs.

Late Neogene sediments within the Santa Clara graben offer a classical example of structurally controlled sedimentation. Pleistocene and Holocene submarine-fan deposits reflect the impingement of the westward-plunging Montalvo anticlinorium. Contrasting northern and southern shelf deposits of similar age also are controlled by early formed structural trends. Many of these structures are the oil and gas fields of the modern Santa Barbara basin.

FISHER, W. L., Bur. Econ. Geology, Univ. Texas, Austin, Tex.
**DELTAIC SEDIMENTATION, SALT MOBILIZATION,
 AND GROWTH FAULTING IN GULF COAST BASIN**

Regional facies mapping and analysis of thick, terrigenous wedges, and mapping of principal structural features (salt intrusion and growth faulting) indicate a direct relation between sedimentation and tectonics in the western Gulf basin. Proximal parts of clastic wedges infilling the basin consist of paralic, mainly deltaic, depositional systems; distal parts are made up largely of deep-water, continental-slope deposits. Two kinds of delta systems are characteristic: (1) high-constructive deltas marked by rapid, large-volume deposition; and (2) high-destructive deltas with large sand content and slower rates of accumulation.

Distribution of 4 major high-constructive delta systems and associated continental-slope systems is coincident with 4 major salt-diapir fields (including about 90% of the domal salt structures of the basin). Mobilization of deep seated, bedded salt occurred by lateral migration to interdeltic areas and by distal migration, establishing diapir fields fronting major delta systems and coincident with slope systems. Salt mobilization related to strike-depositional (barrier bar and strandplain) systems was generally as broad salt ridges rather than domes.

In high-constructive deltas, growth faulting is facies coincident, forming at the boundary of delta-front sands and thick prodelta muds. As with salt mobilization, the principal tectonic grain of growth faulting developed in connection with the 4 major episodes of high-constructive wedges in the offlap filling of the basin, rejuvenating growth faulting and salt mobilization initiated by the high constructive deltas. Accordingly, direct facies correlation in distribution of growth faulting and salt mobilization occurred.

FORRESTER, J. D.

HOW MINING INDUSTRY CONSIDERS THE ENVIRONMENT

No abstract available.

FOSTER, NORMAN H., Trend Exploration Ltd., Denver, Colo.

**PRACTICAL APPLICATION OF REMOTE SENSING TO
 PETROLEUM AND MINERAL EXPLORATION**

The symposium papers concern many aspects of the application of remote sensing to natural resource exploration. Remote sensing is best discussed in the context of the electromagnetic spectrum. Any device which senses a part of the spectrum from a remote position may be classified as a remote sensor. Visible light is the most familiar part of the spectrum, and our eyes and

a camera with its film are remote sensors. In the shorter wavelengths, ultraviolet light has been useful in limestone-dolomite studies. X-ray diffraction patterns long have been used in mineral identification. In exploration for radioactive minerals, sensing parts of the gamma-ray spectrum with airborne or ground scintillometers is a common technique. On the longer wavelength side of the spectrum, near and far infrared are adjacent to visible light. Near or photographic infrared, when taken with a low sun angle, has been very useful in mapping faults and other features which are topographically expressed. Far, or thermal, infrared is also useful in mapping faults. Because water has a tendency to be more abundant in fault zones, they commonly are expressed as slight temperature lows due to the cooling effect of evaporation.

Side-looking-radar imagery also is very useful in fault delineation and for general mapping purposes, particularly in areas where cloudiness is a problem in conventional aerial photography. Side-scan sonar is a technique used to obtain underwater imagery, and should prove useful in bathymetric studies of shelf areas.

FRANKS, S. G., and J. HELWIG, Dept. Geology, Case Western Reserve Univ., Cleveland, Ohio

PETROLOGY AND SEDIMENTATION OF EARLY PALEOZOIC ISLAND-ARC DEPOSITS, NEWFOUNDLAND

The Cambro-Ordovician New Bay Formation (2,000 m) of the Newfoundland Appalachians is a southward-thickening prism of conglomeratic, volcanogenic turbidites interbedded with sandstone and black argillite. Abundant sedimentary structures include tool and scour marks, cross-lamination, normal and multiple graded bedding, slump folds, and channels. The top of the formation is mostly agglomerate interbedded with red-and-green laminated silty argillite. Trails and burrows are the only fossils.

The sandstones are lithic-feldspathic arenites and basaltic wackes composed of fresh, angular unstable mineral grains and little quartz. The conglomerates have volcanic and sedimentary clasts enclosed in an ash matrix altered to chlorite. Volcanic clasts are basalt, andesite, and dacite porphyry. Sedimentary clasts include green cherty siltite, fine-grained feldspathic arenite, abundant rip-up clasts of argillite, and rare jasper.

Analysis of Bouma sequences, amalgamated contacts, and trace fossils indicates abrupt regression late in New Bay deposition. Sole marks, slump folds, and channel axes indicate marginal sediment supply from turbidity currents and slumping down an east-facing submarine slope.

On the northwest the formation thins and apparently passes into marine and subaerial pyroclastic strata of the Wild Bight Group. On the east it is in sedimentary and tectonic contact with the Dunnage mélange which has been interpreted as an ocean trench deposit. The petrology and sedimentology of the New Bay Formation compares closely with certain Miocene volcanic arc deposits described from Malekula Island, New Hebrides. Therefore the New Bay is interpreted as a volcano-genic "apron" deposited in the arc-trench gap of a Cambro-Ordovician island arc.

FRASER, GORDON S., and NORMAN C. HESTER, Illinois State Geol. Survey, Urbana, Ill.

STARVED ROCK MEMBER OF ST. PETER SANDSTONE—OFFSHORE SHOAL

The Starved Rock Member of the St. Peter Sandstone has been described as a clean, medium-grained sandstone that was deposited in a broad band across northern Illinois. This deposit displays a sequence of sedimentary structures that indicates deposition in shallow water. From the base upward the structures include (1) massive beds; (2) small-scale trough cross-strata; (3) high-angle tabular cross strata; (4) large-scale trough

cross strata; (5) alternating beds of horizontal, burrowed laminae and small-scale trough cross strata; and (6) massive and burrowed beds.

Tabular cross strata and small-scale trough cross strata show a consistent northwest orientation, whereas the large-scale trough cross strata show no preferred orientation. Both tabular and large-scale trough cross strata consist of medium-coarse sand, whereas the small-scale trough cross strata and horizontal laminae are of medium sand. Discontinuous horizontal laminae, present at irregular vertical intervals in the section, consist of interlaminated medium to fine sand and silt.

Recently, the sand waves of the North Sea have been proposed as an analog for the St. Peter environment in Wisconsin, because of the presence of such sedimentary structures as very large-scale tabular cross strata. However, the sedimentary structures, directional data, and textures observed in the Starved Rock Member resemble more closely those developed on nonbarred coasts, like that of Oregon. The presence of inferred deeper water sediments within the St. Peter, landward of the Starved Rock Member, indicates that this sand body may have been deposited as an offshore shoal.

FREEMAN, TOM, Dept. Geology, Univ. Missouri-Columbia, Columbia, Mo.

TEMPORAL DOLOMITE-CALCITE SEQUENCE AND ITS ENVIRONMENTAL IMPLICATIONS

Where dolomite and calcite cements occur together, calcite is commonly the younger. This is true of cements that fill both primary (intergranular and intraskeletal) and secondary (vug and vein) pores. Rarely there is an alternation of cement types, but where this occurs, dolomite is still usually the penultimate, and calcite the ultimate.

In primary pores, the most likely unidirectional process producing the sequence, dolomite-calcite, is a change from marine to freshwater phreatic conditions through a lowering of sea level, elevation of the land, or increase in freshwater head.

In secondary pores, fluid inclusion studies of epigenetic dolomite-calcite show the dolomite to be both hotter and saltier than the associated younger calcite. Therefore, this sequence most likely records precipitation concomitant with erosional unloading, which should both reduce the geothermal environment and promote a freshening of the water.

FRIEDMAN, GERALD M., Dept. Geology, Rensselaer Polytech. Inst., Troy, N.Y., and JOHN E. SANDERS, Dept. Geology, Barnard College, New York, N.Y.

ASPECTS OF GEOLOGIC HISTORY AND MARINE GEOLOGY OF NEW YORK BIGHT

The New York Bight consists of two shelf sectors: (1) New Jersey shelf, trending NE-SW; and (2) Long Island shelf, trending east-west, which are separated by the Hudson shelf valley (extending southeast from outer New York Harbor to the Hudson apron at the shelf edge near the head of the Hudson submarine canyon). Although the present shoreline configuration of the bight results from the Holocene submergence, 2 important earlier developments include post-Miocene uplift of the Appalachians and especially of New England, and Quaternary glaciations.

During the Pliocene, uplift inland created the present regional dips of the coastal-plain strata and supplied sediment to fans which prograded a thin sheet of gravel across the marine coastal-plain formations. There followed a great drop in base level and deep cutting of major river valleys. The inner cuesta and adjoining inner lowland of the coastal plain, part of which is now filled by the water of Long Island Sound, were formed and large quantities of sediment built the shelf edge seaward. Erosion in submarine canyons was especially active at this time.

During the Pleistocene, in connection with continental glaciations, many changes of base level and climate occurred. During the Wisconsin Stage, ice flowing from about N20°W depos-

ited 2 prominent terminal moraines atop Long Island's coastal-plain cuesta—the Ronkonkama (older and more southerly) and Harbor Hill (younger and more northerly) moraines. During each deglaciation, wide outwash plains spread across parts of what is now the Long Island shelf and extensive lakes flooded by varved clays formed north of each moraine ridge. Rapid drainage of these moraine-dammed lakes through the Narrows (between Staten Island and Brooklyn) is thought to have been responsible for final erosion of the Hudson shelf valley and for major deposition on the Hudson apron.

As the sea transgressed across sandy outwash (Long Island shelf) and fluvial gravels (New Jersey shelf), it formed barrier chains and reworked the sediments. During parts of the submergence, some barrier chains retreated progressively landward, and the surf zone reworked all shelf sediments. At other times the barriers were drowned in place.

FRIEDMAN, GERALD M., AMIHAI SNEH, and ROY W. OWEN, Dept. Geology, Rensselaer Polytech. Inst., Troy, N.Y.

GENERATION OF LAMINATED GYPSUM IN SEAMARGINAL POOL, RED SEA

In the rock record, laminated gypsum commonly is interpreted as formed by precipitation from a body of brine. A study of a hypersaline pool at Ras Muhamad, the southernmost tip of the Sinai Peninsula, suggests an alternative interpretation of origin. Algal mats carpet a sea-marginal hypersaline pool, which has a salinity of 127‰ in the spring when the sulfate concentration has reached 9,700 mg/l; by mid-summer salinity has risen to 314‰ and, as a result of the precipitation of gypsum, the sulfate level has dropped to 4,570 mg/l. In the spring the SO₄/Cl ratio is almost identical with that of seawater (14.0 × 10⁻² for seawater; 13.9 × 10⁻² for the pool), but has dropped sharply by mid-summer as gypsum precipitates (2.4 × 10⁻²).

Algal mats not only survive, but are remarkably active at this high salinity. Gypsum precipitates as a meshwork between the mats and, depending on the algal growth, concentrates as crystals in laminae ranging in thickness from 1 to 2 cm. Compaction of such laminae as they become part of the rock record would result in much thinner laminae. Dark organic mats alternate with white layers of gypsum. Such a deposit in the rock record has, in the past, been mistaken for a deep-sea deposit.

Laminites of aragonite precipitated by algae alternate with laminae of gypsum. Oncolites, pellets, ooids, and spherulites are particles precipitated within the algal mats.

GEBELEIN, CONRAD D., Bermuda Biol. Sta., St. George's West, Bermuda

SEDIMENTOLOGY AND STRATIGRAPHY OF RECENT SHALLOW-MARINE AND TIDAL-FLAT SEDIMENTS, SOUTHWEST ANDROS ISLAND, BAHAMAS

Shallow-marine and tidal-flat carbonate sediments form a seaward-prograding wedge (maximum 6 m thick) on the southwest coast of Andros Island. Sediments have been deposited in a large (85 × 25 km), arcuate embayment in Pleistocene limestone. IncurSION of the sea over freshwater peat reached the inner, central margin of the embayment 5,000-7,000 years ago. Subsequent lateral progradation (maximum 30 km perpendicular to strike) of marine and tidal-flat sediments has occurred in several phases. 1. Continuous lateral progradation of a very shallow (less than 1 m), wide (3-5 km) subtidal platform adjacent to the shoreline. Sediments on the platform are poorly sorted, massive, fossiliferous, white-pellet muds with prominent filled burrows. 2. Development of 5 major shorelines, roughly parallel and 2-5 km apart, during the past 5,000 years. Each new shoreline developed, presumably by storm action, onto the subtidal platform, and isolated a shallow linear lagoon. Abandoned shorelines form parallel bands of discrete to coalesced,