

COMPARATIVE BIOLOGY AND ECOLOGY OF TROPICAL AND ANTARCTIC FORAMINIFERA

Studies of shallow-water benthonic Foraminifera are under way currently at Eniwetok Atoll in the equatorial Pacific and on the Antarctic Peninsula utilizing scuba gear for direct observation of natural situations and laboratory experimentation. On the atoll, species are distributed by microhabitat without regard for depth (to at least 150 ft), whereas in Antarctica, the benthic Foraminifera do not select specific microhabitats. Instead they are zoned according to depth, mostly in association with changing macrofaunal and floral changes. Because of ice abrasion in the Antarctic, few or no Foraminifera live between the intertidal zone and 18 ft in depth. Below that, a zone characterized by large kelps, sponges, tunicates, and brachiopods contains many Foraminifera, and at 120 ft the association is dominated by large glass sponges with about 25 species of Foraminifera.

Heavy predation on Foraminifera takes place on the atoll by fish and grazing invertebrates; in Antarctica, Foraminifera are fed on mostly by invertebrates. There are at least 3 nutritive strategies in reef Foraminifera, although they seem to feed largely on bacteria in Antarctica. Like some tropical species, some Antarctic ones seem to live a long time (years) before reproducing.

LISCO, RICHARD K., Teton Exploration Drilling Co., Inc., Casper, Wyo.

URANIUM AND ITS FUTURE ROLE IN NATIONAL ENERGY REQUIREMENTS

This paper is addressed generally to the use of uranium as an energy source for our future national energy requirements. It deals in part with the problem areas of; recent and pending legislation in the fields of health and safety, environment, public land laws, exploration, production, lead time between start of exploration and production of yellow cake, industry slippage, intervenors, and the effect of all of these on planning for the future. Uranium will play an important role not only in our future national energy requirements, but also in our balance of payments and national security.

LOHMANN, GEORGE P., Dept. Geol. Sci., Brown Univ., Providence, R.I.

STRATIGRAPHY AND SEDIMENTATION OF DEEP-SEA OCEANIC FORMATION ON BARBADOS, WEST INDIES

Detailed study of the Oceanic Formation on Barbados has revealed a record of pelagic sedimentation from earliest middle Eocene through latest Oligocene (20-50 m.y. ago). A succession of diverse lithologies and lateral facies changes within contemporaneous sediments is recognized. The succession consists of foraminiferal, radiolarian, and nannoplanktonic clays and marls, radiolarites, spiculites, diatomites, cherts, brown clays, and volcanic ash beds. All are eupelagic, deep-sea sediments. Of particular interest are (1) radiolarites and cherts which correlate with middle Eocene cherts shown by Deep Sea Drilling Projects to be widespread in both the Atlantic and Pacific; and (2) regular, periodic fluctuations of carbonate sedimentation rates and Foraminifera-nannoplankton ratios, in the upper Oligocene foraminiferal marls, which are similar to those frequently observed in Pleistocene foraminiferal oozes and attributed to climatic periodicities.

No evidence for progressive shallowing of the Barbados Ridge is apparent until after deposition of both the Oceanic Formation and the overlying Conset Marl. No *in situ* shallow-water sediments on Barbados are older than 10-15 m.y. This is consistent with the predicted arrival of the Caribbean plate into the present eastern Caribbean.

Middle and late Eocene sediments exhibit lateral facies changes. These are characterized by both a decrease and an increase in absolute sedimentation rates of carbonate and terri-

geneous clay, respectively, from north to south. The paleoslope determined from sedimentary structures deepens from north to south and suggests that the observed facies change may be attributed to increasing carbonate dissolution with depth. However, the accompanying increase in absolute sedimentation rate of terrigenous clays indicates that dissolution alone is an inadequate explanation. Either dilution by terrigenous clays from the south, or local sediment redistribution by currents and slumping, must have occurred.

LUCAS, PETER T., Shell Oil Co., Denver, Colo.

ALTAMONT—A MAJOR FRACTURED AND OVERPRESSURED STRATIGRAPHIC TRAP

Altamont field in the Uinta basin of Utah is a much-overpressured accumulation of high pour-point crude, producing from multiple, thin, Tertiary reservoirs in a 40-mi long stratigraphic trap. Postdepositional shift of the structural axis of the basin created an updip pinchout of low-porosity sandstones into a lacustrine "oil shale" sequence. Reservoir performance is enhanced significantly by vertical fractures and initial pressure gradients which sometimes exceed 0.8 psi/ft.

Field limits in large part are undetermined, and few dry holes have been drilled. Average well drilling and completion costs approach \$1 million. The initial potential of field wells ranges from 500 to 2,500 BOPD with an average GOR of about 1,000 cu ft/bbl. Matrix permeability generally is less than 0.2 md; therefore, high well performance is a function of fracture permeability. Inability to define reservoir parameters and oil in place causes individual well reserve estimates to depend mainly on pressure decline-cumulative relations.

Significant engineering problems are related to evaluation and completion of these thin, low-porosity, fractured and overpressured pay intervals, which span up to 2,000 ft of stratigraphic section in some wells. These problems are compounded by the problem of handling 110°F pour-point waxy crude. Inability to pipeline the crude conventionally from the relatively remote area has delayed full production of the field. About 40 drilling rigs are active in development of the trend, and a field potential is suggested of in excess of 250 million bbl with significant delineation of field limits yet to be accomplished.

LUMSDEN, DAVID N., Dept. Geology, Memphis State Univ., Memphis, Tenn.

DOLOMITE, LIMESTONE FACIES, AND INSOLUBLE RESIDUE—A RELATIONSHIP?

Almost 600 insoluble-residue and quantitative X-ray analyses have been performed on samples collected from the Callville and Pakoon Formations (Pennsylvanian and Lower Permian) of southern Nevada. The gross lithologies present are limestone, dolostone, and carbonate-cemented sandstone. The insoluble residues range from 0.7 to 77.0% and consist of fine-sand size quartz with varying amounts of illite (clay).

Plots of the total insoluble residue against percent dolomite show that more than 500 of the samples fall into either limestone (greater than 90% calcite) or dolostone (greater than 90% dolomite) end members. Plots of the dolomite percentage in the 100 remaining, mixed-calcite-dolomite mineralogy specimens versus total insoluble residue showed no obvious trends. Plots of the mixed-mineralogy specimens versus the illitic clay content likewise showed no obvious trends.

Five limestone facies (micrite, sparse micrite, packed micrite, biopelsparite and oosparite) were recognized. No relation was found between limestone facies and percent total insoluble residue or between percent dolomite and limestone facies.

LUNKING, W., and F. R. SIEGEL, Dept. Geology, George Washington Univ., Washington, D.C.; and J. W. PIERCE, Smithsonian Inst., Washington, D.C.

GEOCHEMISTRY OF A SEDIMENTARY PROCESS, GOLFO SAN MATIAS

Geochemical analyses of marginal basin sediments (Golfo San Matias, Argentina) for Ca, Mg, Sr, Co, Ni, V, Ti, Cr, Fe, Cu, Zn, and Mn in the total sediments and their carbonate fractions have yielded distinct halistases, the areal dispositions of which must be controlled by processes operating within the basin. One of the most important of these processes is derived from hydrodynamic forces moving the sedimentary particles and depositing them in rather distinct (at the 1-phi interval) granulometric zones.

For the total sample analyses, there are covariant relations between the dominant granulometric fractions and certain elements (for example, Zn with clay size, Ti with fine-sand size). For other elements, correlations are less definitive (for example, Fe).

Fragmented biogenic material is ubiquitous in the basin sediments but shows strong accumulations in some zones as a result of localized sources and dominant current activity; there is an expected positive correlation between Ca, Mg, and Sr, and the quantity of carbonate material in the sediments, and the distribution of Co appears to be directly influenced by biogenic components.

In the study basin, the disposition and shape of the plotted halistases are determined by the mode of migration of the elements, in solution, as hydrolysates, as resistates, and as biocates. Although homogenization drives exist, localized conditions have been more influential in controlling detrital and elemental distributions and dispersions. Selected elements are examined for their utility in serving as discriminants better to interpret similar data from the geologic column.

MACDONALD, HAROLD C., Dept. Geology, Univ. Arkansas, Fayetteville, Ark.

IMAGING RADAR—TOOL FOR PETROLEUM AND MINERAL EXPLORATION

Remote sensing methods have great potential application in geologic exploration for fuel and mineral resources. Unfortunately, many of the more exotic remote-sensing techniques are still in research and development stages, and most surveys must be conducted in the framework of experimentation rather than routine operation. Sidelooking radar (SLAR) is one of the exceptions to this overall categorization. SLAR systems, originally developed as all-weather military reconnaissance sensors, are providing extremely encouraging results in geologic exploration. Although the success of SLAR surveys has not been widely publicized, more than 6 million sq km of radar mapping has been completed during the past 3 years. Three commercial radar-mapping contractors have conducted geologic-reconnaissance surveys in some of the world's most inaccessible and remote terrain. Radar imagery is providing a first look at many cloud-shrouded regions in Brazil, Venezuela, Colombia, Panama, Nicaragua, Indonesia, and Australia.

The fine resolution of aerial photography is not presently available with imaging radars; however, they do offer the distinct advantage of a large swath of ground coverage (typically at least 20 km). This synoptic presentation allows the interpreter to become quickly familiar with the essential features of structural provinces. Minimal scale distortion allows stereoscopic interpretation on imagery strips that can be enlarged to at least 10 times the acquisition scale. Radar-mosaic construction has provided sufficient base-map information to anticipate and evaluate logistic problems to be encountered during seismic operations or when reconnoitering a territory for favorable drilling sites. Sidelooking radar, like any tool, has limitations as well as capabilities for petroleum and mineral exploration.

MARK, H. B., JR., Dept. Chemistry, Univ. Cincinnati, Cincinnati, Ohio

PROBLEMS IN CHEMICAL ANALYSIS OF ENVIRONMENT

Several recent comparative studies of analytical results and procedures cast serious doubt on the validity of many of the pollution data now being reported. These comparative studies indicate that both sampling techniques and analytic procedures are at fault. For analysis of trace metals in seawater and of oil spills, methods of obtaining valid or representative samples from the field and design of "foolproof" procedures for carrying out the subsequent chemical analysis have been developed, as have remote or automated monitoring devices for natural systems.

MARTIN-KAYE, PETER H. A., Hunting Geology and Geophysics Ltd., Boreham Wood, Elstree, England

GEOLOGY OF EASTERN AND CENTRAL NICARAGUA—INTERPRETATION OF SIDE-LOOKING RADAR IMAGERY

In late 1971, the entire country of Nicaragua was surveyed by side-looking radar for the production of a 47-sheet sequence of 1:100,000-scale mosaics. Interpretation of the imagery of the central highlands and eastward toward the Atlantic Coast has contributed substantially to the elucidation of the geology of this previously little-known region. Although little that is new has been added to the stratigraphic column, the distribution of the main stratigraphic units has been clarified, and the principal structural elements established. The work in Nicaragua is an additional example of the quality of side-looking radar for rapid regional geologic interpretation and consequent guidance of ground programs.

MATTHEWS, MARTIN D., Dept. Geology, Washington State Univ., Pullman, Wash.

AGGREGATE PARTICLES IN SEDIMENTS OF YUKON RIVER, ALASKA

A study of the particle size and mineralogic characteristics of bottom and suspended sediment from Kwikluak Pass of the Yukon River revealed that iron oxide and organic material act as binding agents. The binding agents appear to be equally responsible for the presence of aggregate particles which constitute a minimum of 14% of the sediment weight. The mean and standard deviation of the sediment in the aggregate state was $\bar{x} = 8.2$ phi and $\sigma = 2.0$ phi as suspended material and $\bar{x} = 5.4$ phi and $\sigma = 1.5$ phi as bottom material. These distributions are coarser and less variable than those measured after removal of iron oxide and organic binding agents: $\bar{x} = 8.6$ phi and $\sigma = 2.3$ phi as suspended sediment and $\bar{x} = 6.4$ phi and $\sigma = 2.6$ phi as the bottom sediment. The proportion of aggregates as a function of grain size increases from 10% at 4.0 phi to 50% at 12.0 phi. The size distribution of particles composing an aggregate grain is distinctly bimodal indicating the binding of fine particles to larger grains. However, aggregates also may be composed entirely of fine grains. There was no evidence of mineralogic selectivity within these aggregates. This supports a probabilistic mechanism of formation, whereby the mineralogy of the aggregates is controlled by the frequency of the mineral type in the discrete state. The similarity of aggregates in the bottom and suspended sediments supports the hypothesis that the aggregates were formed in the soil horizon within the Yukon River basin.

The presence of aggregate particles within river sediments can significantly alter conclusions concerning grain-size parameters and resulting mineral-transport characteristics unless proper analytical procedures are utilized.

MCBRIDE, E. F., Dept. Geol. Sci., Univ. Texas Austin, Austin, Tex.

BOULDER BEDS IN TESNUS AND DIMPLE FORMA-