

pears to be geothermally immature in both the onshore and offshore region and neither the Selma Group nor the Tuscaloosa is considered capable of generating producible quantities of either liquid or gaseous hydrocarbons within these areas.

BRYANT, W. R., Texas A&M, College Station, Tex., and J. M. COLEMAN, Louisiana State Univ. Coastal Studies Inst., Baton Rouge, La.

Subaqueous Landslides and Diapirs, Gulf of Mexico

No abstract available.

BUFFLER, R. T., F. JEANNE SHAUB, J. LAMAR WORZEL, Univ. Texas Marine Sci. Inst., Galveston, Tex., and JOEL S. WATKINS, Gulf Research and Development Co., Houston, Tex.

Gravity-Slide Origin for Mexican Ridges Foldbelt, Southwestern Gulf of Mexico

The geometry of a part of the Mexican Ridges foldbelt in the southwestern Gulf of Mexico as detailed by multifold seismic data suggests an origin by large-scale gravity sliding. Here, apparently competent, lower Tertiary-Holocene strata, probably sandstones and shales, have moved downslope along an underlying deformed zone probably consisting of weak and incompetent Upper Cretaceous-lower Tertiary geopressed shales. Gravity-generated horizontal compressive stresses apparently caused uniform folding and imbricate thrusting within the competent strata. This deformation is probably an important mechanism for overcoming resistance to basal sliding within the incompetent layer. Large-scale submarine sliding is also suggested by the grouping of the fold trends into two large lobate areas. Sliding apparently has continued periodically since about middle Tertiary time. This massive sliding may have been triggered by periods of uplift and seaward tilting and perpetuated by sediment loading in a large growth-fault system just landward of the main foldbelt along the upper slope.

BUFFLER, RICHARD T., and KENNETH J. MCMILLEN, Univ. Texas Marine Science Inst., Galveston, Tex.

Miocene Submarine Fans in Deep Western Gulf of Mexico as Interpreted from Seismic Reflection Profiles

Sequences of regularly spaced, seaward-dipping reflections seen on east-west seismic profiles along the lower slope of the western Gulf of Mexico are interpreted to be prograding clinoforms in the distal parts of large submarine-fan complexes. North-south lines define the lobate nature of the fans, some of which are over 50 km across. Some fans appear to be composites of overlapping, individual fan lobes. DSDP hole 90 recovered upper Miocene laminated, silty mud at the top and bottom of one of the prograding sequences, suggesting that any sand in the system must have been deposited farther upslope in the more proximal parts of the fan. Deposition of these sequences may have occurred during a major late Miocene lowering of sea level.

The late Miocene fans appear to be part of an overall fining-upward sequence along the lower slope beginning with the middle Miocene sandy turbidites and continuing through deposition of late Miocene silty clay laminates into Pleistocene deposition of hemipelagic and pelagic sediments. The sandy middle Miocene turbidites were derived from the Mexican mainland on the west and correspond to a zone on the seismic profiles characterized by chaotic and discontinuous reflections. These reflection patterns suggest a fan environment more proximal than the younger fans above—perhaps a suprafan environment with fan channels.

The overall fining-upward sequence and gradual cessation of turbidite deposition probably were due to the late Tertiary growth of the Mexican Ridges foldbelt, which apparently trapped terrigenous sediment and prevented turbidity currents from reaching the deep sea.

BURCKLE, LLOYD H., and NEIL D. OPDYKE, Lamont-Doherty Geol. Observatory, Palisades, N.Y.

Pan-Pacific Neogene Diatom Correlations

On the basis of piston cores and DSDP sites, a Neogene diatom correlation for Pacific Ocean sediments was determined. Where possible, these correlations rely on the paleomagnetic-reversal record in piston cores. In the central Pacific, diatom datum levels are directly tied to this reversal record. Furthermore, they are extended eastward to DSDP Sites 77 and 158. Although datum levels are generally isochronous in the low-latitude Pacific, some forms have substantially longer ranges in upwelling regions of the easternmost equatorial Pacific. In addition, some high-latitude elements are also present in this region. A mixed assemblage containing both high- and low-latitude diatoms is present at DSDP sites and in piston cores in the northwest and southwest Pacific. This fact permits correlation across several zonal schemes and verifies correlations to such high-latitude sites as 178, 183, and 192 in the North Pacific and 278 in the South Pacific. A series of time-slice maps for the Pacific Neogene depicts changes in the temporal and spatial distribution of diatoms. Diatom-distribution maps reflect accurately the productivity of the overlying water with diatom-bearing sediment present in the eastern equatorial regions and the higher latitude North Pacific. In the South Pacific, diatoms are present south of the Subtropical Convergence. Elsewhere, in the Central Water Masses, the surface sediments are free of diatoms. This pattern holds true generally for much of the Neogene, but major changes in diatom species distribution are indicated in the early Miocene and near the middle/late Miocene boundary.

BURKART, BURKE, Univ. Texas-Arlington, Arlington, Tex.

Pre-Strike-Slip Positions of Polochic Fault Blocks as Determined from Geomorphic Evidence in Guatemala

Previously reported left-lateral slip of 132 ± 5 km across the Polochic fault of Guatemala and Chiapas, Mexico, was based on a match of structures and stratigraphic units. This amount of offset can also be established by a match of present topographies in the region

of the Cuchumatanes Mountains of northwestern Guatemala and the once adjacent region in central Guatemala. By moving the blocks back to their pre-slip positions and examining the union of modern topographies, several interesting correlations are noted. In the pre-slip position the present drainage divide of the Cuchumatanes Mountains on the northern block closely matches the present drainage divide between the Polochic and Chixoy Rivers on the southern block. Ridge and drainage trends extend across the fault so that, in the reconstructed position, a missing part is restored to the annular drainage pattern of the Cuchumatanes domal uplift. The Chixoy and Polochic Rivers on the southern block probably flowed into the present Selegua and Cotzal Rivers respectively. Before slip, the region south of the fault drained northward around the highest elevations of the Cuchumatanes Mountains, but after slip the drainage of rivers on the southern block was shunted into the new, fault-controlled Cuilco and eastern Polochic River channels, vastly reducing the discharge across, and erosion of, the northern block. Significantly lower terrane on the southern block is probably due in large part to this erosional factor.

BURKE, RANDOLPH B., Smithsonian Inst., Washington, D.C.

Bank-Barrier Reef Morphogenesis, St. Croix, United States Virgin Islands

Bank barriers are the major structural reef type in the eastern Caribbean, yet their internal structure and formation process are poorly understood. Twenty-six core holes drilled along the northern St. Croix barrier and on a transect combined with a ship-channel section through the southern bank barrier have shown that this is a bar-type structure which consists of a series of patch reefs within a carbonate sand and rubble matrix capped by reef-crest and fore-reef facies. Locally, reef linearity and location are controlled by a 4-m limestone step of Pleistocene age.

These bank reefs apparently were initiated by shelf-current deposition of carbonate traction load in a bar configuration. Scattered corals—*Porites porites*, *Diploria* spp., *Acropora cervicornis*, and dominantly *Montastrea annularis*—acted to stabilize the bank. Framework-dominated patch reefs developed at scattered locations and depths as the bank grew upward. As the reefs built into shallow water, carbonate production increased, eventually coalescing them into a continuous reef crest.

Core return shows that the structure is characterized by little cementation. Cement occurs in localized lenses within the structure just as it does along the present reef surface. Framework construction is concentrated in patch-reef, reef-crest, and fore-reef facies. This type of construction produces highly permeable structures with extremely high local porosity. Extensively cemented caps occur only in areas having low accumulation rates and in areas of high energy.

Bank-barrier reefs are probably also important in the Indo-Pacific in exposed non-atoll situations, but their occurrence is generally not recognized. Descriptions of Mesozoic and Cenozoic reefs suggest that the bank reef is an extremely important but little understood type in the fossil record also.

CAFLISCH, L., Gulf Science and Technology, Houston, Tex.

Oil Exploration in Pre-Salt and Post-Salt Sequence of West Africa—Results in Cabinda

The Aptian evaporites separate two different oil habitats in the offshore of Cabinda: the pre-salt and the post-salt sedimentary sequence, with three-quarters and one-quarter of the oil production, respectively. In the pre-salt sequence, three oil fields are being produced: Malongo North, Malongo West, and the Block 121 trap. In the post-salt, three shallow accumulations are present, in the Malongo North area.

The pre-salt sequence is extremely variable in thickness, and its sedimentary environment is strictly connected with the initial rifting between Africa and South America. In the earliest Cretaceous, a basement peneplane was flooded by a few fluvial streams along which the thick Lucula sands were accumulated. Production from Malongo North and part of Malongo West fields comes from two of these independent sandstone bodies.

Subsequent subsidence of the basin resulted in the deposition of the Bucumazi black shales in a lacustrine environment. These shales may grade laterally into the Toca carbonate rocks, mainly as a function of the water depth of the lake, every shallowing being accompanied by an increase in carbonate deposition. The Toca carbonate rocks around the Malongo West basement high are the second important reservoir of the field. They are largely dolomitized rocks derived from algal biostromes or pelecypod coquinas.

After the Aptian salt deposition and the definitive opening of the Atlantic Ocean, the basin was tilted to the west, its original trough shape being modified into a monocline. During the Albian, the eastern part of Cabinda offshore was covered by continental red sandstones of the Vermelha Formation. Seaward of this, a definite shallow marine environment existed where the Pinda carbonate beds were deposited, frequently substituted by sandstones and siltstones. These are the reservoir rock of GCO and Mibale fields in Zaire. After a brief marine transgression during the Cenomanian, the heterogeneous Senonian sediments of the Iabe Formation form the reservoir rock of the shallow Malongo fields.

It seems that the proximity of a growth fault may enhance the quality of the post-salt reservoirs: silt and clay accumulated along the downthrown sides of the faults, while salt flowage created localized highs where clean carbonate rocks were deposited in a high-energy environment.

CALZIA, J. P., and J. D. SIMS, U.S. Geol. Survey, Menlo Park, Calif.

Deformational Structures in Pliocene Bouse Formation Near Vidal, Southeastern California

Ball-and-pillow, flamelike, and load structures accompanied by convolute laminations in the Pliocene Bouse Formation are thought to be caused by slumping of water-saturated sediments. New structural data suggest that these structures formed by foundering and by lateral movement of unconsolidated sand into hydro-