

nism, (2) deep-sea environments, and (3) frequency and dynamics of depositional events.

SHARIEF, FAROOQ A., Riyadh Univ., Riyadh, Saudi Arabia, and J. L. WILSON, Rice Univ., Houston, Tex.

Regional Aspect of Sediment Distribution Across Middle East During Middle Triassic Time

Middle Triassic rocks across the Middle East provide information on regional paleogeography based on the distribution of different depositional facies that are clearly related to the structural history of the region. Several structural provinces have been recognized: the Arabian-Nubian massif, the Middle East platform, and the Tethyan trough.

Except for the central part of interior Syria and central and northeastern Iran, the Middle Triassic rocks of the Middle East represent shelf sediments consisting of three major depositional systems. A belt of mixed clastic-carbonate rocks with evaporite deposits of nonmarine, supratidal, tidal-flat, and lagoonal origin is present in the central part of Saudi Arabia. This belt grades into carbonate-evaporite facies of the Arabian Peninsula, southwestern Iran, northern Iraq, and the area of the Syrian-Turkish border. The carbonate-evaporite facies grades into strata which are predominantly open-marine carbonate rock of the Tethyan sea shelf facies over most of Iran, northern Iraq, most of Syria, probably Lebanon, Jordan, Palestine, and the Sinai Peninsula. Several positive and negative areas are within this platform: the Jawf-Rutbah-Mardin high, the Central Arabian embayment, and the Southeastern Saudi Arabia uplift.

The basinal marine sediments of the Tethyan trough facies include Alpine-Mediterranean deposits in central Syria and a terrigenous flyschlike sequence in central and northeastern Iran. The northeastern to northwestern part of central Saudi Arabia holds promise for new petroleum exploration because of a possible great thickness of Permian-Triassic rocks of the carbonate-shelf facies capped by Middle-Late Triassic evaporites.

SHAUB, F. JEANNE, RICHARD T. BUFFLER, and J. LAMAR WORZEL, Univ. Texas Marine Sci. Inst., Galveston, Tex.

Post-Middle Cretaceous Seismic Stratigraphy and Geologic History of Deep Gulf of Mexico Basin

A seismic stratigraphic analysis of 5,000 n. mi of regional multifold reflection data allows an interpretation of the post-middle Cretaceous geologic history of the deep Gulf of Mexico basin. Five horizons that are major unconformities along the southern margin of the basin were mapped throughout the deep Gulf and used to define five depositional sequences (seismic units). Ages for the two youngest horizons were obtained by direct correlation with DSDP holes—Pliocene-Pleistocene and late Miocene. The other horizons are tentatively correlated with proposed major global unconformities and eustatic sea-level changes—middle Oligocene, early Tertiary, and middle Cenomanian. The middle Cenomanian horizon is the most prominent subbottom re-

flector in the deep Gulf. It represents a striking unconformity on the seismic data along the base of the Campeche and Florida Scarps against which the younger sequences thin, onlap, and pinch out. Isopach maps of the units indicate a combined maximum thickness of 7 to 8 km for the post-middle Cretaceous sediments in the central basin. Variations in individual unit thicknesses and seismic facies reflect regional changes in depositional patterns through time: during the Late Cretaceous through middle Tertiary the main sediment source was in the west, and deposition consisted mainly of homogeneous, probably fine-grained sediments changing upward to alternating sand and mud; in the middle Tertiary there was a major influx of sand and mud from both the north and west; following a late Miocene-Pliocene starved-basin interval, fine-grained turbidites from a northern source built the thick Pleistocene-Holocene Mississippi fan.

SHEFFIELD, F. C., Tenneco Oil Co., Houston, Tex.

Future Exploration Opportunities in Gulf of Mexico

Those directly responsible for evaluating recent lease sales in the Gulf of Mexico are painfully aware that the quality of prospects and amount of potential reserves are significantly reduced from what they were several years ago. We believe, however, that this may not be an irreversible situation, in that significant reserves may lie in the undrilled structures of the continental shelf, slope, and abyssal plain of the Gulf of Mexico.

Portions of the South Florida basin and deep waters of offshore Louisiana and Texas are specific areas in the Gulf of Mexico that we feel warrant further investigations by industry.

Subsurface facies and structure maps constructed from available well control, regional CDP seismic, gravity, and magnetic coverages indicate favorable conditions exist in these frontier areas for the entrapment and generation of hydrocarbons. The testing and development of these deep-water and/or deep-drilling objectives will require advanced drilling and production technology.

The most economic means by which to initiate the evaluation of these frontier areas is through stratigraphic-test programs (i.e., COST-type wells). Through the use of stratigraphic tests, better "direct" scientific knowledge can be obtained that would supply the needed exploration incentives and business inducements for industry to invest capital dollars in what may now appear to be prohibitive, high-risk areas. If successful, it would also give industry the lead time required to develop and refine technologies needed to operate in these less than favorable environments.

SHERER, F. C., Sarawak Shell Berhad, Lutong, Sarawak, Malaysia

Exploration in East Malaysia over Past Decade

In the past decade two large oil fields and several large gas fields have been discovered on the continental shelf of East Malaysia together with some smaller oil and gas fields. Geologically, the shelf contains most of the postgeosynclinal younger Tertiary sediments of the