those that fall within the province of plate tectonic theory. Relations between several of these parameters are obvious, e.g., bathymetry and physiographic provinces, and paleomagnetic anomalies and seafloor age. New generations of model reliabilities will occur as further interrelations are found.

The current suite of theories describing overall geologic phenomena, e.g., seafloor spreading and marine sedimentation, are adequate in first order, basinwide terms. Second and third order accuracies are not possible without interpretation on a regional scale.

LUDWIG, WILLIAM J., and PHILLIP D. RABI-NOWITZ, Lamont-Doherty Geol. Observatory of Columbia Univ., Palisades, NY

Seismic Stratigraphy and Structure of Falkland Plateau

Multichannel and single-channel seismic reflection profiles and sonobuoy reflection and refraction measurements indicate that the Falkland Plateau is not a simple extension of South America, but largely owes its morphology to sediments deposited in a continental slope-ocean basin floor environment.

The western part of the plateau is a segment of oceanic crust over which has been deposited 4 to 6 km of sediment in a basin bounded by the Falkland Islands platform on the west, a narrow ridge associated with the Falkland Escarpment on the north, M. Ewing Bank on the east, and the North Scotia Ridge on the south. M. Ewing Bank, a subsided continental block sampled by D/V Glomar Challenger Leg 36, forms the eastern part of the plateau.

The sediments in the basin have been deposited in an oblique progradational-type of configuration. Widespread sheets of sediment dip southward from the Falkland ridge and are terminated updip by erosional truncation. They lap out against the Falkland Islands platform and M. Ewing Bank. The lower boundary of the depositional sequence has been disrupted through movement of the North Scotia Ridge toward the plateau, resulting in subduction of the lower sequence of sediment beneath the ridge and deformation and uplift of the upper sequence to outbuild the northern flank of the ridge.

Overall reflection geometry of sediments filling the basin suggests that they were transported from the north. This implies that they are largely continental slope deposits of pre-drift (>130 m.y. ago) age. Strong bottom currents evidently have caused erosion of significant amounts of the post-drift sediments. The drilling results of D/V *Glomar Challenger* Leg 71 will be discussed in interpretation of the depositional environment of the Falkland Plateau.

- LULMAN, P.S., and G. D. LOBB, Syncrude Canada Limited, Calgary, Alta., and W. E. FLEWITT, Esso Resources Canada Limited, Calgary, Alta.
- Geology of Syncrude Canada Limited Mine Site, Athabasca Oil Sand Area

The Athabasca oil sand deposit covers 4.4×10^6 ha., of which 0.2×10^6 ha. are amenable to surface mining.

In-place reserves of crude bitumen are estimated to be 114.5×10^9 cu m (720 $\times 10^9$ bbl), of which 11.8×10^9 cu m (74 $\times 10^9$ bbl) are within the surface minable interval of less than 46 m. The Syncrude mine site covers 2,850 ha., has in-place reserves of 0.24×10^9 cu m (1.5×10^9 bbl), and commenced production in 1978 with a plant design capacity of 20,500 cu m (129,000 bbl) of synthetic crude per day. The geologic complexities of the oil-bearing McMurray Formation and the overburden zone have had a major impact on engineering considerations at the Syncrude mine.

The Cretaceous McMurray Formation was deposited along a transgressive shoreline between two regional highlands and is interpreted to be mainly estuarine. Paleotopographic lows in the underlying Devonian limestone are filled with salt marsh clays and fluvial water sands. The overlying oil-bearing part of the McMurray Formation is subdivided into a basal fluvial sand, a middle thick estuarine unit with interbedded tidal flat clays, and an upper low-marine unit. Dipping beds contributing to possible highwall instability are associated with estuarine and marine channels.

The overburden zone ranges up to 30 m thick and is composed of marine mudstones and indurated siltstones of the Cretaceous Clearwater Formation and overlying Pleistocene tills, lacustrine clays, and glaciofluvial granular materials, all of which impact on the mine plan.

Detailed documentation of the depositional facies is a prerequisite in geotechnical consideration and mine planning of an oil sand mining operation.

- LYNN, HELOISE BLOXSOM, GEORGE A. THOMPSON, LISA R. KANTER, et al, Stanford Univ., Stanford, CA
- Depth Migration and Interpretation of Cocorp Wind River, Wyoming, Seismic Reflection Data

To better understand the effects of Laramide deformation, deep seismic reflection data in the Wind River (WR) Range area of Wyoming have been migrated using a $45^{\circ} \omega$ -Finite-difference depth migration. The algorithm allows velocity to vary laterally and with depth, contains the thin lens (or shifting) term, and correctly migrates energy in laterally varying media within the limits of a 20 algorithm and 2D dataset. Each migrated section shown is the best from a series of migrations.

Major structural features displayed in the migrated data are the Pacific Creek (PC) anticline and the WR thrust. The PC anticline is underlain by a thrust fault similar in geometry to the WR thrust. The base of the Green River basin sediments has a seismically observed vertical offset of 0.6 km. The intra-basement PC thrust reflections are as conspicuous as the WR thrust reflections, yet the movement along the PC thrust was 0.02 of that of the WR fault. The reflectivity of the PC fault is attributed to the change in seismic impedance of the fault-zone constituents. The thinning of sediments over the anticline in Late Cretaceous time (possibly Lewis, certainly earliest Lance time) indicates that the anticline is part of the Laramide deformation. The anticline continued to grow by faulting and buckling the lower 2 km