

PITTMAN, EDWARD D., Amoco Production Co., Tulsa, OK

#### Effect of Cementation on Physical Properties of Sandstones

Cementation reduces porosity and permeability, and modifies pore geometry. Cementation also increases bulk density, compressive strength, tensile strength, modulus of elasticity, and sonic conductivity. These properties vary with the amount and type of cement as well as the location of the cement, at least for small amounts of cement. Crystal growth at pore apertures has a pronounced influence on permeability, rock strength, and sonic conductivity. The amount, type, and nature of the cement affect wireline log response, upon which many decisions are based.

Major mineral cements in sandstones are quartz, carbonates (calcite, dolomite-ankerite, and siderite), clay minerals, anhydrite-gypsum, and zeolites. Growth habits, crystal shape, crystal size, and location of crystals within the pore system influence pore geometry. Tabular micropores commonly occur among the cement crystals. The nature of the cement affects tortuosity and rugosity.

PLAFKER, GEORGE, and GARY WINKLER, U.S. Geol. Survey, Menlo Park, CA, and GEORGE CLAYPOOL, U.S. Geol. Survey, Denver, CO

#### Geology of Continental Slope Adjacent to OCS Lease Sale 55, near Yakutat, Eastern Gulf of Alaska

Forty-two samples of probable outcrop dredged along the 250-km-long northwest-trending continental slope between long. 138°00'W and 142°30'W from water depths of 3,150 to 200 m provide control on the geology of the seaward margin of the Tertiary basin underlying the outer continental shelf (OCS) Lease Sale 55 on the Yakutat segment of the shelf.

Six rock units apparently in approximately normal stratigraphic position are tentatively recognized. (1) Hard graywacke, argillite, and possible intrusive rocks of inferred Late Cretaceous age crop out from Cross Sound to 139°W and probably underlie much of the Fairweather Ground. (2) 800+ m of shallow-marine basaltic flow and pyroclastic rocks, at least in part of early Eocene age, makes up much of the rugged lower slope from 139°W to about 141°30'W, and rocks of this type are sporadically present west of Yakutat Sea Valley. (3) 500+ m of shallow-marine fine to coarse clastics including organic-rich shale of early and middle(?) Eocene age overlies and intertongues with the volcanic unit from 139°30'W to 142°30'W. (4) 1,500+ m of upper to middle bathyal organic-rich shale, tuffaceous shale, and sandstone of late Eocene and early(?) Oligocene age, apparently overlies the older units between about 140°W to 141°30'W. (5) 450+ m of siltstone with sandstone of probable late Oligocene age underlies the upper slope between 139°W and 140°W. (6) Neogene marine glacial deposits and mud occur on the OCS as slope-basin fills and slope-rise prisms, and as a mantle over the slope at the mouths of the Alsek Canyon and Yakutat Sea Valley.

The lower Tertiary sequence sampled on the conti-

mental slope differs significantly from coeval rocks onshore or penetrated by wells on the Yakataga segment of the OCS to the west. The Eocene and Oligocene units sampled include abundant thermally mature source rocks. Most dredged sandstones are poor reservoirs, but some upper Eocene to lower Oligocene sandstones have fair to good porosity and permeability. Seismic reflection data indicate that the Eocene and Oligocene sequence dips beneath Lease Sale 55 where it could be a favorable exploratory target.

PLUMLEY, WILLIAM J., Chevron Oil Field Research Co., La Habra, CA

#### Effective Stress and Abnormally High Fluid Pressure

Published hypotheses of the causes of abnormally high fluid pressure have not explored explicitly the relation between the present effective stress ( $s_p$ ) and the maximum effective stress ( $s_m$ ) to which a sediment has been exposed during its history. In basins where sediments are at their maximum burial depth, the present effective stress can be used to evaluate quantitatively different hypotheses for abnormal fluid pressure development. The underlying assumption of this approach to hypothesis testing is that shale porosity is principally a function of maximum effective stress.

"Nonequilibrium compaction" as a cause of abnormally high fluid pressure requires that  $s_p = s_m$  and predicts porosities that are higher than would be obtained if another cause, such as "aquathermal pressuring" or "clay transformation" (which require that  $s_p < s_m$ ), were the predominant mechanism producing the same abnormally high fluid pressure. Observed porosities in Gulf Coast high-pressure shale formations commonly are too low to be solely the result of nonequilibrium compaction. In two field examples, shale porosities predicted with the nonequilibrium compaction model are about 20 porosity units higher than porosities determined from  $\gamma - \gamma$  density logs and cores. In one example, a combined mechanism of nonequilibrium compaction and clay transformation is one possible, and internally consistent, interpretation of the fluid pressure and porosity data. In the other example, the fluid pressure and porosity data are consistent with clay transformation being the mechanism for the generation of the abnormally high fluid pressure, but are not consistent with the nonequilibrium compaction model.

POGORSKI, LOUIS A., and G. STEWART QUIRT, Chemical Projects Ltd., Toronto, Ontario

#### Helium Surveying for Deeply Buried Uranium Deposits

Helium emanometry is rapidly becoming a widely employed exploration tool in searching for deeply buried uranium deposits. It has been used in many uraniumiferous and potentially uraniumiferous regions including the Colorado Plateau, Texas Gulf Coast, Wyoming, Appalachian orogenic belt, Athabasca basin, and Canadian Northwest Territories.

Helium escaping from radioactive mineral deposits can be detected and serves as a guide to the location of uranium ore. This gas is an almost ideal geochemical indicator of uranium because it is inert, stable, slightly

soluble in ground fluids, practically non-adsorbable, highly mobile, and a direct product of radioactive decay.

The technique can be applied by collecting soil, soil gas, water or bottom sediment samples in reconnaissance, and semi-detailed and detailed arrays under a wide range of environmental conditions. Helium analyses are made by gas-source mass spectrometry. The resultant data are interpreted and presented with the aid of computers. In interpreting the helium data, it is necessary to consider the effect of some parameters which must be determined for each sample.

Helium anomalies have been found in near-surface soil and soil gas over known sandstone-type deposits in New Mexico, Texas, and Wyoming; hydrothermal(?) ore in Washington; unconformity-type mineralization in the Athabasca basin; and pegmatitic ore zones in Ontario. Anomalies have also been detected in lake bottom water and sediment overlying these types of deposits and in the groundwater recovered from wells and boreholes located close to them. The results from resurveys over several of these deposits indicate that even though the magnitude of the helium anomalies may vary from season to season, the anomalies themselves persist and hence define the location of the mineralization. This technique therefore seems to offer great promise as an economical indicator of deeply buried uranium deposits in a wide range of geologic environments.

POOLE, FORREST G., and GEORGE A. DESBOROUGH, U.S. Geol. Survey, Denver, CO

Oil and Metals in Ordovician and Devonian Kerogenous Marine Strata of Central Nevada

Kerogen-rich mudstone, siltstone, dolomite, and chert units as much as 50 m thick in the Vinini (Ordovician) and Woodruff (Devonian) formations contain potential resources of syncrude oil, V, Zn, Mo, Se, Ag, and Cr. Most kerogenous rocks originally consisted of organic-rich siliceous muds, slimes, and oozes. Organic matter is mostly amorphous, flaky, and stringy sapropel composed of planktonic organisms. The strata are within strongly deformed eugeosynclinal Paleozoic marine rocks of the Roberts Mountains allochthon.

Many fresh black rocks are low-grade oil shales which, upon pyrolysis, yield <40 l of oil per metric ton of rock; some thin layers yield as much as 125 l per metric ton. In these rocks, solid bitumen and liquid oil commonly fill voids and microfractures. Such early-phase hydrocarbons probably were released during diagenesis and formed without any major thermal degradation of the kerogen. Geochemical data suggest that the organic matter is thermochemically immature to mature and has not been subjected to temperatures above 60°C since deposition. Hydrocarbon contents (<100 to 5,400 ppm) and organic carbon contents (<1 to 25 weight %) vary widely.

V, Mo, Se, Ag, and Cr in fresh black rocks occur chiefly in organic matter; Zn occurs as sphalerite and Ni in iron sulfides. Concentrations are as much as 5,000 ppm V, 18,000 ppm Zn, 1,000 ppm Mo, 100 ppm Se, 20 ppm Ag, 150 ppm Ni, and 600 ppm Cr in unoxidized

rocks. Enrichment of V and Se and depletion of Zn, Mo, Ni, and organic matter occur in oxidized rocks.

POWERS, RICHARD B., U.S. Geol. Survey, Denver, CO

Oil and Gas Potential of Wyoming-Utah-Idaho Overthrust Belt—Relation to Canadian Foothills Province Analog

The Cordilleran orogenic belt is generally considered to be a single tectonic element extending from northern Alaska to Central America. Two segments of the North American part of this element are oil and gas producers—the Canadian foothills thrust belt and the Wyoming-Utah-Idaho Overthrust belt.

Turner Valley was the first field discovered in the Canadian foothills. In the 55 years since that discovery, 32 fields have been found containing 9.3 Tcf of initially recoverable gas, 143 million bbl of natural gas liquids and 132 million bbl of oil. The first significant field discovery in the U.S. Overthrust belt was made in 1975 at Pineview. By the end of 1979 eleven new fields had been found, containing an estimated 500 million bbl of recoverable oil and 5.5 Tcf of recoverable gas, plus natural gas liquids.

The Canadian province is considered to be an appropriate geologic analog to the U.S. Overthrust belt, based on a number of characteristics common to both provinces. These include general structural configuration, trap types, reservoirs, stratigraphy, timing of migration of hydrocarbons, depth of burial, and age of tectonic movement. However, significant differences include age of major source rocks and paleothermal histories.

The future potential of the immaturely explored Wyoming-Utah-Idaho Overthrust belt is assessed by using a volumetric method, wherein hydrocarbon yields in barrels of oil and cubic feet of gas per cubic mi of sediments are established for the more densely explored Canadian foothills. These yields are applied to the less densely explored U.S. Overthrust belt. The usefulness of this method lies in the correct interpretation and analysis of similar, as well as dissimilar, geologic characteristics of the analog province. In addition, the assessment of the U.S. Overthrust belt is further refined by using the southerly productive part of this province as an "internal" analog which is applied to the remaining area of the province.

PRATT, BRIAN R., Petro-Canada Exploration Inc., Calgary, Alta.

Algal-Metazoan Bioherms of Lower Ordovician Age—St. George Group, Western Newfoundland

Bioherms are common in the St. George Group, a sequence of shallow-water carbonate rocks deposited on the western continental shelf of Iapetus Ocean. The cores of these bioherms are composed of thrombolites (unlaminated, branching, columnar stromatolites), "calcareous algae," and corals. On the basis of framework-building components, three types are distinguished: (1) thrombolite mounds, (2) *Lichenaria*-thrombolite mounds, (3) thrombolite-*Lichenaria-Renalcis* reefs. Associated with these structures is a diverse fauna of bur-