

sabkha and dune facies were deposited over marine carbonate rocks. Desert conditions prevailed at the eastern edge of the sea as it transgressed eastward and regressed westward across northern Arizona. Eolian dunes with south-dipping cross-beds were formed by trade winds blowing southward toward the paleo-equator south of Arizona during the Permian. Extensive coastal and continental sabkhas formed between the restricted mud flats and the dune fields. Westward, the restricted marine deposits of dolomite characterized by bivalves and gastropods gave way to shallow open-marine deposits of brachiopods, byzozoans, corals, and crinoids. The westward regression ended with a rapid transgression that deposited the Kaibab Limestone across the sabkhas and dune fields of the Toroweap and Coconino formations.

READ, J. F., Virginia Polytechnic Inst. and State Univ., Blacksburg, VA, and J. R. MARKELLO, Chevron U.S.A. Inc., Lafayette, LA

Carbonate Depocenters and Facies Distribution on Passive Cambrian Shelf and Middle Ordovician Foreland Basin, Appalachian Orogen, Virginia

Depocenters in the southern and central Appalachians in Virginia appear to have been a major influence on thickness and carbonate facies distribution on the passive Cambrian-Ordovician shelf and in the Middle Ordovician foreland basin.

The Upper Cambrian Nolichucky Formation illustrates sedimentation patterns associated with a shelf embayment (located above the southern Appalachian depocenter) on the passive Cambrian shelf. The Nolichucky Formation is an onlap-offlap, shallow ramp to deep shelf sequence that consists of peritidal stromatolitic carbonate rocks, ooid grainstones, deeper ramp ribbon carbonate rocks, and embayment plain shale/siltstone/flat-pebble conglomerate facies. Facies bands are normal to the regional shelf edge and parallel the outline of the shelf embayment.

The Middle Ordovician sequence illustrates the influence of depocenters on foreland basin evolution during a time of profound tectonism, when the shelf edge was uplifted, deformed, and subjected to erosion. The Middle Ordovician ramp-to-basin sequence is an onlap-offlap package, that consists of peritidal fenestral lime mudstones, shallow subtidal cherty wackestones, ramp and downslope skeletal buildups, deeper ramp shaly skeletal wackestones, and basal black limestones and shales. Widespread ramp and basin deposition commenced in southwest Virginia during south to north transgression from the southern depocenter. Widespread downwarping extended the basin into northern Virginia. Rapid clastic influx coupled with progradation of the carbonate ramp caused the southern basin to fill.

Recognition of depocenters associated with carbonate sequences in orogenic belts is important if controls on direction of transgression, facies distributions, and thicknesses of units are to be better understood. Furthermore, the carbonate depocenters appear to localize development of subsequent clastic wedges.

REAUGH, A. B., and G. S. BAYLISS, GeoChem Laboratories, Inc., Houston, TX

Fluorescence of Acritarchs in Study of Marine Kerogen

Acritarchs, the organic-walled palynomorphs of algal affinities, exhibit strong fluorescence colors depending upon species type, stage of thermal maturity, and degree of oxidation.

This fluorescence readily enables identification of acritarchs in gross kerogen concentrates and provides a means of distinguishing acritarch fragments in poorly preserved kerogen assemblages.

Although acritarchs show similar fluorescence colors in an autochthonous assemblage, there may be minor differences as a result of different types and wall thicknesses. Similarly, the varying degrees of oxidation of individual palynomorphs can give differing colors.

Thick-walled marine algae also fluoresce but can be distinguished from acritarchs by the differences in intensity of the fluorescence colors.

REAUGH, A. B., and G. S. BAYLISS, Geochem Laboratories, Inc., Houston, TX

Depositional Environments and Kerogen Types

No abstract available.

REDWINE, LOWELL, Consulting Geologist, Costa Mesa, CA

Hypothesis Involving Dilation and Natural Hydraulic Fracturing to Explain Petroleum Reservoirs in Monterey Shale, Santa Maria Area, California

Fractured reservoirs in the siliceous Monterey Shale of the Santa Maria area represent anomalous lithology and anomalous type of fracturing. Some, perhaps all, reservoirs are not fractured chert but chert embrittled by dolomitization. Abundant reservoir extension fractures are disordered and open, commonly containing epigenetic dolomite breccias. These fractures are partly dolomite-cemented but contain common open voids, many 15 cm across, some larger. Breccias locally have an exploded appearance and contain matched fragments separated by veins, which apparently were injected as a slurry of water, oil, and fragments of dolomite and dolomitic Monterey Shale.

The highly organic Monterey served also as source rock and probably originated as richly diatomaceous slope sediment beneath an oxygen-minimum zone at a depositional site much larger than the Santa Maria area, and not confined to a specific silled basin. Local dolomitization may have been due, at least in part, to rising solutions and injected slurries.

These reservoirs are explained by an hypothesis involving repeated episodes of rock dilation followed by natural hydraulic fracturing, all produced by continued episodic tectonic compression of the region (principal, maximum, effective stress oriented northeastward). High fluid pressures enlarged underpressured dilation microfractures into macrofractures and produced breccias by hydraulic fracturing. Viscous oil derived from the Monterey Shale was forced into voids as part of overpressured slurries whose breccia fragments were

propping agents. Dolomite precipitated from slurries partly healed the fractures.

REID, FRED S., and CLAUDE E. BERGHORN, Peppard-Souders and Associates, Denver, CO

Lower and Middle Paleozoic Potential of Paradox Basin

Successful oil exploration in the Paradox basin began in 1907 with the discovery of Pennsylvanian oil at Mexican Hat field in San Juan County, Utah. Accelerated exploration began with the discovery of oil in the Pennsylvanian Paradox Formation at Aneth field in 1956 and lasted into the 1960s. Cumulative production at greater Aneth field exceeds 271 million bbl of oil. A second major oil producing zone, the Mississippian Leadville, was established at Lisbon field in 1960. By 1978, its 25 wells had produced 40 million bbl of oil and 310 Bcf of gas. Recently, exploration in the area has been minimal with the exception of the current exploration for carbon dioxide for secondary recovery projects.

The Paradox Formation, the major producing unit in the basin, can be subdivided into both genetic units and facies subzones using sample and mechanical log information. Recognition and mapping of the major facies—hypersaline, penesaline, and marine shelf—for each genetic unit can delineate fairways most favorable for the development of algal mounds in which the best production occurs.

Perhaps the most significant remaining potential is in the Mississippian Leadville formation. Exploration for structural traps within this carbonate unit is difficult owing to the thick overlying Pennsylvanian salt section present over much of the basin. A pilot gravity modeling study in the Lisbon field area has shown that the salt effect can be removed and the underlying structure mapped. Regional gravity modeling of the remaining salt basin may define similar Lisbon-type structures.

REIMER, G. M., A. A. ROBERTS, and M. E. HINKLE, U.S. Geol. Survey, Denver, CO

Recent Advances in Helium Analysis as Exploration Tool for Energy "Deposits"

Recent research by the U.S. Geological Survey demonstrates that helium-gas analysis of waters and soils holds great promise as a cost-effective exploration technique for uranium, oil and gas, and geothermal energy sources. The technologic advances include assembling a helium analyzer, almost entirely from commercially available equipment, and packaging the equipment into a mobile laboratory capable of performing as many as 100 analyses a day at a field location. Helium is an attractive indicator element for many exploration programs because of its unique properties: it is highly diffusive, chemically inert, radioactively stable, and not produced or affected by biologic activity. Many associations of helium with uranium have been observed, in which helium is produced by natural radioactive disintegration; with oil and gas, where helium is trapped by structural and stratigraphic features; and with hot-water geothermal systems, in which the cooling and reduced pressure of rising water causes dissolved helium to be released. The following are examples of

distinctive helium anomalies found associated with energy "deposits": for uranium, the Ambrosia Lake district, New Mexico; for oil and gas, the Cement oil field, Oklahoma, and the Cliffside gas field, Texas; and for geothermal, the East Mesa known geothermal resource area in the Imperial Valley, California. With respect to an ambient air background of 5.24 ppm helium, the highest observed concentrations of excess helium in soil and soil gas were typically 0.5 ppm for uranium, 10 ppm for oil and gas, and 100 ppm for geothermal; water samples usually had several hundred parts per million helium for all types of energy deposits. Helium analysis can be used as a rapid and inexpensive reconnaissance tool and as complementary support for other geophysical and geochemical prospecting techniques.

REINSON, GERALD E., and GUSTAV VILKS, Geol. Survey Canada, Dartmouth, Nova Scotia

Seabed Characteristics and Sand Dispersal on Bedrock-Dominated Inner Shelf of Southern Labrador

The Labrador Shelf is dissected by the Marginal Trough into a narrow inner rocky shelf, which is the submerged extension of the Precambrian Shield landmass, and an outer shelf zone consisting of broad, flat banks mantled by thick deposits of glacial drift. The inner shelf north of Groswater Bay is 35 km wide, with a highly irregular bedrock-dominated bottom topography. Unconsolidated materials consist of sand, and coarse gravel "pavement" deposits. Sand deposits are less than 1 m thick and limited in areal extent to the flat-bottomed, low-lying areas between bedrock highs. Coarse gravel deposits occur as veneer pavements on the flanks of highs. The sands are underlain either by cohesive muds (early Holocene?) which were deposited in former basinal depressions, or by coarse gravels in local areas marginal to bedrock outcrops. The coarse gravels are probably relict lag deposits formed by the reworking of glacial drift, but the sands are thought to be derived from contemporary nearshore and beach sediments situated about 10 km west of the study area.

The thin and patchy sand distribution suggests that transport mechanisms are more than sufficient to disperse the volume of sand that is being supplied to the inner shelf. Preliminary analysis of near-bottom velocity measurements indicates that the seabed is subjected to a strong southeasterly current (Labrador current) which induces a net southeasterly sand flux across the shelf. The predominantly resistant substrate of the shelf would likely be swept clean of sand, if it were not for the irregular bottom configuration which provides local and temporary sinks for sand deposition.

The most important sediment-transport process on the inner shelf is the southeasterly directed Labrador current. Wave-generated currents are of lesser importance (except in shallow nearshore areas) as a sand-dispersal mechanism, and iceberg-scouring is more effective in redistributing sediment in areas seaward of the inner shelf edge.

REITER, MARSHALL, and CHARLES SHEARER, New Mexico Bur. Mines and Min. Resources and New Mexico Inst. Mining and Technology, Socorro, NM