

sized tangential, wedge and trough sets of cross-beds. The Atarque Member is about 11 ft (3.4 m) thick and contains fossil-rich lenses of poorly sorted, dark yellow-brown-weathering sands that are friable to moderately indurated and calcite cemented. Shale galls are present in the fossil-rich lenses and, with wood fragments, abundant turtle bone fragments, some crocodile tooth and scute fragments, amid vertebrae and teeth indicate a nearshore environment with a nearby source of fresh water. Selachian teeth and probable coprolites are richly varied and abundant. The following genera have been recognized: *Hybodus*, *Lonchidion*, *Squalicorax*, *Squatina*, *Brachaelurus*, *Scapanorhynchus*, *Odontaspis*, *Cretoxyrhina*, *Cretolamna*, *Plicatolamna*, *Paranomotodon*, *Ischyryza*, *Ptychotrygon*, *Rhombodus*, as well as several as yet indeterminate genera. The dominant invertebrate genus is the oyster *Crassostrea* although at least two genera of gastropods are present.

BARGHOORN, STEVEN F., Columbia Univ., New York, NY

Geomagnetic Polarity Stratigraphy of Type Tesuque Formation, Santa Fe Group, in Espanola Valley, New Mexico

Sampling of the several type sections of the Tesuque Formation, lower Santa Fe Group, exposed near Espanola, New Mexico, has yielded a composite magnetic-polarity stratigraphy in which the 760-ft (230 m), fine-grained, tuffaceous middle (Skull Ridge) Member is characterized by a long, uninterrupted interval of reversed polarity; approximately 900 ft (275 m) of the coarser grained upper (Pojoaque) member is characterized by rocks predominantly of normal polarity. Laboratory results from AF demagnetization in fields up to 200 oe, and from the acquisition and removal of IRM (often of multiple specimens cut from single samples), indicate that the magnetic signals in these rocks, probably borne by detrital magnetite, are a record of the geomagnetic field at or near the time of deposition. The stratigraphic utility of the results is confirmed by replication in geographically separate, lithostratigraphically equivalent sections in different fault blocks. Within the context of the medial to late Miocene time span long acknowledged for this part of the Santa Fe Group, this composite polarity stratigraphy appears to correlate best with the standard Tertiary polarity time scale as follows: fossiliferous Nambe Member: early Epoch 16; Skull Ridge Member: late Epoch 16 and early Epoch 15; Pojoaque Member: mid-Epoch 15 and younger. These results indicate an age for at least the Pojoaque Member significantly younger (by about 3 to 4 m.y.) than fission-track ages recently reported for ash beds in the type Pojoaque section.

BLACK, BRUCE A., Colorado Plateau Geological Services Inc., Consultant, Farmington, NM

Effect of Oil Prices and "Windfall Profit" Taxes on Life or Death of an Oil Field—a Case History

The dramatic effect of price increases and subsequent taxing of profits is obvious for small oil operators who are attempting to produce marginal oil and gas properties. The Red Mountain oil field and associated leases in the San Juan basin of New Mexico are a prime example of the recent "yo-yo effect" of oil and gas pricing and subsequent taxing.

An analysis of the profitability of these operations, and its subsequent effect on oil field employment and purchases of equipment, are a clear, if small, example of the impact recent

government policies have had throughout the oil industry.

Increases in the price of oil and gas made operations temporarily economic, and inspired additional drilling. This activity actually resulted in discoveries of new oil and gas sands and increased oil and gas reserves. However, the effect of a positive price has been partly or totally cancelled by taxing through the "windfall profit" tax. "Windfall profit" taxes have had a negative effect by shutting down additional exploration, cancelling planned waterflood projects, and generally hindering profitable operations.

This microstudy can be multiplied by hundreds and perhaps thousands of similar examples throughout the United States, with the effect of shutting in millions of barrels of oil, which would otherwise have been discovered and recovered. This is a prime example of illogical and arbitrary taxing procedures and policies.

The ramifications are felt not only by the oil and gas operators but also by suppliers, oil field workers, and their families. The economics of the area, and thus the state and the nation as a whole are adversely affected.

To compound this problem, we pay over \$40 per barrel for imports to make up for the oil we have in these areas but do not recover. This \$40 per barrel would have been paid to Americans, in America, and would have bought American products and paid American labor. This in turn would have increased the American labor force and the American inventory, and would have used American goods to find and increase American reserves.

Instead this money is diverted to foreign lands where it buys foreign goods and services. When, and if, it is reinvested in America it may be used to buy political influence, or buy control of American lands and industry. This microexample shows the economic insanity of such a program.

BROOKINS, DOUGLAS G., Univ. New Mexico, Albuquerque, NM

Geochronologic Studies near WIPP Site, Southeastern New Mexico: Summary and Interpretation

The WIPP site is currently being investigated for pilot storage of defense-generated radioactive waste. A critical facet of the overall study is the question as to whether the rocks have remained closed systems with respect to their bulk chemistry and isotopic composition since formation in the Late Permian as opposed to later episodic or continuous recrystallization. Early attempts at K-Ar dating of sylvites mixed with other salts yielded inconclusive results, primarily because sylvite is not well suited for K-Ar study owing to loss of radiogenic ^{40}Ar . Rb-Sr study of sylvites yields a 214 ± 15 -m.y.B.P. isochron indicating closed system conditions to Rb and Sr since latest Permian or earliest Triassic. Pre-200-m.y.B.P. K-Ar dates have also been determined for pure langbeinites and polyhalites. When mixed with sylvite, age lowering results. The Rb-Sr systematics of aeolian clay minerals known to have interacted with the evaporite brine yield a poorly defined isochron of 390 ± 75 m.y.B.P., but the apparent date indicates that the clay mineral-brine interactions were not so severe as completely to rehomogenize Sr isotopes despite the clay-mineral alteration. A 34 ± 1.5 -m.y.B.P. lamprophyre dike intrudes the evaporite sequence 16 km north of the WIPP site. Contact effects, including recrystallization of polyhalite, are restricted to within 10 m of the dike. Finally, polyhalite inclusions in one rubble chimney yield a pre-200-m.y.B.P. age indicating no major recrystallization effects due to this disturbance of the evaporite sequence. Collectively, the geochronologic studies argue for