

OLSEN, CLAYTON E., Los Alamos Scientific Lab., Los Alamos, NM, and D. G. BROOKINS, Univ. New Mexico, Albuquerque, NM

Distribution of Uranium as Function of Sediment Particle Size

Uranium concentrations as a function of size were determined for 100 samples from the Estancia Valley, New Mexico, 100 samples from northeast of the Grants mineral belt, and 46 samples from the Black Hawk mining district, southwestern New Mexico. For the Estancia and Grants samples, statistical analysis shows uranium concentrations in stream sediments size fractions greater than 0.125 mm to be nearly independent of particle size whereas uranium concentration increases with decreasing particle size less than 0.125 mm. The standard deviations of uranium concentrations from both areas vary only slightly for all size fractions. For the Black Hawk mining district the uranium concentrations are strongly dependent on sediment particle size with uranium increasing rapidly with decreasing particle size. A correlation coefficient of 90% is calculated for increasing uranium content with surface area of sedimentary particles. In contrast, a much lower correlation coefficient for uranium-to-surface area is observed for the Estancia and Grants samples. We propose that the observed statistical differences are related to the fact that the stream sediments of the Black Hawk mining district represent control by mechanical disintegration of igneous and metamorphic rocks with little effect of chemical weathering, whereas the sediments of the Estancia Valley and Grants area are more chemically mature. The data imply that the nature (maturity) of the sediments sampled must be considered when sampling on a large grid, such as one sample/10 km², or else risk false anomalies (positive or negative) or masking any anomalies which may be present.

OTTON, JAMES K., U.S. Geol. Survey, Denver, CO

Model for Uranium, Humate, and Silica Fixation in Lacustrine Carbonaceous Sediments at Anderson Mine, Western Arizona

Sedimentation at the Anderson mine in western Arizona was dominated by subaerial and subaqueous debris-flow and lacustrine-turbidite processes. The typical debris flow at the mine is 30 to 100 cm thick, massive, internally chaotic, composed of various amounts of reworked tuff, mica, quartz, and silicified plant and shelly detritus, and has undergone varying degrees of zeolitization and silicification. These debris flows grade laterally to thinner debris flows with graded bedding and laminated lacustrine turbidites that are rich in carbonaceous organic detritus. The more distal flows and turbidites are the principal uranium host at the mine. Uraniferous silica, amorphous uranium silicate, and coffinite are commonly associated with internally structureless though crudely stratified organic matter (humate?) in the carbonaceous siltstones. Uranium is not associated with fragments of coaly plant debris. Uranium, silica, and humic(?) material were probably leached from proximal subaerial and subaqueous debris-flow facies and transported to the anoxic lake-bottom environment where reducing conditions, low permeability, and porosity prevailed. In the anoxic environment, uranium, silica, and humate all tended to precipitate from open water at an Eh-pH boundary zone either at the sediment-water interface or at some depth in the sediment. The carbonaceous laminated lacustrine turbidites were the favorite host, but individual orebodies cross lithologic boundaries, suggesting that uraniferous silica apparently precipitated wherever reducing conditions existed on the lake bottom.

PARKER, MARK D., and GEORGE R. JIRACEK, Univ. New Mexico, Albuquerque, NM

Evaluation of Geothermal Potential, Albuquerque Area, New Mexico

The first comprehensive evaluation of the geothermal potential of the Albuquerque, New Mexico, area, recently completed, included gravity and magnetic surveys, electrical resistivity measurements, temperature logging of water wells and of shallow (25 m) temperature-gradient boreholes, and geochemical analysis of local waters. Local seismic history and temperature records from municipal water wells were also analyzed. Geothermal anomalies of the Albuquerque area were delineated if temperature gradients in excess of 35°C/km were detected, as this figure is the average temperature gradient for the Rio Grande rift.

Temperature logging of city water wells resulted in the identification of two well fields having anomalously warm water at shallow depth. The Walker well field, on Albuquerque's north side, has 32°C water at a depth of 320 m, representing a gradient of 53°C/km. At the West Mesa well field, on Albuquerque's west side, recent pumping prevented the obtaining of an accurate temperature gradient. However, repeat measurements at selected depths suggest a gradient of approximately 65°C/km.

A third site, about 8 km west of Albuquerque, was found to have shallow temperature gradients in excess of 120°C/km. Near this site, magnetic surveys indicated that several igneous bodies are buried beneath 300 m of alluvial sediments. Because no water wells exist at this remote location, shallow boreholes were drilled for temperature logging. The resulting values of temperature gradient show a distinct high of greater than 120°C/km superposed on a background of 60°C/km.

PAULL, RACHEL K., Univ. Wisconsin-Milwaukee, Milwaukee, WI

Stratigraphic Relations of Type Section of Lower Triassic Dinwoody Formation in Wyoming to Exposures in Idaho Overthrust Belt and Northwest Utah

The Lower Triassic Dinwoody Formation, the oldest Triassic unit in the Cordilleran miogeosyncline, rests unconformably on underlying Permian rocks. The type section of the Dinwoody is on the northeastern flank of the Wind River Mountains in Wyoming. As this locality is near the eastern margin of the Early Triassic depositional basin, the formation is relatively thin (52 m) and largely composed of clastic sediments. To the west, the Dinwoody thickens abruptly and the amount of interbedded limestone increases.

In the Bear Lake-Montpelier region of the southeastern Idaho overthrust belt, the thickness increases to about 380 m. Although significant amounts of interbedded carbonate are present here, intertonguing red beds and evidence of periodic shoaling also characterize this transitional area. In northwest Utah, the Dinwoody thickens to 410 m and includes numerous interbeds of limestone.

Conodont biostratigraphic zonation of the Dinwoody Formation permits detailed correlation and establishes the timing of the Early Triassic transgression eastward over the Wyoming shelf. This short-lived event was followed by westward progradation of terrigenous sediments from the shelf into the depositional basin.

On the basis of color change of conodont elements, which relates to depth and duration of burial, thermal maturity of the Dinwoody Formation increases southwestward. The color