

---

---

# Association Round Table

---

---

**AAPG SOUTHWEST SECTION**  
**Annual Meeting, February 25-27, 1980**  
**El Paso, Texas**

## Abstracts of Papers

AIKEN, CARLOS, and DAVID GARVEY, Univ. Texas, Dallas, MAURICIO DE LA FUENTE, Consejo De Recursos Minerales, Mexico City, et al.

Geophysical Studies of Chihuahua City Region, Mexico

No abstract available.

BARROWS, L. J., Sandia Laboratories, Albuquerque, NM

Geophysical Studies of Evaporites in Northern Delaware Basin, Texas

Geophysics has been an integral part of the Waste Isolation Pilot Plant since its inception. Previous activities have included electrical resistivity surveys, review of petroleum exploration seismic lines, purchase of aeromagnetic and gravity surveys, installation of an earthquake seismometer, a seismic refraction survey, and five seismic reflection surveys. Ongoing or planned activities include installation of a multistation seismometer array, subsurface radar, vertical-gradient aeromagnetics, and high-resolution seismic work.

The seismic reflection and gravity data indicate complex structure in the northern part of the site. The seismic lines show a deterioration in reflector quality, character, and continuity within this disturbed zone. Events within the Castile Formation have a blocky nature with abrupt offsets and angular discontinuities between blocks. The lower members of the Castile Formation are thinner than normal. Both the underlying Delaware Mountain Group and overlying Salado Formation are less deformed than the Castile. Well control is consistent with the seismic interpretation. An interesting feature is an "expandable scale" contraction of the lower Salado in two wells in which individual beds are present but have been reduced 10 to 50% in thickness.

BOCKOVEN, NEIL T., Exxon Minerals Co., Albuquerque, NM

Geology of Sierra del Gallego Area: Comparison with Pena Blanca Uranium District, Mexico

A 1,000-m section of volcanic rocks overlying Lower Cretaceous limestone in the Sierra del Gallego area 200 km south of El Paso can be divided into four lithologic sequences. About 44 to 38 m.y. ago, at least five separate crystal-poor sanidine-bearing outflow units of rhyolitic tuff covered parts of the area. The tuffaceous activity was closely followed by lava flows of fritted feldspar-bearing tholeiitic andesite. These

flows were followed about 37-35 m.y. ago by a sequence of rhyolite flows, flow domes, and intrusions. A 6-m.y. period of quiescence was followed about 29 m.y. ago by massive outpourings of basalt with minor associated rhyolite tuff. Parts of the oldest sequence are probably correlative with stratigraphic units in the Pena Blanca area in New Mexico. The Pozos limestone conglomerate at the deposit has a counterpart in the Sierra del Gallego area. Petrographic similarities, thickness relations, geodes, and K-Ar dating directly link the H member in the lower part of the Liebres Formation in the Sierra del Gallego area with the Nopal Formation at Pena Blanca. The Pozos, Nopal, Escuadra, Pena Blanca, and Mesa Formations are, as a group, lithologically and chronologically similar to the oldest sequence (the Liebres Formation) of the Sierra del Gallego area.

BORNHORST, THEODORE J., WOLFGANG E. ELSTON, and RICHARD S. DELLA VALLE, Univ. New Mexico, Albuquerque, NM, et al

Distribution of Uranium in Middle Tertiary Volcanic Rocks, Mogollon-Datil Volcanic Field, New Mexico

The uranium abundances in middle Tertiary volcanic rocks of the Mogollon-Datil volcanic field, southwestern New Mexico, have been determined as part of a major petrogenetic study. Over 350 samples of middle Tertiary to Quaternary volcanic rocks have been analyzed for their uranium content by delayed neutron activation analysis.

Of the volcanic associations previously proposed for southwestern New Mexico, calc-alkalic andesite,  $\pm 43$  to  $\pm 35$  m.y., has a mean of 2.3 ppm U (range 0.9 to 5.4 ppm); calc-alkalic quartz latite to rhyolite,  $\pm 35$  to  $\pm 29$  m.y., has a mean of 3.9 ppm U (range 1.7 to 6.2 ppm); basaltic andesite and associated rocks,  $\pm 32$  to  $\pm 18$  m.y., has a mean of 2.3 ppm U (range 0.8 to 6.9 ppm); and high-silica rhyolite,  $\pm 32$  to  $\pm 18$  m.y., has a mean of 5.2 ppm U (range 1.6 to 9.4 ppm). Anomalous values in the range of 14 to 35 ppm U were found for a riebeckite-bearing lava from the central San Mateo Mountains, a sample of intrusive andesite from the Alum Mountain area, and a lithophysal rhyolite lava and associated ash-flow tuff from the Sierra Cuchillo. Post-13-m.y. bimodal basalt-rhyolite is sparse within the Mogollon-Datil volcanic field. A few determinations from this study, and published and unpublished data for other localities in New Mexico, indicate U abundance from 0.3 to 1.5 ppm U in post-13-m.y. basalt and about 7 to 8 ppm U in rhyolite.

BURT, DONALD M., and MICHAEL F. SHERIDAN, Arizona State Univ., Tempe, AZ

Model for Formation of Uranium/Lithophile Element Deposits in Fluorine-Enriched Volcanic Rocks

Many uranium and other lithophile element deposits are

within or adjacent to small fluorine-rich rhyolitic dome complexes or tuff sheets. Examples studied include Spor Mountain, Utah (Be-U-Li-F), the Honeycomb Hills, Utah (Be-U-Li-F), Wah Wah Mountains, Utah (U-F), and the Black Range, New Mexico (Sn-Be-F). The formation of these and similar deposits begins with the emplacement of a rhyolitic magma, fractionated in lithophile metals and complexing fluorine, that rises to a shallow crustal level, where its roof zone may become further enriched in volatiles and the ore elements. During initial explosive volcanic activity, aprons of lithic-rich tuffs and surge deposits are erupted around the vents. These early pyroclastic deposits commonly host the mineralization, owing to their initial enrichment in the lithophile elements, their permeability, and the reactivity of their foreign lithic inclusions (particularly carbonates). The breccias are capped by thick topaz rhyolite lavas or welded tuffs that can serve as a source of heat and of additional quantities of ore elements. Devitrification, vapor-phase crystallization, fumarolic alteration, or the formation of lithophysae may free the ore elements from the glass matrix and place them in a form that is readily leached by percolating meteoric waters. Heat from the rhyolitic sheets drives such waters through the system, generally into and up the conduit and out through the early tuffs. Secondary alteration zones (K-feldspar, sericite, silica, clays, fluorite, carbonate, and zeolites) and economic mineral concentrations may form in response to this low temperature (less than 200°C) circulation. After cooling, meteoric water continues to migrate through the system, modifying the distribution and concentration of the ore elements (especially uranium). In this model, the ore elements are derived essentially from the volcanic vent complex itself, although contributions from the underlying magma chamber are not excluded. Plutons and country rocks beneath such vent complexes may themselves contain disseminated, vein, or replacement deposits of U, Th, Be, Sn, Mo, W, Nb, Ta, or associated elements.

CAPPS, RICHARD C., East Carolina Univ., Greenville, NC  
Geology of Rancho el Papalote Area, Chihuahua, Mexico

The Rancho el Papalote area is 35 km north of Chihuahua City and includes Minas Terrazas. The oldest rocks are highly fractured (17% dilation), massive Cretaceous biomicrites, containing rudistid bioherms, that crop out on ridges and peaks surrounded by volcanic rocks of two eruptive periods. The older volcanics are altered rhyolite and/or dacite flows, flow breccias, and minor lithic tuffs. The flow breccias are thickest around Cerro Choloma which may represent a vent. A boulder breccia containing limestone and chert clasts is along the limestone-older volcanic contact. The younger volcanics are rhyolite ash flows with minor basalt and andesite flows. There are at least five cooling units. A 44-m.y. (initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio = 0.7048) old vitric-crystal tuff (22% phenocrysts, sanidine/quartz = 4.5) overlies the older volcanics 1 km west of the ranch buildings. The "Red Platy," an ash-flow rhyolite, overlies the 44-m.y. tuff and is a distinctive marker bed. A small dike cutting the tuff may represent a vent for the Red Platy. The remaining units, vitric tuffs in the eastern part of the area, dip gently east and are cut by northwest-trending normal faults. The first of these ash flows has a basal lithic zone and contains 3% sanidine phenocrysts (Or 45). The next unit is vesicular at its base and has 6% sanidine phenocrysts. The youngest ash flow contains sanidine, quartz, and fayalite phenocrysts (sanidine/quartz = 1, phenocrysts = 4%). A basalt above the Red Platy contains distinctive gabbroic

xenoliths. On the south the older volcanics are overlain by an andesite flow. At Minas Terrazas, copper deposits occur in skarns near a felsic intrusive. These skarns are mainly andradite and contain small amounts of pyrite and chalcopyrite.

CARRAWAY, KENNETH, and PHILIP GOODELL, Univ. Texas, El Paso, TX

Gamma Ray Spectrographic Study of Stream Sediment Samples, Pena Blanca, Chihuahua

A stream sediment survey was conducted in the Pena Blanca uranium district, Chihuahua, Mexico, to determine the applicability of gamma ray spectrometry as an exploration method for volcanogenetic uranium deposits. The Pena Blanca district is 40 km northeast of Chihuahua City, in outflow facies of ash flow tuffs, which are 45 to 35 m.y. old. The mineralization in the district is confined to these ash flows and the underlying Cretaceous limestones.

Stream sediment samples were collected and 500-g, 100-mesh separates were analyzed for eU, eTh, and eK. The samples were also analyzed for both acid extractable uranium and total uranium.

The results of the study show that, with the use of two standard deviations from the mean as an anomalous value, the gamma ray data show anomalies, not over the deposits, but at the break in slope in the streams and at the proximal edge of alluvial fans. Other anomalies were detected by this method, but they were single-sample anomalies. The extractable uranium and the total uranium data show a high degree of correlation (0.92), and also correlate well with the eU values determined for the stream sediments. Uranium being mobilized from the Pena Blanca deposits produced geochemical anomalies at the tops of alluvial fans. This knowledge can be applied to exploration programs elsewhere.

CASTOR, S. B., and M. R. BERRY, Bendix Field Engineering Corp., Spokane, WA

Geology of Lakeview Uranium District, Oregon

About 400,000 lb (181.4 Mg) of  $\text{U}_3\text{O}_8$  have been mined from Tertiary volcanic rocks near Lakeview, Oregon, mostly from uraninite-coffinite ore bodies at the White King mine. At this deposit, tuffs and tuffaceous sedimentary rocks are overlain by a basaltic lahar which is capped by basalt flows. The tuffaceous rocks are intruded by flow-banded rhyolite. Most of the ore is in the rhyolite and adjacent tuffaceous rocks, but uraniferous shear zones cut the lahar, and the basalt is highly altered. Most of the uraniferous rock is silicified and brecciated, and has high As, Hg, Mo, and Pb together with secondary U minerals.

Within 10 km of the mine are nine smaller uranium occurrences, four of which are in, or adjacent to, flow-banded rhyolite domes which are considered endogenous. However, five occurrences are distant from surface exposures of rhyolite: three in volcanic sandstone and tuff near contacts with basalt, and two in a thick sequence of rhyolitic to dacitic ash flows.

At the White King mine and most of the other occurrences, uranium was probably deposited from epithermal fluids released during rhyolite intrusion. These fluids could have migrated some distance from the intrusions along fractures or through permeable sedimentary rocks. Capping basalts, which are cut by the intrusions, appear to have arrested the movement of uranium at several occurrences.