

calcite-aragonite sand sequence to temporal variations in rates of CO₂ production and loss. The model can be used to help predict the spatial distribution of zones of dissolution and cementation in various coastal marine carbonate sediments.

HARMS, JOHN C., Marathon Oil Co., Littleton, CO

Lateral Seals of Small Stratigraphic Traps in Cretaceous Rocks, Western Nebraska

The lateral seals of stratigraphic traps located on monoclines offer some interesting perspectives on problems of hydrocarbon entrapment. On the eastern flank of the Denver basin, oil reservoirs in Cretaceous valley-fill sandstone bodies afford a particularly instructive example. There the updip seal is provided by sandy, silty, and clayey marine facies into which were cut the slightly younger valleys. A subsequent marine transgression covered the entire sequence with clay shale, providing a top seal.

The lateral seal can be visualized as a maze of pipes and capillaries through which the oil phase finds its way, until blocked at each potential escape route by entry pressures that exceed that generated by the buoyancy of the oil column. Cores of the sealing facies show that burrowed silty sandstone is a small-scale maze, where silty laminae shortly block oil invasion. Wave-rippled shallow-shelf sandstone beds offer more continuous conduits, but are lenticular on a scale of a few hundred meters and are separated by shale beds that ultimately baffle oil escape. Cross-stratified shoreline sandstone is also a sealing facies by diagenetic kaolin; stained sets of cross-laminae are enclosed in unstained rock, a testimony to slightly different capillary pressure characteristics.

Careful examination of cores of rock facies can give vital information on the geometric nature of a seal; on how to convert laboratory measurements of capillary pressure using mercury and air to natural oil-brine systems; on critical oil saturation levels that distinguish seal and non-seal rocks; and on the height of hydrocarbon column than may be trapped by a seal.

HARRIS, PAUL M., Gulf Science and Technology Co., Houston, TX

Upper Jurassic Evaporites and Carbonates of Gulf Coast

Shallow-water carbonate rocks of the Upper Jurassic Smackover Formation contain some of the most promising exploration targets of the Gulf Coast. Evaporites of the Louann, the Buckner, and within the Smackover have been important factors in forming the reservoirs.

The Louann, a thick salt lying stratigraphically below the Smackover, localized production and influenced facies patterns and diagenesis. The updip limit of the Louann coincides with peripheral basin faults, so fault traps in the Smackover are to some degree related to salt movement. Salt flowage created highs on the sea floor that localized high-energy oolite bars and, in some places, reef development. Some Smackover porosity was created secondarily during early freshwater diagenesis in association with the highs. Late-stage burial diagenesis in the Smackover, which is also important in forming reservoir porosity, may have been caused by interstitial fluids that were expelled from the Louann during salt compaction.

Anhydrite occurs in the Buckner and Smackover in several forms: (1) nodules, (2) microcrystalline replacement, (3) laminations of blocky crystals, (4) poikilitic cement, (5) poikilitic replacement, (6) fracture fill of blocky or lath-shaped crystals, (7) lath-shaped replacement crystals, and (8) plugging

of oomoldic porosity. The Buckner anhydrites and shales are important seals for Smackover reservoirs. Commonly the Buckner interfingers with the reservoir at the updip limit of a field and as a result of basinward progradation forms an impermeable cap. The Buckner may have influenced early diagenesis of the uppermost Smackover by introducing brines into the lime sands. Thickness maps of the Buckner serve as a valuable exploration tool for the Smackover, because porous facies are associated with Smackover highs underlying the Buckner thins. Anhydrite within Smackover grainstones can locally reduce porosity to form internal seals. The presence of minor amounts of anhydrite in Smackover carbonate rocks can confuse log interpretations on neutron and density log crossplots by causing overestimates of the dolomite percentage and underestimating of porosity.

HARRISON, RAND S., Alberta Geol. Survey, Edmonton, Alberta, Canada

Subaerial Exposure Surfaces: Variations on a Theme

Carbonate sediments are commonly subjected to subaerial exposure and surficial alteration relatively early in their history through real or relative changes in sea level. The resultant diagenetic overprint may be superficial or pervasive, subtle or profound, depending upon the interplay between a number of controlling variables. It is particularly important to realize that, to a first approximation, the nature and extent of near-surface subaerial diagenesis is not exclusively a time-dependent function, but rather commonly reflects the prevailing subaerial (climatic) regime. Because of this, the course of such diagenetic modification is highly variable.

Studies on Pleistocene deposits from localities such as Barbados and Florida demonstrate a tremendous range in exposure-surface fabric. These fabrics encompass lithified but otherwise little-altered horizons, thin but texturally variable calcrete crusts, thick complex caliche profiles, in-situ pseudobreccias, and solution karst with pits and channels of high relief. It is common to find sharply contrasting fabrics developed in close proximity on the same exposure surface.

To recognize such breaks in outcrop or core, it is necessary to maintain a critical appreciation that variability is the rule rather than the exception. There is no question about the importance of such breaks, for they serve as major time-stratigraphic datums, provide insight into basin history, and may either locally enhance porosity or act as substantial vertical permeability barriers, commonly within stratigraphic sequences that otherwise appear to be lithologically uniform.

HART, ORVILLE D., Chevron U.S.A. Inc., Oildale, CA

Geologic Studies of Steam Drives Pay Off in Kern River Field, California

The Kern River field in California currently produces over 100,000 bbl of heavy crude oil per day from several steam-drive projects. Conservatively, recoverable reserves of 600 million bbl remain as the prize of enhanced oil recovery techniques. Operating costs for steam-drive projects are high because for every three barrels of crude produced, one barrel is burned to generate the steam. To effectively operate such projects, geologic conditions of the reservoir must be thoroughly understood. Close coordination between the geologist and the engineer is needed to evaluate and follow steam-drive performance on a pattern-by-pattern basis. Detailed cross sections and isopach maps of the reservoir sands and their overlying