

Recognition of Shelf-Slope Break Along Tectonically Active, Ancient Continental Margins

Tectonically active continental margins include transform, protoceanic-rift, and subduction settings. Shelf-slope breaks in these settings tend to be transient in time and space. Wrench and fault-block systems feature irregularly shaped basins and uplifts that have abrupt vertical movements and facies changes. Subduction systems form elongate basins and ridges in response to the interplay of tectonics and sedimentation.

Because of the nondepositional character of narrow, tectonically controlled shelves, shelf-slope breaks commonly are expressed as unconformities separating shallow-marine/nonmarine and slope/basinal deposits. Alternatively, shelf and basinal deposits may be separated by a biostratigraphically compressed interval, including glauconite and phosphorite, which represents the clastic-starved shelf-slope break. Subtly expressed shelf-slope break deposits of active margins commonly are masked by abrupt sedimentation in adjoining areas along with abrupt facies migration. Consequently, bracketing deeper and shallower marine facies is the usual key to locating ancient shelf-slope breaks.

Basinal facies are distinctive in many wrench and fault-block systems, with two common sedimentary motifs: (1) ponded submarine-fan deposits displaying little proximal-to-distal facies segregation, and (2) clastic-starved sections of laminated (commonly biogenous) sediment reflecting anoxic conditions in isolated silled basins. These facies, together with slump-dominated slope facies, adjoin shelf-slope break deposits.

Variable geometries characterize subduction-controlled shelf-slope breaks, but generally, ancient deposits overlie highly deformed, deep-marine components (trench-slope and trench sediments) and are overlain by relatively undeformed fore-arc deposits. Underlying deposits may have exotic sources and may contain recycled materials. Overlying deposits consist of predominantly land-derived components.

INGLE, JAMES C., JR., Stanford Univ., Stanford, CA

Origin, Depositional History, and Correlation of Miocene Diatomites Around North Pacific Margin

Distinctive Miocene diatomites and genetically related porcellanites form a remarkably widespread lithofacies within bathyal marine sequences around the margin of the North Pacific Ocean from Mexico to Korea. The Monterey Shale of California and the Onnagawa Formation of Japan typify these deposits. Planktonic correlations, radiometric dates, and oxygen isotope records indicate that the initial accumulation of the diatomites commenced in the early middle Miocene (15 Ma), coincident with initiation of a major climatic event marked by massive ice accumulation in Antarctica, steepening of the latitudinal temperature gradient, and increasingly vigorous surface circulation and primary productivity. Microfaunal and sedimentologic evidence demonstrates that the diatomites were commonly deposited in subsiding marginal basins characterized by sills intersecting intensified oxygen minima allowing preservation of laminated muds. Moreover, the diatomites occur within strikingly similar stratigraphic successions that suggest closely parallel tectonic, sedimentologic, and oceanographic development of Neogene marginal basins in this region. Typically, four major events can be recognized in these sequences: (1) deposition of Oligocene-lower Miocene volcanic rocks and continental and/or neritic marine clastics followed by (2) rapid margin subsidence, development of silled basins, and deposition of middle and upper Miocene diatomaceous sediments in essentially empty depocenters, (3)

climatically induced elevation of the carbonate compensation depth and increasing influx of terrigenous debris in latest Miocene-earliest Pliocene time producing a carbonate-poor mudstone facies, and (4) introduction of rapidly deposited wedges of coarse terrigenous clastics during Pliocene-Pleistocene time that cap the underlying diatomites. This widespread and correlative stratigraphic pattern appears to be the combined product of (1) a major middle Cenozoic readjustment of Pacific plate margins resulting in synchronous development of Neogene marginal basins around the North Pacific rim and (2) coincident acceleration of diatom productivity in this region in response to severe deterioration of Neogene climate.

INGRAM, ROY L., and LEE J. OTTE, Univ. North Carolina, Chapel Hill, NC

Peat—Potential Energy Bridge for North Carolina

North Carolina has an estimated 1,000 sq mi (2,600 sq km) of peat land containing about 600 million tons of moisture-free peat. Because North Carolina is deficient in energy resources, there is considerable activity aimed at using this peat as a fuel.

Peat deposits in North Carolina are of three main geologic types representing the accumulation of organic matter in: (1) pocosins—broad shallow depressions on an uplifted sea floor; (2) river flood plains; and (3) Carolina Bays—elliptical depressions of unknown origin.

The largest pocosin deposits are: (1) Albemarle-Pamlico peninsula, 360 sq mi (936 sq km), 210 million tons moisture-free peat; (2) Dismal Swamp, 100 sq mi (260 sq km), 60 million tons; and (3) Croatan Forest, 40 sq mi (104 sq km), 23 million tons. These deposits normally range in thickness from 1 to 8 ft (0.3 to 2.4 m).

River flood-plain deposits are of unknown extent. Peat occurs as lenses in alluvial sands and clays and may attain a thickness of 25 ft (8 m).

Five to six hundred Carolina Bays from 0.2 to 3 mi (0.3 to 5 km) in length are scattered over the coastal plain. Many contain high quality peat up to 15 ft (5 m) thick.

Most North Carolina peat is black, fine grained and highly decomposed, with an ash content commonly less than 5%. Sulfur content is low (median 0.2%), and heating value is high (median to 10,300 Btu/lb moisture-free).

ISAACS, CAROLINE M., U.S. Geol. Survey, Menlo Park, CA

Porosity Reduction During Diagenesis of Monterey Formation, Santa Barbara, California

In the Monterey Formation of the Santa Barbara coastal area, porosity decreases with increased burial diagenesis from an average of 60% in diatomaceous (opal-A) rocks to an average of 10% in rocks bearing diagenetic quartz. In carbonate-free siliceous rocks, porosity was lost principally in two abrupt reductions of 10 to 30% porosity during the two silica phase transformations (opal-A to opal-CT and opal-CT to quartz). Large differences in porosity among interbedded rocks with different silica phases show that porosity losses resulted directly from the two silica phase transformations rather than from specific conditions of temperature and burial depth. Porosity reduction was thus probably due to brief loss of strength during solution-precipitation of silica in the rocks. Similar compositions and different thicknesses of sedimentary

units in adjacent stratigraphically equivalent sections with different silica phases show that most porosity reduction during these transformations resulted from compaction, not from addition of silica.

In calcite-bearing siliceous rocks, the pattern of porosity reduction was similar, and minor additional reduction was caused by silica filling of some foraminiferal tests. In calcite-bearing rocks, with moderate (10 to 40%) silica, only 5 to 10% of porosity was lost during each silica phase transformation, and additional gradual compaction occurred in the interval between the two transformations, presumably in response to increased burial load. Formation of disseminated dolomite, in contrast to the rarer formation of highly cemented dolomite beds and nodules, was accompanied by only moderate reduction in porosity (0 to 10% porosity).

JAMES, NOEL P., Memorial Univ. Newfoundland, St. John's, Newfoundland, and COLIN F. KLAPPA, Gulf Canada Resources, Inc., Calgary, Alberta, Canada

Composition of Particles and Cements in Lower Cambrian Reefs and Precipitation of Metastable Carbonates in Paleozoic Ocean

Bioherms and biostromes of the Lower Cambrian Forteau Formation in southern Labrador and western Newfoundland are rich in skeletal and nonskeletal components and display a wide spectrum of syndepositional and postdepositional cements. Through petrography, cathodoluminescence and microprobe analysis, three types of components can be distinguished: (1) molds filled with iron-rich or iron-poor calcite spar or iron-rich dolomite (archaeogastropods, hyolithids, brachiopods, ?coelenterates, and *Chancelloria*); (2) particles composed of iron-poor calcite and showing original or relic texture (trilobites, salterellids, echinoderms, and sponge spicules); and (3) components illustrating both the above fabrics (ooids, archaeocyaths, and *Renalcis*). Syndepositional cements are (1) rays or botryoids in which each acicular crystal is a spar-filled mold, or (2) rinds of fibrous calcite commonly with fascicular-optic properties.

Comparison with the petrographic characteristics of Phanerozoic and modern carbonates indicates that those Cambrian particles and cements which are now spar-filled molds were originally high porosity aragonite; those without a void stage and retaining original fabric were calcite or Mg-calcite; those with a fibrous habit were Mg-calcite, and those which exhibit two or more of the above characteristics were low porosity aragonite or high porosity Mg-calcite.

These styles of preservation confirm that organisms in Early Cambrian were secreting skeletons with the same spectrum of mineralogies as those to today and that metastable carbonates (aragonite and Mg-calcite) were being precipitated on the shallow sea floor, as they are in the modern ocean. The absence of similar fabrics at other times in the Phanerozoic probably reflects epeiric sea rather than continental margin sedimentation and not a gradual change in seawater chemistry or atmospheric CO₂ through time.

JAMES, NOEL P., Memorial Univ., Newfoundland, St. John's, Newfoundland, and ERIC W. MOUNTJOY*, McGill Univ., Montreal, Quebec, Canada

Fossil Carbonate Platform Margins

The margin of carbonate shelves and platforms is a critical zone. During construction these margins control both the

geometry and the style of sedimentation on the adjacent platforms. After burial they remain a major interface between carbonate and detrital sediments, a preferred site for both hydrocarbon and metallic mineral entrapment. During later deformation the structural style commonly changes across this zone.

Fossil carbonate margins occur in two realms: (1) along the craton margins, bordering Paleozoic and earlier ocean basins that are now fragmented and deformed in major mountain belts or buried beneath Mesozoic and Cenozoic successions along modern continental margins, or (2) around the edges of Paleozoic platforms developed in major basins on large cratons.

Regardless of setting, two different sedimentary styles recur. Rimmed margins, those composed of an elevated rim of bioherms or carbonate sand shoals, exhibit a marked declivity between shelf and basin. This slope is either gentle, gradually changing from shallow into deeper water sediments (depositional slope), or is a steep incline, with little sediment accretion, over which sediments move on their way to the basin (bypass slope). Slope and toe-of-slope sediments in these settings exhibit a full spectrum of hemipelagic ribbon and parted limestones, hardgrounds, slides and slumps, and sediment gravity flows together with submarine erosion. Open margins or ramps, those in which there is no marked shelf break, gradually change from inner shelf to slope strata. The high-energy zone is located well inside the craton margin on the platform or at the strandline. Deeper water shelf limestones are particularly common in these situations while slope deposits are characterized by hemipelagic deposits but fewer examples of gravity flow deposits.

JANSA, LUBOMIR F., Geol. Survey of Canada, Dartmouth, Nova Scotia, Canada

Mesozoic Carbonate Platforms and Banks of Eastern North American Margin

The Jurassic-Lower Cretaceous carbonate platforms and banks form a discontinuous belt extending from the Grand Banks to the Bahamas, a distance of over 6,000 km. Six types of carbonate buildups are recognized that document the variability of depositional, paleo-oceanographic, and tectonic processes on the eastern North American margin. The texture of the carbonates closely resembles recent deposits of the western Great Bahama Bank in that oolitic grainstones were present near the shelf edge, and skeletal, peloid wackestones and mudstones (biomicrites) were deposited in the inner part of the platform. Coral-stromatoporoid and sponge bioherms were rare constituents of the carbonate banks.

The thickness of carbonate buildups progressively increases southward along the margin attaining a thickness of more than 5 km in the Bahamas. The platforms also become younger southward, which is thought to reflect the northward movement of less than 1.5 cm/year of the North American plate. The carbonate platforms were seeded over the continental basement following the taphrogenic period of plate tectonics.

The building of carbonate ramps characteristic of the Early Jurassic began during a transitional period between continental rifting and early drift of the continental plates in the North Atlantic. The second stage in construction of the carbonate platforms and offshore banks proceeded mainly after separation of the continental plates.

JOHNSON, GARY D., South Dakota Geol. Survey and Univ. South Dakota, Vermillion, SD