

carbonate rock. The primary matrix porosity averages about 3%. However, weathering during the early Tertiary enlarged fractures and previously existing porous zones to a depth of 100 to 150 m below the top of the carbonate rocks. This secondary porosity, in combination with an extensive fracture network, has converted the otherwise dense carbonate rock into a commercially exploitable reservoir.

A contour map on the top of the eroded Mesozoic carbonate reservoir defines a closure 11 by 2.5 km. The field is elongated parallel to and bounded by Miocene faults with an overall configuration of a rounded limestone ridge.

The ridge is covered by middle Miocene organic rich shales, the oil source. These and younger shales cap the accumulation.

The use of a paleogeomorphic model aids the interpreter in mapping data in the Casablanca area which otherwise would seem uninterpretable, or at least difficult to interpret. The model may be of use in other carbonate areas.

WEAVER, FRED M., Exxon Production Research Co., Houston, TX, and RICHARD E. CASEY and ANNA MARIA PEREZ, Rice Univ., Houston, TX

Stratigraphic and Paleo-Oceanographic Significance of Early Pliocene to Middle Miocene Radiolarian Assemblages from California to Baja California

Comprehensive studies of radiolarian faunal assemblages recovered from outcrop sections at Centerville Beach, Newport Back Bay, Palos Verdes Hills, and Chalk Hills in California and at Bahia Tortugas and Maria Madre Island, Baja California, provide the data base for regional paleo-oceanographic and stratigraphic analyses of Monterey and associated siliceous sediments of middle Miocene to early Pliocene age within Neogene California marginal basins.

The sequential succession and development of characteristic radiolarian assemblages and the temporal and spatial variance in environmentally sensitive species in Luisian, lower and upper Mohnian, and "Delmontian" sediments provide significant insight to regional paleo-oceanographic evolution. Documentation is provided by interpretation of paleoclimatic trends, variation in sediment accumulations rates, changes in water-mass interaction and distribution, and upwelling and productivity events.

Stratigraphically important species of radiolarian genera such as *Theocorys*, *Lamprocyrtis*, *Stichocorys*, *Diartus*, *Eucyrtidium*, *Cyrtocapsella*, and *Botryocyrtes* are used to differentiate potential biohorizons which are useful for inter- and intra-basinal and broad regional correlations. Within key intervals, radiolarians provide accurate identification of the low latitude *Dorcadospyrus alata* and *Diartus petterssoni* Zones in lower Monterey formation sediments.

WEIMER, ROBERT J., Colorado School Mines, Golden, CO

Relation of Unconformities, Tectonics, and Sea Level Changes, Cretaceous of Western Interior, United States and Canada

Intrabasin tectonics have influenced patterns of deposition and geographic distribution of major unconformities within the Cretaceous of the Western Interior. Eight major regional to subregional unconformities have been identified. Five of these have been related by previous workers to sea level changes and to well-documented regressive-transgressive cycles.

New studies of recurrent movement on basement-controlled

fault blocks suggest a synchronous relation among fault block movement, sea level changes, and unconformities. Which fault blocks moved on the basin floor, and when, can be explained by stress fields generated by direction and rates of plate motion. Unconformities associated with north-northwest fault trends are caused by more westerly movement, and those associated with east-northeast trends by more northerly plate motion. Expansion of the Gulf of Mexico, the Atlantic Ocean, or the Arctic Ocean during these plate motions may account for associated sea level changes. The 81 to 82 m.y. unconformity and shoreline regression in the Western Interior and synchronous volcanic events on the northern margin of the Gulf of Mexico illustrate the relations.

Uncertainty exists in dating many of the unconformities. However, by use of the time scale of Obradovich and Cobban, the approximate dates for unconformities are estimated as follows: (1) late Neocomian to early Aptian, >100 m.y.; (2) late Aptian-early Albian, 100 m.y. ±; (3) Albian, 96 to 97 m.y.; (4) early Cenomanian, 93 m.y. ±; (5) Turonian to early Coniacian, 87 to 88 m.y.; (6) late Coniacian-early Santonian, 81 to 82 m.y.; (7) late Campanian 71 to 74 m.y.; and (8) late Maestrichtian, 64 to 69 m.y.

Several billion barrels of oil have been found in sandstones associated with unconformities in the Cretaceous. Future stratigraphic trap exploration will be guided by a knowledge of tectonic influence on sedimentation during sea level changes and how these factors controlled distribution of source rock, migration patterns, reservoir rock, and seal.

WELDER, FRANK A., U.S. Geol. Survey, Meeker, CO

Ground-Water Potential for Oil Shale Development in Northwestern Colorado

Rocks in northwestern Colorado contain large amounts of oil shale which constitutes perhaps the richest hydrocarbon resource in the United States. Efforts to develop oil shale will increase demand for water in a region where surface water is fully appropriated. To meet additional water needs associated with industrial and population growth, sources of ground water need to be investigated.

It has been 15 years since investigators determined that large quantities of ground water occur above, within, and below rich oil shale deposits in the Eocene Green River Formation in the Piceance basin of northwestern Colorado. Estimates of the amount of ground water stored in the Piceance basin are as much as 25 million acre-ft. The specific conductance of ground water discharged during the drilling of 24 test holes ranged from 100 to 50,000 micromhos per cm at 25°C.

Another potential major source of ground water in northwestern Colorado may be the Leadville Limestone of Mississippian age. Solution cavities in the outcrop of the Leadville Limestone in northwestern Colorado indicate that the formation may store and transmit large quantities of water. Where fractured and near the surface, the Leadville Limestone has been exposed to ground-water movement, resulting in the development of solution cavities that have enhanced the hydraulic conductivity and storage capacity of the aquifer. Where the Leadville is exposed on or near various structural uplifts in northwestern Colorado, the opportunity for ground-water recharge, movement, and storage may be extensive. Other potential aquifers such as the Dakota Sandstone and the Entrada Sandstone are also under consideration.

WELTE, D. H., and A. YUKLER, Kernforschungsanlage Julich GmbH, Julich, West Germany