

Three boreholes in New York State were investigated, all of which had porosity, sonic velocity, and electrical resistivity well-log data available. Thermal conductivities of drilling chips were measured on a needle probe apparatus. These data were correlated with the geophysical well-log data using multiple linear regression statistics.

An empirical relation for eastern New York State is $K = -42.5\phi + 0.31 \times 10^{-1} \Delta t + 0.12 \times 10^{-3} \alpha + 3.24$ where K is thermal conductivity in watts/m°C, ϕ is porosity in percent, Δt is the interval transit time in $\mu\text{sec}/\text{ft}$, and α is electrical resistivity in ohms m^2/m . A goodness of fit of 0.885 and a multiple correlation coefficient of 0.941 indicate that the technique can be used for analysis of the variation in temperatures in the subsurface.

The data fell into two groups: (1) clastics with thermal conductivities ranging from 3.39 w/m°C to 3.92 w/m°C; and (2) nonclastics with thermal conductivities ranging from 3.04 w/m°C to 4.89 w/m°C.

The evaluation of temperatures in sedimentary basins can be greatly enhanced if empirical relationships are established for particular areas.

MONCURE, GEORGE K., RICHARD W. LAHANN, and ROBERT M. SIEBERT, Conoco Inc., Ponca City, OK

Diagenesis of Sandstone/Shale Package, GCO/DOE No. 1 Well, Brazoria County, Texas

Petrographic analysis of closely spaced sandstone samples (GCO/DOE No. 1 well) of a sandstone/shale package from the Frio Formation (Oligocene) indicates that sandstone reservoir quality was influenced by shale diagenesis. Three alteration zones at increasing distance from the sandstone/shale contact are observed. This zonation may be explained as follows.

Organic and inorganic maturation processes modified shale fluids which, upon expulsion into the sand, resulted in the precipitation of thin, isopachous chlorite grain coatings in the contact sand. Late in the chloritization process and thereafter, unstable framework grain silicates began to dissolve within the sand. We believe that aluminum from framework grains was removed from the contact zone by mobile organic complexes. Silica released from grain dissolution reprecipitated as quartz cement. This contact zone is about 1-ft (0.3 m) thick.

As fluids passed into the second zone (about 1-ft (0.3 m) thick) the sandstone framework grain leaching continued but to a lesser degree. Kaolinite was produced from internal mass sources and from aluminum imported from the first zone. The net addition of alumina from the contact zone prevented development of quartz overgrowths in this zone and the third zone.

In the third zone, dissolution of framework grain silicates was least thorough because of greater distance from the shale and alteration appears to have been aluminum conservative.

MONTAGGIONI, LUCIEN, Univ. Francaise de l'Ocean Indien, Ste. Clotilde, France

Dolomitization of Pleistocene Reef Sediments by Magnesium Leaching Out of Overlapping Volcanics, Mauritius Island, Indian Ocean

A hole drilled through basaltic rocks on the west coast of Mauritius Island encountered 3 m thick, partly dolomitized reefal deposits of middle Pleistocene age, 120 m below mean sea level. Above this carbonate bed is a sequence, about 50 m thick, of strongly weathered basalts and paleosols.

The degree of dolomitization increases toward the overlying basalt. Low-magnesian calcite and dolomite are distributed along

the core as follows: (1) entirely dolomite 0.5 m downward from the volcanic rock, (2) from 0.5 to 2 m below the base of the basalt, both calcite and dolomite, and (3) below 2 m only calcite. High calcium/magnesium ratios (1.2 to 1.4) indicate that protodolomite has replaced micrites (as 0.5 to 3 μm crystals) or sparites (as 10 to 30 μm , subhedral to euhedral crystals) and fills pores created by the dissolution of calcite, demonstrating the near contemporaneity of calcite solution and dolomite precipitation. The isotopic composition of the dolomite ($\delta\text{O}^{18} = +0.1$ to $+3.6\text{‰}$, $\delta\text{C}^{13} = -1.7$ to $+2.00\text{‰}$) and its relatively high strontium content (>500 ppm) suggest the system initially must have been partly closed in order to retain the O^{18} , C^{13} , and strontium of reef carbonates during dolomitization.

The calcium/magnesium ratio of the volcanic rocks decreases from the upper to the lower part of the series (0.66 to 0.075), confirming that higher magnesium contents present in the uppermost reef layers are derived from weathered basalts.

The chemistry of the dolomite and associated volcanic rocks strongly suggests that dolomitization was probably caused by leaching of magnesium out of the volcanic rocks and redistribution within carbonate sediments by descending waters.

MOORE, MICHAEL G., Exxon Co., U.S.A., New Orleans, LA

Depositional Environments and Processes of Calico Bluff Formation (Carboniferous), East-Central Alaska

The Calico Bluff Formation (Carboniferous) is an interbedded limestone and shale unit which is exposed along the Yukon River in east-central Alaska. A prograding submarine-fan model explains the observed lithofacies succession and sedimentary structures.

The rocks of the Calico Bluff Formation are subdivided into five lithofacies: (1) coarse-grained bioclastic limestone, (2) fine-grained bioclastic limestone, (3) platy micritic limestone, (4) calcareous shale, and (5) siliceous shale. Shallow-water fauna in the bioclastic limestones contrast with deep-water goniatites, straight cephalopods, and brachiopods found in the platy micritic limestone and calcareous shale. Sedimentary structures include graded bedding, sole marks, Bouma sequences, and soft-sediment deformation.

Depositional processes of the Calico Bluff Formation range from basinal pelagic sedimentation to sediment gravity flows. Both proximal and distal turbidites are recognized. Distal turbidites represent deposition in a lower-fan environment, whereas proximal turbidites are deposited in a mid-fan environment. Several thickening and coarsening upward cycles are recognized in outcrop. Individual cycles may represent shifting suprafan depositional lobes. Paleocurrent measurements indicate direction of transport to the south-southeast.

Micritic beds are probably calcareous pelagic oozes that have been transported downslope by turbidity currents and deposited in the basin. Calico Bluff Formation shales were deposited by suspension and weak suspension currents.

MORGAN, KEN M., Texas Christian Univ., Ft. Worth, TX, and DON MORRIS-JONES, Environmental Research Inst. Michigan, Ann Arbor, MI

Use of Computer Enhanced Landsat Imagery as Planning Tool for Resource Exploration

Computer-processed NASA Landsat data were used as an exploration planning tool in four potential hydrocarbon-bearing areas in the United States and Israel. The project areas are located in Montana, Washington, west Texas, and the Dead Sea (Israel).

Structural, stratigraphic, and geomorphic analyses have been performed to locate geologic anomalies. Image analysis provides a better understanding of the regional stress-strain relations for tectonic correlation.

Digital Landsat (satellite) data were processed to produce a variety of images (i.e., edge-enhanced, high pass filter, false color, and ratio). Some of the images were geometrically corrected with map controls and nonlinear deconvolution resampling techniques (coverage 13,000 mi²; 33,670 km²). This helped facilitate more detailed mapping, interpretation, and data integration. These specially processed images have been used to map surface geology, lineament systems, and tectonic anomalies in relation to subsurface geologic and geophysical data.

Project areas are defined in terms of tectonic genesis, structural trends, and hydrocarbon potential. Numerous exploration targets and several modes of hydrocarbon entrapment were identified by geologists at the TCU Center for Remote Sensing and Energy Research. This information is being used by various companies for planning their seismic programs in these frontier drilling areas.

MORROW, DAVID W., Inst. Sedimentary and Petroleum Geology, Calgary, Alberta, Canada

Manetoe Facies—Gas-Bearing Late Diagenetic Dolomite of Northwest Territories, Canada

The white coarsely crystalline space-filling and replacement, late-diagenetic Mississippi Valley-type dolomite of the Manetoe facies occurs across a broad area of at least 15,000 sq km in the southern Mackenzie Mountains of the Northwest Territories. Through most of this area it is stratiform and confined to a thin (<100 m thick) stratigraphic interval under a shale unit within a lower Paleozoic carbonate sequence. Some large vertical developments of the Manetoe facies, such as at the Kotaneelee and Pointed Mountain gas fields, occur in the eastern part of this region where the overlying shale of the Headless Formation is thin. These large dolomite masses have a core of dolomite-cemented breccia and are surrounded by a halo of replacement dolomite. Solution-collapse breccias and large solution cavities are common throughout. Quartz and bitumen are the final vug infillings. The pronounced curvature of Manetoe dolomite crystal faces is similar to that displayed by many Mississippi Valley-type dolomites.

These dolomites are nearly stoichiometric with a mean of 51 mole % CaCO₃, and they exhibit a high degree of cation order. The range of carbon isotope values (+1.33 to -2.99 δ C¹³PDB) and a sodium concentration of ~100 to 350 ppm are typical for this type of dolomite. But the range of oxygen isotope values (-8.03 to -17.33 δ O¹⁸ PDB), the extremely high strontium content of ~200 to 1,000 ppm, and an iron content less than 120 ppm is atypical and must reflect precipitation from a medium of unusual composition, enriched in strontium but depleted in iron and in the O¹⁸ isotope.

MORTON, JOHN P., and LEON E. LONG, Univ. Texas at Austin, Austin, TX

Rb-Sr Dating of Illite Diagenesis

In the Woodford Shale (Upper Devonian), apparent Rb-Sr ages decrease as clay grain size decreases, which in turn correlates with increasing abundance of diagenetic illite. Analyses of the fine-clay size fraction (<0.2 μ) from widely spaced wells in the Delaware basin of west Texas, plot on a single isochron indicating an age of 300 ± 4 m.y. (Middle Pennsylvanian). At this time the Woodford was buried only 200 to 600 m; consequently diagenesis

must have been triggered by a circumstance other than deep burial. Possibly diagenesis was accomplished by hydrothermal fluids moving toward the craton out of the Ouachita geosyncline, which at that time was experiencing horizontal compression. These fluids may have been responsible for petroleum migration and lead-zinc mineralization.

In the Frio Formation (Oligocene) of the Texas Gulf Coast, samples of fine clay-size material (<0.06 μ) from the 3 to 5 km depth interval in a single well also provide a well-defined isochron corresponding to 21.6 ± 2.2 m.y. Burial here was possibly so rapid that transformation of smectite to illite approximated an episodic event over the entire depth interval. Alternatively, because the sediment is geopressed, the age might record the time of geopressure development which was accompanied by a rapid rise in temperature.

Clay diagenesis at the surface is illustrated by a paleosol developed on Pennsylvanian shale in the Llano uplift of central Texas. The paleosol was buried by Cretaceous basal conglomerate and records the time of marine transgression 119 ± 3 m.y. ago. Constituents of the shale were degraded by soil-forming processes which erased previous isotopic memory, then reconstituted by coming in contact with marine water. This field relation offers a new way to date directly a time of sedimentary deposition.

MOSHIER, STEPHEN O., Mobil Producing Texas and New Mexico, Houston, TX, and DOUGLAS W. WAPLES, Mobil Research and Development, Dallas, TX

Was the Mannville Group the Source for Alberta's Heavy Oils?

The Western Canadian basin hosts about 12 billion bbl of conventional oil in Devonian to Cretaceous reservoirs. Lower Cretaceous heavy-oil sands contain 1,300 to 2,600 billion bbl in place. They represent the biodegraded remnants of supergiant conventional deposits, the source for which has been thought to be mature rocks of the equivalent-age Mannville Group. This work shows, however, that the Mannville rocks alone are incapable of generating the required volume of hydrocarbons.

Volume of hydrocarbons generated in the Mannville under central Alberta was calculated by combining measured geochemical and geologic data with a model (modified from Lopatin's method) for thermal maturation. Original hydrocarbon generative capacity of the Mannville rocks was calculated from geochemical analyses of immature samples. Using average values for TOC (1.3%) and Rock-Eval Hydrogen Index (100 mg HC/g TOC), maximum hydrocarbon generation per unit volume of source rock was calculated. The maturation model was then employed to estimate the extent to which maximum yield has been achieved.

Total volume of source rock in the basin was obtained from isopachs of Mannville shale. Multiplication of actual oil generation per unit volume by source rock volume gave a generated volume of 450 billion bbl. Inclusion of oil generated in the Foothills belt would less than double this number. These calculated values are exceedingly optimistic, however, because they ignore inefficiencies in expulsion and migration. It is therefore clear that the Mannville Group cannot be the major source of the heavy oils. Dominant contributions probably come from Paleozoic and other Mesozoic rocks.

MROSZCAK, V. E.,* D. R. HORN,* and D. S. TURNER, Dome Petroleum, Calgary, Alberta, Canada

Petroleum Geology of Central Beaufort Sea, Northwest Territories, Canada