

A recently drilled Cambrian test well in Waterford Township, Pennsylvania, penetrated Precambrian basement at a total depth of 7,342 ft (2,238 m). The well site was determined on the basis of structural contour mapping of the top of the Queenston Shale. This work has led to the definition of a conspicuous structural high (anticline, horst-block?) characterized by about 250 ft (76 m) of relief. The steeply inclined southeast limb of the structure may control gas production within the younger Paleozoic formations and ultimately in older formations (e.g., Cambrian Gatesburg and Potsdam). Interpretation of structure, lithologic characteristics, and log analysis reveals two target formations—the Gatesburg Sandstone and the Potsdam Sandstone. The upper sandy section of the Gatesburg Sandstone, a clean, yellowish-white quartzarenite of Late Cambrian age was penetrated at a depth of 6,280 ft (1,914 m) and was found to be water saturated between 6,340 and 6,370 ft (1,932 and 1,942 m). The productive zone is targeted between 6,415 and 6,447 ft (1,955 and 1,965 m) and contains approximately 13 ft (4 m) of pay that averages 8% porosity. Silica is the dominant cementing agent. The Potsdam Sandstone was encountered at a depth of 7,143 ft (2,177 m), just below a dolomitic interval of the lower Gatesburg. The productive zone ranges from 7,250 to 7,294 ft (2,210 to 2,223 m) and is characterized by 10 ft (3 m) of sand that averages 7% porosity. Calcite is the dominant cementing agent within the target zone.

MOTIA, BENJAMIN D., and ARTHUR E. BURFORD, Univ. Akron, Akron, OH

Stratigraphic Relation of Silurian Sandstone in Part of Summit and Portage Counties, Ohio

The Silurian sandstones of northeastern Ohio, informally termed white "Clinton," red "Clinton," and stray "Clinton," in ascending order, have been described by various investigators as belonging to a major Silurian delta complex sourced from an easterly direction. Detailed study of geophysical logs of a limited area in Summit and Portage Counties supports this interpretation. Isopach maps, sand thickness maps, and sandstone-shale ratio maps show a pattern of probable deltaic distributary streams flowing generally from east to west, bifurcating at places to probable crevasse splays, and terminating westerly in nearshore beach and marine bar sands. The units are probably marine regressive. In ascending order, the white Clinton represents partial bars, beaches, and delta front; the red Clinton represents poorly drained swamp and/or upper delta or fluvial environments; and the stray Clinton may be marine transgressive representing environments similar to the white Clinton.

MURIN, TIMOTHY M., and JACK DONAHUE, Univ. Pittsburgh, Pittsburgh, PA

Upper Devonian First Bradford Formation of Southwestern Pennsylvania: Environment of Deposition and Factors Affecting Gas Production

Core descriptions and thin-section analyses from five wells in southwestern Pennsylvania, in addition to regional cross sections and subsurface maps, provide the basis for subdividing the Upper Devonian First Bradford formation into three distinct members. These include upper, middle, and lower sandstones. Although individual members vary in geometry and areal distribution, similar characteristics in sedimentation persisted over a distance of 125 km (78 mi) along strike and 25 km (16 mi) along dip. The sandstones are generally very fine to fine grained, exhibit upward coarsening, and are calcite cemented. Depositional structures range from planar bedding to cross-laminations, which include flaser, wavy, and lenticular stratification. Bioturbation and disseminated organic debris are common.

Isopach maps show the sandstones to be strike-oriented linear ridges and/or pods 1-3 km (0.6-1.9 mi) wide and 3-20 km (1.9-12.4 mi) long. Thicknesses rarely exceed 10 m (33 ft). Regionally, sand percentage for the overall interval (approximately 46 m or 151 ft) decreases west and southwest of Clearfield County where only the lower sandstone is preserved in parts of Armstrong and Westmoreland Counties. The postulated depositional environment is a shallow marine, inner to mid-shelf bar facies. The sediments accumulated as a result of current reworking; in places, deposition was controlled by shelf topography. Sediments were probably derived from a deltaic complex in the vicinity of Clearfield County.

Gas production is confined to the stratigraphic limits of the clean sandstones and is locally enhanced by a high structural position. This "combination trapping" permits the segregation of gas and water and is evidenced by production in Westmoreland and Fayette Counties. Core porosities average 6.7%; permeabilities, with an effective overburden pressure of 1,000 psi (6,895 kPa), average 0.04 md. Production capabilities are ultimately affected by various diagenetic influences, which include compaction, cementation, authigenesis, replacement, and leaching.

NURMI, ROY D., Schlumberger-Doll Research, Ridgefield, CT

Basin Analysis and Petroleum Potential of Michigan Basin: Deposition and Subsidence History from Middle Ordovician (Trenton Formation) to Early Devonian

The history of the Michigan basin (Early Ordovician to Early Devonian) is that of a nonuniformly subsiding basin, with the Michigan basin, at times, nearly disappearing as either a topographic feature or a depositional center. This history is interpreted from the analysis of lithostratigraphic units, time stratigraphic features, and log formats (term by J. Forgotson). These units are defined for wells throughout the Michigan basin, and they extended eastward into the Appalachian basin. The definition and thickness mapping of these lithostratigraphic units and formats are accomplished using well cuttings, cores, and wire-line geophysical well logs. Furthermore, these units are correlated to both outcrops and stratigraphic type sections for areas adjacent to the Michigan basin. From these data, it is possible to interpret the major aspects of both the subsidence and depositional history of the basin. During deposition of both the Trenton limestones and Early Silurian carbonates and shales, the Michigan basin behaved as if it were part of the greater Appalachian basin, whereas prior to the deposition of the Trenton (Middle Ordovician) and during Middle and Late Silurian, the Michigan basin was an entity separate from, and with an apparent structural independence of, the greater Appalachian basin. However, even during times of Michigan basin structural independence, the stratigraphic and sedimentologic characteristics of these two basins were closely related.

The structure and topography of the Trenton prior to the deposition of the Utica Shale was mapped throughout Michigan to provide insight into the nature of petroleum entrapment in the Trenton formation. The structural entrapment of petroleum in southeast Michigan contrasts with the combination diagenetic to structural Albio-Scipio trend of south-central Michigan. Evidence is available that more of these two types of traps occur in unproducing areas of the Michigan basin.

O'NEIL, CARON, and THOMAS ANDERSON, Univ. Pittsburgh, Pittsburgh, PA

Lineament Analysis near French Creek, Northwestern Pennsylvania

We are attempting to characterize the nature of linear topographic features in the vicinity of French Creek, northwestern Pennsylvania. Straight valleys that trend north to northwest and that are partially filled with glaciofluvial deposits distinguish prominent lineaments in Crawford, Venango, and Mercer Counties. The traces of these linear valleys are 1-15 mi (1.6-24 km) long and 10-4,000 ft (3-1,220 m) wide. The vertical extents of the lineaments are assumed to be proportional to their lengths. We assume that the topographic lineaments mark easily eroded zones coincident with fractures.

The French Creek lineaments have been plotted on a Landsat photo, glacial map, regional map, and topographic maps. Subsurface structural and isopach maps may show facies changes or anomalous thickness across the lineaments. Geochemical surveys determine if there is any enhanced vertical migration of hydrocarbons along a lineament, and very low-frequency magnetic surveys would indicate any increase in groundwater concentration associated with the fracture zones. Well log data on initial production and breakdown pressures have been compared to lineament positions for possible correlations.