

all the basin's oil in addition to having its own potential carbonate reservoirs. The faulted margins of the basin fed a system of alluvial fans, sand flats, and mud flats. Alternating dry and rainy periods regulated the size and nature of contemporaneous basinal alkaline lakes. Dry periods corresponded to contracted playa lakes with ostracod carbonates and euxinic shales; rainy periods corresponded to expanded pluvial lakes with pelecypod banks. Subaqueous intrusions of basaltic magma generated hyaloclastites with kerolitic ooids and hyalotuffs.

Petrographic analysis reveals 5 diagenetic stages: (1) syndepositional alteration of lithoclasts to trioctahedral smectites; (2) early dolomitization, early silicification, and cementation by bladed-rim calcite and zeolites; (3) freshwater-vadose dissolution of bioclasts and lithoclasts, freshwater-phreatic sparite cementation, and neomorphism; (4) mixed saline-freshwater silicification; and (5) burial with compaction, late dolomitization, and partial conversion of smectites to illite.

Pelecypod limestones with primary interparticle, secondary intraparticle, moldic, and moldic-enlarged porosities are the potential reservoirs. Ideal conditions for porosity generation and preservation were subaerial exposure followed by rapid lake expansion and burial.

BIEBER, DAVID W., Independent Geologist, Englewood, CO

Exploration Model for Unconformity-Related Hydrocarbon Accumulations in Cherokee Group of Western Kansas

The sandstones of the Desmoinesian Cherokee Group in western Kansas are important hydrocarbon producers. The Start oil field in Rush and Ness Counties is an example of an unconformity-related Cherokee accumulation from which an exploration model can be made. In this field, the upper Cherokee member is economically important and is interpreted to be a marine unit deposited on the distal portion of an alluvial plain. Traps and reservoirs in this unit were formed by winnowing of clay and silt-sized material from sediments deposited on the crests of paleohighs.

Four maps are useful in exploring for upper Cherokee hydrocarbon accumulations such as Start. An isopach map of the Cherokee Group is useful for locating thins that coincide with paleohighs on the basal Pennsylvanian unconformity. An isopach map from the Cherokee top down to the first sandstone porosity is useful. "Thins" of this interval define areas where wave and current action have winnowed finer material from sands. Closed anticlines on a Cherokee structure map are areas where Cherokee reservoirs are likely to be oil bearing rather than water bearing. An isopach map from the Cimarronian Stone Corral anhydrite top down to the Missourian Lansing Group top is also useful. "Thins" of this interval correspond to paleohighs on the basal Pennsylvanian unconformity. This interval can be picked from seismic records. Prospective areas occur where isopach thins of Stone Corral to Lansing, of Cherokee Group, and of Cherokee top to first sandstone porosity coincide with Cherokee anticlinal structure.

BINDER, ALAN B., NASA, Johnson Space Center, Houston, TX

Lunar Resources—Enabling Factor in Industrialization of Near-Earth Space

The industrialization of near-earth space has already begun during the last few space shuttle flights and will continue at an ever-accelerating pace as the first United States space station is built in the next 8 yr. However, the economic return from near-earth space industry is limited by the high cost per kilogram of launching into orbit the structural elements needed to build the space stations and the raw materials that are to be made into products in them. This limiting factor can be overcome if the moon is used as a source of the material needed to build the space structures and as the source of the raw materials needed in the processing. First, O, Si, Mg, Fe, Ca, Al, and Ti are the major constituents (> 1% by weight) in lunar rocks, and can be obtained directly from them by one of several proposed processes. Of these, O is needed as a rocket fuel, Si for making solar cells to generate space-station electricity, and light weight Mg, Al, and Ti to make structural elements for the space stations. Second, the rocket fuel per kilogram of payload needed to reach low earth orbit from the moon is 68% of that needed from the earth's surface, assuming that decelerating into earth orbit is achieved by a rocket maneuver. The amount of fuel is reduced to 15% if orbit is achieved by aero-braking. If the payload is launched from the moon by a "mass driver" and aero-braking is used, the

cost in the rocket fuel needed to reach low earth orbit is reduced to nearly zero.

BINGHAM, M. P., ARCO Exploration Co., Denver, CO, and R. H. BAILEY, Northeastern Univ., Boston, MA

Depositional Environment of Lower Cambrian Limestones, Nahant, Massachusetts

About 130 m (427 ft) of Lower Cambrian Weymouth Formation is exposed at Nahant, Massachusetts, 11 km (7 mi) northeast of Boston. Several beds of white to light-gray, fossiliferous limestone, up to 3 m (9.8 ft) thick, occur in a sequence of dark, very thinly bedded argillite. Portions of the argillite contain altered and chertified carbonate nodules. Limestone beds contain irregular, very thinly laminated chert layers with structures characteristic of silicified laminate stromatolites.

The limestone is comprised of 4 microfacies: (1) thinly bedded unfossiliferous micrite, (2) irregular intraclasts surrounded by sparry cement or biomicrite, (3) biomicrite (wackestone to packstone) containing small shelly fossils, primarily hyolithids, and (4) biosparite (hyolithid grainstone).

Conoidal hyoliths do not show a strong preferred current orientation on slabs and peels, but are often irregularly disposed with long axes at high angles to bedding. Biomicrites (packstones) and biosparites occasionally overlay irregular scour surfaces. Irregular bioclastic-rich pockets are surrounded by and grade into micrite or biomicrite.

A shallow subtidal, partially protected shelf of platform environment best explains the textural and bedding characteristics of limestones. Storm events are recorded as bioclastic-rich and intraclastic sediments over scour surfaces. Irregular cyclic repetition of microfacies and lack of progressive shoaling or deepening during carbonate deposition suggest that limestones represent periods of stillstand at relatively low sea level positions. Carbonate deposition ceased when extrabasinal mud input increased, possibly during episodes of rapid sea level rise.

BISHOP, WILLIAM F., Tenneco Oil Exploration and Production, Houston, TX

Eocene and Upper Cretaceous Carbonate Reservoirs in East-Central Tunisia

Regionally, well-defined belts of lowest Eocene (Ypresian) Metlaoui carbonates trend northwest-southeast. On the northeast is an open-marine, basinal facies of planktonic foraminiferal micrite and marl. Thick bars of shallow marine nummulitic wackestone, packstone, and grainstone trend northeastward at an angle to the paleoshelf. Lagoonal or supratidal carbonates are widespread between the shelf deposits and thick evaporites that crop out in intermontane basins.

The reservoir is confined largely to nummulitic packstone, and visible effective porosity is best developed between forams in zones filled with sand-size debris where secondary solution-enlargement has occurred. Porosity within nummulite chambers, while abundant, is ineffective, although a few open fractures were observed in cores. This lithology tested oil in 2 recent wildcats and is a commercial reservoir at Sidi El Itayem and Ashtart fields.

Distribution of Zebbag carbonates of Late Cretaceous (Turonian) age is more complex. A northwest-southeast-trending platform is bounded on 3 sides by basinal shale and micrite with planktonic forams which grade into a transitional facies of micrite and wackestone that shows some evidence of shallow-water deposition, such as dolomitization, bioclasts, rare ooliths, etc. Predominately back-reef and lagoonal bioclastic wackestones and packstones occur in narrow belts, apparently controlled at least locally by block faulting. The rest of the platform lithology comprises mostly dolomite and dolomitic limestone.

The most significant porosity is interparticle (generally solution-enlarged) in foram packstones, but intraparticle porosity in forams and rudists commonly enhances the reservoir. Intercrystalline porosity in dolomitized zones is common, and fenestral porosity occurs in a few places. All are modified by nonfabric-selective channel and vuggy porosity and in some instances by fractures.

BLACKWELL, DAVID D., Southern Methodist Univ., Dallas, TX

Measurement of Contemporary Thermal Properties in Sedimentary Basins