

Mesozoic sediments are a mixture of clastics and carbonates. Cenozoic sediments are predominantly limestones, dolomites, and anhydrites with some sandstones and shales. Environments of deposition range from continental to deep marine.

The Cyrenaican area has not been heavily explored and, until recently, no commercial hydrocarbons were found. Drilling on surface structures of some of the first wells in Libya resulted in one Devonian gas well. A reported 5,600 BOPD Cretaceous discovery offshore Benghazi in mid-1984 demonstrates that hydrocarbon potential exists where thick sediments have been preserved.

ROTHAMMER, CHRISTINE M., and LOWELL R. MORRISON, California State Univ., Northridge, CA, and STEVEN L. WARKENTIN, Texas Christian Univ., Fort Worth, TX

Shoreline, Grain-Size, and Total-Carbon Distribution Changes Before and After Hurricane Alicia, Galveston Island, Texas, 1983

Shoreline, grain-size, and sediment total-carbon changes were monitored, on a monthly basis, on three Galveston Island beaches, from January through December 1983. The study area included: (1) East Beach, obstructed by groins and a seawall; (2) Galveston Island State Park, obstructed by fences artificially stabilizing the dunes; and (3) West Beach, an unobstructed beach.

Beach profiles revealed the effects of beach obstruction, such as erosion and undercutting at East Beach, and truncation of the dunes at Galveston Island State Park. Approximately 20 m of expansional cutback occurred on the beaches after Hurricane Alicia hit on August 18, 1983.

Contour maps of grain-size and total-carbon distributions reflect the movement of beach sand by either onshore-offshore transport during low-energy periods, or longshore, edge-wave transport during high-energy periods.

Statistical analyses revealed a small variation in grain size throughout the year. There were well-defined times of either no correlation or strong correlation between total carbon vs. mean grain size, skewness vs. mean grain size, kurtosis vs. mean grain size, total carbon vs. percent sand, total carbon vs. skewness, and skewness vs. kurtosis. Strong correlation was found in response to high-energy events, whereas no correlation was found in response to low-energy events.

Galveston Island is undergoing net erosion and appears to be in a metastable state, still capable of responding to oceanographic conditions. The economic effects of Hurricane Alicia include considerable loss of the shoreline and destruction of property. Beach nourishment appears to be the only economically feasible solution to counteract the extensive erosion.

ROWLEY, D. B., A. L. LOTTES, and A. M. ZIEGLER, Univ. Chicago, Chicago, IL

North America–Greenland–Eurasian Relative Motions: Implications for Circum-Arctic Tectonic Reconstructions

The Mesozoic-Cenozoic tectonic evolution of the Circum-Arctic region is based on constraints imposed by (1) relative motion histories of the three major plates (North America, Greenland, and Eurasia) and a number of smaller pieces; and (2) distribution and age of sutures, accretionary prisms, volcanic arcs, fold-thrust belts, stretched continental crust, strike-slip faults, and ocean floor. We conclude that: (1) North America and Eurasia remained relatively fixed to each other until the latest Cretaceous-Paleocene opening of the Labrador Sea–Baffin Bay and Greenland-Norwegian and Eurasian basins (earlier convergence between North America and Eurasia in the Bering Sea region shown on many reconstructions are artifacts of incorrect plate reconstructions); (2) the North Slope–Seward-Chukotka block has constituted an isthmus connection between North America and northeast Asia since at least the middle Paleozoic and did not rotate away from the Canadian Arctic; (3) the Canada basin opened behind a clockwise-rotating Alpha Cordillera-Mendeleyev ridge arc during the Early to middle Cretaceous and consumed older, Paleozoic(?) Makarov basin ocean floor (the Chukchi cap is a detached continental fragment derived from the Beaufort Sea; the North Slope Arctic margin is a left-lateral transform fault associated with the opening of the Canada basin); and (4) the Nares Strait fault has a net relative displacement of approximately 25 km, but actual motion between

Greenland and northern Ellesmere was about 250 km of strongly transpressive motion that resulted in the Eurekan and Svalbardian orogenies.

RU, KE, Nan Hai West Oil Co., Zhangjiang, Guangdong, People's Republic of China, and JOHN D. PIGOTT, Univ. Oklahoma, Norman, OK

South China Sea Tectonic Evolution and Hydrocarbon Potential: New Geological and Geophysical Constraints

The South China Sea has yet to receive a DSDP investigation; consequently, numerous untested models have been proposed for its post-Cretaceous evolution. From a compilation of regional oceanographic heat-flow measurements with offshore and onshore bore-hole temperatures, we thermally model and constrain possible interpretations of its tectonic evolutionary path. The heat-flow data, together with magnetic profiles, depth to basement determinations, and regional sediment isopachs, characterize two principal subbasin extensional elements—one trending east-west (northern area between Hainan and Luzon) and the other trending northeast-southwest (central area between Palawan and Vietnam). The thermal models of simple lithosphere cooling suggest the central area began spreading 55–58 m.y.B.P. and the northern region 34–35 m.y.B.P. These dates of incipient extension correspond to two principal unconformities in Paleocene and Oligocene strata of both offshore China and Palawan, and together they indicate regional uplift of the South China Sea owing to thermal expansion prior to the spreading events. The thermal models also suggest that in the northern region, spreading ceased approximately 19 m.y.B.P., which compares favorably with published magnetic estimates of 17.7 m.y.B.P. A late Cenozoic heating event is evidenced by a thermal anomaly in the southern portion of the central region (southern Vietnam margin) that may be related to incipient spreading along a zone of crustal weakness inherited from the Jurassic-Cretaceous Sunda-Tethys suture. Overall, these data tend to support the hypothesis of spreading occurring first in the central region and then in the northern region.

From the geophysical data and observations of Cretaceous ophiolites cropping out to the south in Sabah and Brunei, we kinematically border our model to the south and propose the Palawan Ulugan fault to be a right-lateral suture between continental and intermediate crust. Structurally, within limits of the data presently available, our model further predicts the southern China shelf to have experienced two principal episodes of extension with a net result of younger (< 34 m.y.B.P.) east-west trending graben normal faults superimposed upon older (< 55 m.y.B.P.) northeast-striking pregraben normal faults. In terms of thermal maturity, geochemical kinetic modeling of Late Cretaceous source rocks suggests depths to the oil ceiling to range from 1.3 to 1.8 km in the northern region and from 0.98 to 1.6 km in the central region. Similarly, depths to the oil floor are estimated to range from 2.4 to 3.4 km and from 1.8 to 3.2 km in the two respective regions.

RUPPEL, STEPHEN C., Bur. Econ. Geology, Austin, TX

Facies and Geochemical Characterization of Mississippian Rocks in Palo Duro and Hardeman Basins, Texas

Mississippian rocks in the southern Texas Panhandle constitute a complex sequence of carbonate deposits formed in a platform-to-basin setting. Following relatively rapid transgression and inundation of the area from the north and east, the Hardeman basin area was characterized by outer platform conditions in which isolated carbonate buildups developed surrounded by relatively deep water. The Palo Duro basin to the west was the site of shallow-water, inner platform deposition. In intermediate areas, limestone turbidites, perhaps derived from carbonate buildups to the east, accumulated in quiet water. After this initial transgression, an upward-shallowing trend resulted in the formation of ooid to skeletal shoals throughout the area.

Although current production in the area is coincident with the distribution of organic-rich Upper Mississippian shales in the eastern part of the Hardeman basin, TOC studies indicate that potential carbonate source rocks are present in the western Hardeman and eastern Palo Duro basins. Mississippian rocks in the Palo Duro basin proper have little source rock potential. Vitrinite reflectance studies indicate that Hardeman basin rocks are well within the oil window. However, correlative deposits at

equivalent depths in most of the Palo Duro basin are only marginally mature.

Although thermal maturity seems to be mirrored by the present geothermal gradient, and source rock quality appears related to depositional setting (depth of water), successful exploration outside currently productive areas will require a detailed analysis of organic geochemistry and depositional facies.

RUZYLA, KENNETH, Exxon Production Research Co., Houston, TX

Pore Size Distributions by Analysis of Back-Scattered Electron and Fluorescence Images

Computerized image analysis provides direct, rapid, and highly accurate measurement of pore size in thin or plane sections. The method can be applied to a wide range of rock types and requires only conventional sample-preparation techniques. Back-scattered electron or fluorescence microscopy images of impregnated samples are scanned, converted to digital form, and stored and processed on a microcomputer. Individual picture points from a matrix of up to 600,000 elements per scan are classified as rock or pore based on gray level. Pore size is obtained by area measurements of individual pores and by Feret's diameter, the maximum spacing between parallel tangents to a pore in up to 56 directions. The measurements are readily summarized as pore size distributions.

Cumulative porosity vs. pore diameter crossplots indicate how porosity is distributed and can be used to aid reservoir evaluation and production assessment. The crossplots can also be used to determine the amount of porosity contributed by different pore types. Individual pores, or groups of pores classified by size or shape, can be interactively identified on the image analyzer monitor, enabling the user to make a visual association of pore type with size. Rocks having a wide range of pore size can be analyzed at more than one level of magnification, and the data can be merged to form a composite pore size distribution for the sample.

Pore size distributions of carbonate rocks containing complex pore systems show changes in slope that are indicative of pore type. Other carbonates, including fine crystalline dolomites and microcrystalline limestones, show relatively uniform pore size distributions, which reflect a single pore type.

SAID, F. M., National Oil Co., Benghazi, Libya, and W. H. KANES, Univ. South Carolina, Columbia, SC

Paleozoic Producing Sequences in Ghadames Basin of Libya, Tunisia, and Algeria: New Stratigraphic Concepts for Hydrocarbon Exploration

The Ghadames basin covers an area of more than 3 million km² in Libya, Tunisia, and Algeria. Although the basin has been productive since the mid-1950s, recent construction of a pipeline in western Libya has spurred additional exploration. The results of this increased drilling activity and the incorporation of earlier work have provided new data that indicate the following. (1) The clastic sequence underlying the Tanezuft shales has porous and permeable fairways that may be related to the Hogar glaciation to the south. (2) The clastic sequence overlying the Tanezuft shales was deposited as a northerly sourced fluvial-deltaic clastic wedge, which thins and becomes less porous to the south. Only in the area of the Gargaf arch does this sequence become cleaner and more porous; there, a shoreline-sand depositional sequence dominates. An easterly source of the clastics is indicated.

With the lack of large structures in the basin, the facies model presented here for the depositional history provides indications of more productive porous and permeable fairways.

SALLER, ARTHUR H., Cities Service Oil and Gas Corp., Tulsa, OK

Marine Diagenesis on Enewetak Atoll

Marine waters currently circulate through Eocene and Miocene limestones deposited in slope, fore-reef, and reef-related environments on Enewetak Atoll. Subsurface core samples (375-1,400 m deep) demonstrate that marine waters have extensively altered the original carbonate sediments. Evidence supporting alteration by marine water includes: (1) calcite cements with isotopic compositions characteristic of a marine ori-

gin ($\delta^{13}\text{C} = 1.3$ to 2.5 ‰; $\delta^{18}\text{O} = -1.8$ to 0.4 ‰ PDB), (2) dolomites with average isotopic compositions that are in equilibrium with cold normal-marine waters ($\delta^{13}\text{C} = 2.3$ ‰; $\delta^{18}\text{O} = 2.5$ ‰ PDB), (3) consistent magnesium concentrations in fossil coralline algae, and (4) a lack of diagnostic evidence for meteoric diagenesis. Products of marine diagenesis on Enewetak Atoll are dependent on the CaCO_3 saturation state of the diagenetic waters. Above the aragonite saturation depth (350 m), marine diagenesis is dominated by aragonite and high-magnesian-calcite cementation. Between aragonite and calcite saturation depths, marine waters apparently have dissolved aragonite and precipitated low-magnesian, radial calcite cements. Below the calcite saturation depth (1,000 m), deep marine waters have apparently dissolved some calcite and have precipitated dolomite. Effective marine diagenesis requires limestones with good permeability and a hydraulic drive, such as tidal pumping or thermal convection, to move large volumes of marine water through the rock.

SAPPINGTON, JOHN M., Elan Exploration Inc., Mt. Pleasant, MI, and F. B. DAVIES, Aangstrom Precision Corp., Mt. Pleasant, MI

Computer Mapping of Silurian Pinnacle Reefs in Northern Michigan

Current availability of high-quality well completion records accessed via computer data bases in Michigan has allowed further exploitation of the northern Michigan reef trend. Integration of these data with computer mapping enables investigation of large regions in terms of structural and stratigraphic relationships. Inferiority of subsea structure maps is clearly demonstrated because of contamination by irrecoverable surface elevation errors. Use of thickness maps of reef units and Middle Silurian salt deposits has resulted in delineation of new regional reef patterns and an increased resolution of local reef development. Case studies from the Michigan reef trend demonstrate the use of "play maps," where computer-generated maps are constructed in yearly intervals to test exploration strategies for less developed regions.

SAVIN, SAMUEL M., MINGCHOU LEE, and JAMES L. ARONSON, Case Western Reserve Univ., Cleveland, OH

Thermal History of Sandstones and Shales: Oxygen Isotope and K/Ar Evidence

The combined use of oxygen isotope and K/Ar dating can yield information about the diagenetic history of clay minerals in shales and sandstones. In the Tertiary shale sequences of the United States Gulf Coast, in which the dominant detrital clay mineral is mixed-layer illite/smectite, the progressive conversion of expandable clay layers to illite layers can be monitored by the isotope systematics. Oxygen isotopes of fine-grained clay and quartz approach equilibrium with one another, raising the (as yet unrealized) possibility of an O-isotope geothermometer. Fine-grained quartz becomes isotopically zoned as detrital grains are overgrown by diagenetic quartz that forms, accompanying the smectite-to-illite transformation. The K/Ar clock of the diagenetically formed illite layers is set to zero age at the time of diagenesis (although that of existing illite layers within the crystals remains unaffected), and it is therefore possible to estimate the mean time of diagenesis in such shale sequences.

Illite is a common cement in sandstones. The time of cementation by illite can be estimated from K/Ar systematics in cases, such as that of the Permian Rotliegendes of the North Sea, where the clay-sized fraction of the original detritus was relatively free of illite or other K-bearing phases. Conditions of cementation can be inferred from oxygen isotope measurements, augmented by knowledge of the geologic/tectonic history of the sandstone. In the case of the Rotliegendes, the timing and conditions of illite cementation were relatively uniform within fault blocks but varied from block to block. Oxygen isotope measurements indicate that meteoric water components were important in illite formation at times when nearby sections of the Rotliegendes were exposed to the surface by uplift and erosion.

SAWYER, D. S., Y. NAKAMURA, W. O'BRIEN, and J. EBENIRO, Univ. Texas at Austin, Austin, TX

Crust Type and Structure, Northern Gulf of Mexico: an Ocean Bottom Seismograph-Air Gun Seismic Transect