In November to December 1983, the University of Texas and the U.S. Geological Survey conducted a seismic experiment that attempted to measure crustal and upper mantle depths and velocities along a transect of the Gulf of Mexico south of Galveston, Texas. The transect is composed of 5 along-strike lines spaced to span the continental shelf and slope and reaching the deep basin. Two 2,000-in.³, 2,000-psi air guns were fired simultaneously at 30-sec intervals at 5 knots for a shot spacing of 77 m. The signals were recorded by digital ocean-bottom seismographs with vertical geophones. Four seismographs were placed along each 90-km line. The seismic sections obtained are densely sampled and fully reversed, about half showing seismic arrivals at the full range of the line.

We find that the deep Gulf of Mexico is, as expected, underlain by oceanic crust. On the outer slope, we see deeply penetrating arrivals from below thick salt, and we find that this crust is thicker than the oceanic crust. North of the thick salt, we see the crust thinning to nearly the thickness of oceanic crust. Although this crust may be seismically indistinguishable from oceanic crust, we believe it to be highly extended continental crust. We interpret the two northern lines to show northward thickening, extended continental crust.

SCHAFER, CHARLES T., Geol. Survey of Canada, Bedford Inst. of Oceanography, Dartmouth, Nova Scotia, Canada, and DAVID B. SCOTT, Dalhousie Univ., Halifax, Nova Scotia, Canada

Micropaleontology, Textural and Isotopic Characteristics of Turbidites on Continental Slope and Rise off Newfoundland

Graded turbidite deposits observed in x-radiographs of gravity and piston cores collected from the lower slope and rise (2,300-3,210 m) occur below a well-documented 9,300 y.B.P. ash zone. The largest turbidite observed on the upper rise is 35-cm thick. C¹⁴ dates of foraminifera tests from intervals above and below this deposit average about 13,000 y.B.P. suggesting deposition during late glacial time. The base of this deposit is marked by a textural change to comparatively coarse sediments. It is also marked by an anomalous concentration of foraminiferal species (up to 50%) belonging to the miliolid family, that constitutes 2-4% of modern assemblages. A pronounced increase in the percentage of *Valvulinaria artica* in the turbidite suggests a middle-slope source. The thinner gravity flows (10-15 cm thick) that occur in the deeper sections of the upper-rise cores have extrapolated C¹⁴ ages of less than 22,000 y.B.P. Their foraminiferal assemblages suggest outer-shelf to upper-slope sediment sources.

A comparatively thick (150 cm), graded turbidite sequence of unknown source occurs on the lower slope, just above ash zone 2 (approximate age is 64,000 y.B.P.). The related turbidity-current event apparently took place during the latter part of isotope stage 4. Texturally, this sequence is similar to the 35-cm thick upper-rise turbidite, but it lacks the anomalous concentrations of foraminiferal species in its basal interval. Oxygen isotope values (sinistral *N. pachyderma*) are relatively homogeneous throughout the sequence at a value of +3.75 ppt compared to +2.75 ppt for the upper Holocene section of the core.

SCHAMEL, STEVEN, Univ. South Carolina, Columbia, SC, HABIB LAZREG, ETAP, Tunis, Tunisia, and KENNETH CIRIACKS, Amoco International, Houston, TX

Tectonic Evolution and Hydrocarbon Prospects of Tunisia-Sicily Shelf

The stratigraphic evolution, structural styles, and principal hydrocarbon prospects of the Tunisia-Sicily shelf are all linked to the crustal "template" created during the middle Mesozoic rifting of the Tethyan margin of North Africa. Transtensional stretching and fragmentation of the Tunisia-Sicily shelf during the Late Triassic to Jurassic at the junction of the south Sahara and Gilbraltar fracture zones created a complex array of ridges and furrows and localized pull-apart basins. This block-faulted shelf region was buried during the Cretaceous and early Cenozoic beneath a cover stratigraphy that ranged widely from a thin pelagic limestone succession devoid of terrigenous components in Sicily to a considerably thicker neritic Tunisian succession containing terrigenous and carbonate rocks. Differences in stratigraphic character across the shelf relate to the relative position of Sicily and Tunisia between the unstable and subsident Tethyan margin and the stable and emergent Saharan platform. As the North African continental margin subducted northward beneath the accretionary Kabyle-Calabrian belt in the late Cenozoic, the

thin carbonate successions of Sicily responded by detaching in a series of southward-migrating thrust sheets. Along strike in Tunisia, the thrust sheets die out and stress was taken up by folding the thick marly succession weakened by numerous Late Triassic evaporite diapirs. Despite differences in structural style, the orientations of the structures in both Tunisia and Sicily have been guided by the crustal template created by middle Mesozoic rifting.

Hydrocarbons on the Tunisian-Sicily shelf are related to (1) Mesozoic or younger structures that deform the early Paleozoic black shaleorthoquartzite succession of the Saharan platform, (2) synrift anoxic basin facies, (3) the Cretaceous-early Cenozoic shelf margin in the Tunisian section, and (4) late Cenozoic synorogenic clastic facies.

SCHUEPBACH, MARTIN, Aminoil Inc., Houston, TX, STEVEN SCHAMEL and STEPHEN K. PERRY, Univ. South Carolina, Columbia, SC, and DONALD U. WISE, Univ. Massachusetts, Amherst, MA

Structural Style, Stratigraphic Variability, and Hydrocarbon Traps in Gulf of Suez

Hydrocarbon traps in the Gulf of Suez, whether of a structural or a stratigraphic type, fundamentally are all products of the highly complex late Cenozoic fault-block mosaic. Understanding the geometry and the evolution of this tilt-block mosaic is the key to further successful exploration in the Gulf.

The major faults forming the Gulf of Suez basin are listric normal faults broken along strike by synchronous cross faults that serve as transform or "transfer" structural elements. The cross faults are a consequence of the inability to form uniform and laterally continuous listric fault surfaces within the mechanically anisotropic and inhomogeneous crystalline basement terrane of the Gulf. The combination of contemporaneous Gulf-parallel listric faults and cross-strike "transfer" faults gives rise to the complex mosaic of polygonal tilt blocks. Vertical variations in the structural behavior result from a synrift stratigraphy dominated by thick marls and evaporites. The ductile synrift rocks exhibit shallow secondary detachments, drape folds, and compaction and/or diapiric structures that are decoupled from, but appear controlled by, the brittle underlying basement-rooted tilt-block mosaic.

Lateral stratigraphic variability observed in the Gulf is the consequence of sea level rising and falling relative to the higher structural elements in the tilt-block mosaic. This has resulted in deeply eroded basement highs, clastic apron and perched reef reservoirs associated with the highs, and thick organic-rich marl accumulations in the intervening lows that serve as "kitchen" areas.

SEGAL, D. B., M. D. RUTH, and I. S. MERIN, Earth Satellite Corp., USA, Chevy Chase, MD, H. WATANABE and K. SODA, JAPEX Geoscience Inst., Tokyo, Japan, and O. TAKANO and M. SANO, ERS-DAC, Tokyo, Japan

Remote Detection of Anomalous Mineralogy Associated with Hydrocarbon Production, Lisbon Valley, Utah

Diagenetic mineral assemblages within the Wingate formation are closely associated with hydrocarbon production at Lisbon Valley, Utah. The Wingate formation, exposed along the southwestern flank of the anticline, has a relatively uniform composition and appearance over the entire Colorado Plateau, except at isolated localities such as Lisbon Valley, where it is locally bleached. Previous workers have suggested that hydrocarbon microseepage may account for bleaching of the Wingate Sandstone and the presence of uranium mineralization in outcrops overlying the reservoir at Lisbon Valley. Using broad-band, Landsat Multispectral Scanner (MSS) and airborne Thematic Mapper Simulator (TMS) data, the bleached facies was mapped on the basis of brightness and lack of ferric iron. Examination of the TMS data provided further discrimination of bleached facies because relative abundances of clay minerals are detectable with this sensor.

Analysis of high-resolution airborne spectroradiometric data, thin sections, and x-ray diffraction suggests that the bleached rocks overlying the reservoir at Lisbon Valley contain abundant kaolinite and minor amounts of feldspar. Unbleached exposures contain substantially less clay and abundant feldspar.

This study shows a correlation between the abundance of clay minerals, particularly kaolinite, and hydrocarbon production at Lisbon Valley. Because one of the principal differences between the bleached and unbleached rocks is in the relative abundance of clay minerals (and the TMS is very sensitive to clays), areas of potential hydrocarbon-induced diagenetic alteration may be mapped using broad-bank sensors.

SELF, GREGORY A., S. Q. BREARD, HOWARD P. RAEL, JEFFREY A. STEIN, and MARTIN O. TRAUGOTT, Amoco Production Co., New Orleans, LA, and WILLIAM D. EASOM, Callon Petroleum, Natchez, MS

Lockhart Crossing Field: New Wilcox Trend in Southeastern Louisiana

A 1982 Wilcox oil discovery in southeastern Louisiana constituted one of the more significant onshore United States discoveries for the year and illuminated a new oil trend. Prior to this discovery, Lockhart Crossing field was known for its gas and condensate production from Cretaceous lower Tuscaloosa sandstones at depths of 17,000-18,000 ft (5,200-5,500 m). Oil production from sandstones of the lower Eocene of the uppermost Wilcox Group was recently established at 10,000 ft (3,000 m) with successful completion (630 BOPD), of the Callon 1 Reed Erickson. To date, 30 producers and 5 dry holes have been drilled in the field, with production peaking at 10,000 BOPD and recoverable reserves of 15-20 million bl of oil after secondary recovery.

The main field reservoir is a 40-80 ft (12-24 m) marine sandstone with two distinct facies present. The most dominant of these two facies is a coarsening-upward sequence of very fine to fine-grained glauconitic sand deposited as a nearshore bar. The second is a tidal channel facies with a fining-upward sequence of medium to very fine grained sand. The spatial relationship of these two facies defines a progradational sequence of tidal channels overstepping and eroding into existing nearshore bars. The primary trapping mechanism, however, is structural in the form of a rollover anticline.

In April 1983, Callon and Amoco extended the trend southeast with the discovery of Livingston field. Chances are good that this new trend will yield additional discoveries with continued exploration.

SELF, GREGORY A., Amoco Production Co., New Orleans, LA, and ROBERT W. PIERCE, Eastern New Mexico Univ., Portales, NM

Growth Movement of Citronelle Dome and Its Genetic Relationship to Other Salt Structures in Southern Mississippi Salt Basin: an Approach in Computer Applications

Citronelle field, one of Alabama's largest oil fields, was created by one of the many salt structures that occur throughout the southern Mississippi salt basin.

The history of domal movement was determined using computer techniques of conventional mapping and trend-surface analysis based on structural tops and thicknesses of 22 lithostratigraphic field units. Unique statistical parameters for each trend surface were then compared using cluster analysis. The definition of three major episodes of domal movement resulted.

The first episode occurred in the Cretaceous during Tuscaloosa deposition. The second occurred in the Paleocene during Midway deposition, and the third occurred in middle Eocene during Claiborne deposition. A similar history of structural development was determined for two other salt structures along the southern margin of the southern Mississippi salt basin.

This genetic relationship appears to reflect subsidence characteristics within the southern Mississippi salt basin. As such, this relationship may be valuable in better understanding the occurrence of oil and gas reserves in the basin and in future exploration efforts in this mature area.

SEYLER, BEVERLY, Illinois State Geol. Survey, Champaign, IL

Petroleum Reservoirs in Silurian Dolomite of Western Illinois

The basal Silurian Alexandrian carbonates, sandwiched between the Ordovician-Silurian and Silurian-Devonian unconformities, have produced more than 3 million bbl of oil from the Buckhorn East and Brooklyn fields on the western margin of Illinois. These fields are shallow (450-670 ft), remote from the most-productive portion of the Illinois basin, and potential exists for additional production from this unconformity play. Three lithofacies were recognized in Alexandrian carbonates in detailed core and wireline studies that included thin-section, x-ray, and SEM analyses. These carbonates are regressive sequences composed of shallow-shelf, restricted-subtidal, and intertidal facies. The shallowshelf facies is primarily biosparite composed of crinoid and brachiopod debris; this facies is dolomitic in places. The restricted-subtidal facies consists of both bioturbated and undisturbed gray micrites containing stylolites, green chloritic clay, and glauconite; sparry clacite fills vertical cracks. The intertidal facies is tan, mottled, vuggy dolomite composed of algal mats, anhydrite-filled birdseye structures, burrows, and scattered marine fossils.

Petroleum reservoirs apparently developed as a result of dolomitization in intertidal and shallow-shelf carbonates. Vuggy porosity, a possible result of karst leaching, is found in pay zones within the dolomitic intertidal facies. Evidence of emergence and subaerial weathering include the presence of vadose silt, anhydrite, and nontectonic breccia.

SHAFFER, FRANK R., Shaffoil & Associates, Inc., Harahan, LA, and ALLEN LOWRIE, U.S. Naval Oceanographic Office, Picayune, MS

Glacial Cycles, Hydrocarbon Generation and Salt Tectonics on Louisiana Slope

In the northern Gulf of Mexico, maximum growth on regional downto-the-south faults, maximum salt tectonics, and the timing of maximum oil generation are synchronous. Sedimentary pulses triggered by glacial cycles are recognized to correlate well with the maxima. Synchrony of glacial, structural, and sedimentary events suggest a cause and effect relationship.

During major glacial events, up to an 8-fold increase in melt water had an order of magnitude increase in the amount of sediment brought to the northern Gulf of Mexico. Eight discrete upper Pleistocene sediment packages have been identified in seismic profiles. Likewise, approximately 8 major growth fault trends are recognized on the Louisiana slope. The causal effect of increased lithostatic pressure owing to rapid deposition on underlying plastic salt promotes flowage that activates regional faults and accelerates salt tectonics. Higher biological productivity during glacial maximas and contemporaneous regressive sands explain higher concentrations of hydrocarbons during maximum structural and stratigraphic activity.

SHANMUGAM, GANAPATHY, JOHN E. DAMUTH, and RICH-ARD J. MOIOLA, Mobil Research and Development Corp., Dallas, TX

Is Turbidite Facies Association Scheme Valid for Interpreting Ancient Submarine-Fan Environments?

Although turbidite facies reflect only their processes of deposition, turbidite facies associations are routinely used to identify ancient submarinefan systems and their subenvironments (e.g., upper fan, channel, etc). This credulous leap from interpreting process of deposition to interpreting environment of deposition may not be valid for the following reasons: (1) Mutti and Ricci Lucchi's facies association scheme was developed exclusively from the study of ancient turbidite sequences; however, the true relationship between turbidite facies associations and related environments has not been adequately documented in modern settings; (2) channel-levee complexes of many modern submarine fans such as those of the Amazon and Mississippi are large enough to contain most ancient fan systems in their entirety; and (3) comparable channel-levee complexes have not been recognized in outcrop. Consequently, the validity of turbidite facies associations for interpreting ancient submarine-fan systems and their subenvironments must be considered tenuous until confirmed by detailed core studies of a variety of modern fans.

SHEPHERD, R. G., Univ. Kentucky, Lexington, KY

Anisotropic Permeability, Eolian Lyons Sandstone

Directional permeabilities from nine attitude-oriented core samples of well-laminated eolian Lyons Sandstone from Larimer County, Colorado, were measured with a gas permeameter. Intrinsic permeabilities across the laminae, parallel to lamination and dip, and parallel to lamination