Berriasian-Cenomanian Carbonate Platform Successions of the Southwestern U.S. and Northern Mexico

Charles Kerans¹, Robert Loucks², Lowell Waite³, Ryan Phelps¹, and Laura Zahm²

¹Bureau of Economic Geology, The University of Texas at Austin, University Station, Box X, Austin, Texas 78713-8924

²Jackson School of Geosciences, The University of Texas at Austin, 1 University Station C1100, Austin, Texas 78712-0254

³Jackson Pioneer Natural Resources Company, 5205 N. O'Connor Blvd., Ste. 200, Irving, Texas 75039

ABSTRACT

The Cretaceous carbonate platform of the southwestern U.S. and northern Mexico illustrates a spectrum of shelf-slope morphologies, facies architectures, and reservoir styles that reflect the changing structural underpinnings, subsidence patterns, and sediment supply variables in each region. Five 2nd-order sequences are tracked between the Bisbee Basin-Chihuahua Trough, western Comanche Shelf/Maverick Basin, and Llano Uplift to Stuart City Margin. The 2nd-order supersequences (SS) are (1) the Berriasian– Early Aptian Hosston/Sligo-Cupido, (2) the mid Aptian–Early Albian Pine Island/La Pena/Pearsall–Cow Creek/James/Lower Mural, (3) Early-Mid Albian Glen Rose/Mural, (4) Late Albian Edwards/Georgetown–Stuart City, and (5) Early-Mid Cenomanian Del Rio–Buda.

The Llano-Stuart City passive margin succession illustrates all five 2nd-order supersequences. SS1 (Sligo) establishes a continuous reef-rimmed platform margin, followed by Early Aptian eustatic rise and deposition of the Early Aptian Pearsall–La Pena supersequence 2. Sequence 3 Early-Mid Albian Glen Rose carbonates prograde dramatically, reclaiming the sequence 1 margin. Sequence 4 once again is dominated by aggradational stacking and development of a well-defined reefal rim/lagoon/shallow shelf with reservoirs developed in rim grainstones. Sequence 5 records the demise of the platform terminated by the mid-Cenomanian sea level fall. The western Comanche Shelf/Maverick Basin is best known for productive SS3 Glen Rose transgressive sequence set (TSS) buildups and highly progradational SS4 grainstones. The Mural shelf of the Bisbee Basin/Chihuahua Trough is dominated by siliciclastics. The Early Aptian SS2 eustatic rise initiates carbonate sedimentation in the Lower Mural. SS3 Upper Mural includes TSS patch reefs and platform-margin reefs. SS4 highstand sequence set (HSS) deposits are forced regressive intercalated highstand grainstones and lowstand siliciclastic wedges.

Kerans, C., R. Loucks, L. Waite, R. Phelps, and L. Zahm, 2010, Berriasian-Cenomanian carbonate platform successions of the southwestern U.S. and northern Mexico: Gulf Coast Association of Geological Societies Transactions, v. 60, p. 807.