

## TECTONIC DEVELOPMENT OF IDAHO- WYOMING THRUST BELT

(R. M. A. G. Evening Meeting, October 22, 1964)

FRANK C. ARMSTRONG and STEVEN S. ORIEL\*  
U. S. Geological Survey, Denver, Colorado

**ABSTRACT:** Three stages are evident in the tectonic development of southeastern Idaho and western Wyoming. First, the changing patterns of tectonic elements during deposition, second, development of northward-trending folds and thrust faults; and third, development of block faults that produced horst ranges and graben valleys.

During Paleozoic time about 50,000 feet of marine sediments, mostly limestone and dolomite, were deposited in a miogeosyncline and about 6,000 feet of mixed marine sediments were deposited on the shelf to the east. Detritus came from both east and west recurrently from Cambrian time on. Starting in Mississippian time, the belt between shelf and miogeosyncline, where thicknesses increase markedly, shifted progressively eastward.

During Mesozoic time about 35,000 feet of marine and continental sediments were deposited in the western part of the region and about 15,000 in the eastern, terrestrial deposits becoming increasingly dominant. Western positive areas became the chief source of detritus. The belt of maximum deposition shifted progressively eastward; maximum thicknesses of succeeding geologic systems are not superposed. In Late Triassic a belt to the west rose and the miogeosyncline started to break up. As Mesozoic time progressed the western high spread eastward, until by the end of the Jurassic the miogeosyncline gave way to intracratonic geosynclinal basins that received thick deposits, particularly in Cretaceous time. Cenozoic sedimentary rocks are products of orogeny in the region.

The second stage, which overlapped the first, produced folds overturned to the east and thrust faults dipping gently west in a zone, convex to the east, 200 miles long and 60 miles wide. Stratigraphic throw on many larger faults is about 20,000 feet; horizontal displacement is at least 10 to 15 miles. Lack of metamorphism and mylonite along the faults is striking. From west to east, the thrust faults cut progressively younger beds, have progressively younger rocks in their upper plates, and are estimated to be successively younger. Thrusting started in the west in latest Jurassic and ended in the east perhaps as late as Early Eocene time; detritus shed from emergent upper plates is preserved in coarse terrestrial strata of corresponding ages.


West of the thrust belt is a northwestward-trending area underlain mostly by lower Paleozoic rocks and flanked on east and west by upper Paleozoic and Mesozoic rocks. Scattered pieces of eastward-dipping thrust faults have been reported west of the older rocks. This central area of old rocks has been interpreted as (1) part of a large continuous thrust sheet moved scores of miles from the west, or (2) an uplifted segment of the earth's crust from which thrust sheets to the east and west were derived. Both interpretations have defects; relative thrust ages are difficult to explain under the first; a large positive gravity anomaly, expectable under the second is apparently absent

\* Speaker

Block faulting, the third stage of tectonic development, started in Eocene time. Faulting has continued to the Recent, as indicated by broken alluvial fans, displaced basalt flows less than 27,000 years old, and earthquakes. North-trending and east-trending fault sets are recognized. Old, east-trending steep faults in the Bear River Range may be tear faults genetically related to thrusting. Movement along many faults has been recurrent. Patches of coarse Tertiary gravel on the flanks and crests of ranges, for which there is no provenance with present topography, may record reversed vertical movement along some north-trending faults. Present topographic relief of basins and ranges is tectonic.

<p style="text-align: center;"><b>LEON J. PANKAU</b>  <b>CONSULTING PHOTOGEOLOGIST</b>  <b>STRUCTURAL GEOMORPHOLOGIST</b>  <i>Specialist in interpreting and mapping</i>  <i>obscure surface Geology</i></p> <p>PHONE: <b>466-4726</b>                      <b>3980 W. 134TH PL.</b>  <b>BROOMFIELD, COLO.</b></p>	<p style="text-align: center;"><b>JOHN C. OSMOND</b>  <b>CONSULTING GEOLOGIST</b></p> <p style="text-align: center;">REGIONAL STUDIES, PROSPECT EVALUATIONS  EXPLORATION MANAGEMENT  OIL, GAS AND SOLID HYDROCARBONS</p> <p>P. O. BOX 8152, FOOTHILL STATION                      PHONE 326-0531  SALT LAKE CITY, UTAH 84108                      (AC-801)</p>
<p style="text-align: center;"><b>HAROLD H. BROWN</b>  <i>Consultant in Petroleum Geology</i></p> <p>PHONE: 325-2188                      •                      2705 MESA DRIVE  FARMINGTON, NEW MEXICO</p>	<p style="text-align: center;"><b>TROLLINGER-GOSNEY</b>  <i>and</i>  <i>Associates</i></p> <hr style="width: 20%; margin: auto;"/> <p style="text-align: right;">70 W 6th Ave.  Denver 4, Colorado  Phone 255-0042</p> <p><i>Quality Photogeology—</i>  <i>Geomorphic Analysis</i></p>

# HOW TO S-T-R-E-T-C-H your exploration dollar



**GEOPHOTO** has available for immediate delivery detailed photogeologic maps in important petroleum provinces including Rocky Mountains, Sonora-Las Cruces Basins, Southeast New Mexico, Delaware Basin, Marfa Basin, Val Verde Basin, Kerr Basin, Arkoma Basin and Appalachian Area. Geology is ink drafted to copyrighted Geophoto planimetric bases compiled from township plats. You may inspect any of these maps. Price quotations gladly provided. Discounts for quantity purchases.

Investigate These Other GEOPHOTO Services • Quantitative Photogeology  
• Structural Contouring • Topographic Mapping • Detailed Stratigraphic Studies • Contract Surface Parties • Comprehensive Geomorphic Studies



## **GEOPHOTO SERVICES INC.**

BOX 22293 • 2149 SOUTH HOLLY ST. • DENVER, COLO., 80222 • PHONE 303/757-4941